



Bladder Function as a Representation of Systemic Health: An Integrative Framework for Understanding Lower Urinary Tract Symptoms

Jin Wook Kim^{1,2}

 <https://orcid.org/0000-0003-4157-9365>

¹Department of Medical Informatics, Chung-Ang University, Seoul, Korea

²Department of Urology, Chung-Ang University Gwangmyeong Hospital, Gwangmyeong, Korea

Email: jinwook@cau.ac.kr

In the broad overview of the physiological function of the human body, the bladder seems like an addendum, an anatomical appendage posited at the physiological end of the excretory branch. Though traditional methodologies constantly revisited urine as a medium to detect systemic changes, the complication of a reservoir which diffused the physiologic signal resolution over time diluted the utility vis-à-vis serologic or even salivary based analyses. In this context, understanding bladder pathophysiology itself as a reflection of underlying systemic changes represents an underexplored paradigm.

Circadian rhythm disruption represents a major external environmental influence on internal systemic change, and nocturia emerges as a reciprocal response to this environmental perturbation [1, 2]. Shift workers on alternating schedules demonstrate increased nocturia primarily through decreased nocturnal bladder capacity rather than nocturnal polyuria, implicating central circadian machinery over simple volume overproduction [2]. The NHANES (National Health and Nutrition Examination Survey)-based study in this issue extends this framework, demonstrating that nocturia independently predicts all-cause and cardiovascular mortality, with serum uric acid acting as a partial mediator [3]. The J-shaped curve relating uric acid to mortality risk indicates that physiological balance, not suppression of any single metabolite, remains the therapeutic imperative.

The proteomic investigation of urothelium in obstruction-re-

lated overactive bladder (OAB) moves us from population associations to molecular mechanisms [4]. Analyzing tissue from men undergoing transurethral resection, investigators identified numerous differentially expressed proteins between OAB and non-OAB phenotypes. The pathways implicated extend far beyond the bladder wall itself. Cytoskeletal proteins — ACTA2, CFL2, MYLK, PPP1R12B — showed consistent downregulation, suggesting structural reorganization at the cellular level. Serotonergic signaling components were disrupted, with GNAO1 suppressed and GNA13 elevated, pointing toward aberrant neurotransmitter processing. Immune pathway activation, particularly neutrophil degranulation markers, indicates inflammatory involvement. The urgency these patients experience is not merely a symptom; it reflects tissue-level transformation involving architecture, neural signaling, and immunity simultaneously.

The question of cognitive safety with benign prostatic hyperplasia (BPH) medications has troubled prescribers for years. Alpha-blockers cross the blood-brain barrier, and 5-alpha reductase inhibitors alter neuroactive steroid metabolism. Animal studies demonstrate memory impairment with tamsulosin [5, 6]. The network meta-analysis pooling millions of patients provides reassurance, at least provisionally [7]. Neither drug class demonstrated significant dementia risk elevation compared to untreated controls. Yet the SUCRA (surface under the cumulative ranking curve) rankings deserve attention: finasteride and



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

dutasteride ranked highest for dementia probability, over alpha-blockers, reflecting the systemic involvement of decreased androgen signals influencing systemic senility. The absence of statistical significance does not equal absence of effect, particularly for outcomes requiring decades to manifest.

Younger patients showed substantially better recovery odds. Male sex paradoxically favored recovery — a finding explained by etiology rather than anatomy, since female retention typically reflects complex pelvic floor pathology rather than simple prostatic obstruction [8]. Postoperative retention resolved more readily than medical causes. Larger retention volumes compromised recovery, likely through ischemic detrusor injury that impairs contractility [9]. This volume threshold carries immediate practical implications: early recognition and decompression matter.

Viewed together, these studies present a coherent if challenging picture. The bladder operates within interconnected physiological networks — metabolic pathways governing uric acid handling, neurotransmitter systems modulating both detrusor function and central cognition, inflammatory cascades triggered by mechanical obstruction, and regenerative capacity determining recovery from acute insults. Bladder symptoms may therefore serve dual roles: as consequences of systemic dysfunction and as early indicators of developing disease elsewhere. The patient with new-onset nocturia warrants cardiovascular risk assessment. The man with OAB unresponsive to antimuscarinics may harbor inflammatory processes amenable to different interventions.

Practical applications follow from this framework. Metabolic optimization — weight management, glycemic control, perhaps urate-lowering therapy — deserves consideration alongside conventional pharmacotherapy for lower urinary tract symptoms. Evidence that allopurinol use correlates with reduced BPH risk supports this direction [10]. The proteomic identification of serotonergic dysregulation in OAB opens avenues beyond muscarinic blockade, potentially including agents modulating central and peripheral serotonin pathways. For acute retention, risk-stratified management using simple bedside parameters could guide catheter duration decisions and identify patients requiring closer follow-up or urodynamic evaluation.

The studies assembled here invite a conceptual shift. We have grown accustomed to treating bladder symptoms as isolated problems requiring localized solutions — antimuscarinics for urgency, alpha-blockers for obstruction, catheterization for retention. This approach will persist because it often works. But

the evidence increasingly points toward the organ oft considered as a passive conduit may in fact serve as a sensitive integrator — its dysfunctions clustering with metabolic syndrome, its molecular machinery responding to inflammatory and neural signals originating far beyond the pelvis, its recovery capacity indexing whole-body physiological reserve. Effective management of lower urinary tract symptoms may ultimately require attending not just to the bladder itself but to the systemic milieu in which it operates [11]. The challenge for neurourology lies in translating this integrative understanding into clinical practice that benefits individual patients.

• **Conflict of Interest:** No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Kim SJ, Kim JW, Cho YS, Chung KJ, Yoon H, Kim KH. Influence of circadian disruption associated with artificial light at night on micturition patterns in shift workers. *Int Neurourol* 2019;23:258-64.
2. Kim JW. Effect of shift work on nocturia. *Urology* 2016;87:153-60.
3. Kim SJ, Park SG, Pak S, Kwon O, Lee YG, Cho ST. Mediating effect of serum uric acid in the association between nocturia and mortality. *Int Neurourol J* 2025;29:277-85.
4. Lee SY, Lee JY, Yun SH, Lee M, Lee DE, Min JH, et al. Urothelial proteome changes underlying overactive bladder associated with bladder outlet obstruction. *Int Neurourol J* 2025;29:226-35.
5. Kim YJ, Tae BS, Bae JH. Cognitive function and urologic medications for lower urinary tract symptoms. *Int Neurourol J* 2020;24:231-40.
6. Holanda VAD, Oliveira MC, de Oliveira Torres CI, de Almeida Moura C, Belchior H, da Silva Junior ED, et al. The alpha1A antagonist tamsulosin impairs memory acquisition, consolidation, and retrieval in a novel object recognition task in mice. *Behav Brain Res* 2024;469:115027.
7. Niaga KJK, Rinaldi FX, Nathania N, Supinto PA, Lionardi SK, Yo R, et al. Do alpha-blockers and 5-alpha reductase inhibitors increase dementia risk? A network meta-analysis. *Int Neurourol J* 2025;29:236-47.
8. Mevcha A, Drake MJ. Etiology and management of urinary retention in women. *Indian J Urol* 2010;26:230-5.
9. Mayo ME, Lloyd-Davies RW, Shuttleworth KE, Tighe JR. The damaged human detrusor: functional and electron microscopic changes in disease. *Br J Urol* 1973;45:116-25.

10. Kukko V, Kaipia A, Talala K, Taari K, Tammela TLJ, Auvinen A, et al. Allopurinol and risk of benign prostatic hyperplasia in a Finnish population-based cohort. *Prostate Cancer Prostatic Dis* 2018;21: 373-8.
11. He Q, Wang Z, Liu G, Daneshgari F, MacLennan GT, Gupta S. Metabolic syndrome, inflammation and lower urinary tract symptoms: possible translational links. *Prostate Cancer Prostatic Dis* 2016; 19:7-13.