

Subjective and Objective Assessments of Olfactory Function in Patients Taking Anti-obesity Medications

Hye Jun Lee¹  · Il-Youp Kwak²  · Hyun Jin Min³ 

¹Department of Family Medicine, Chung-Ang University College of Medicine, Seoul, Korea

²Department of Applied Statistics, Chung-Ang University, Seoul, Korea

³Department of Otorhinolaryngology-Head and Neck Surgery, Chung-Ang University College of Medicine, Seoul, Korea

The global increase in obesity emphasizes the importance of effective treatment strategies, including pharmacological therapy [1]. Anti-obesity medications (AOMs), such as glucagon-like peptide-1 (GLP-1) receptor agonists (liraglutide, semaglutide), support weight reduction by modulating appetite and energy intake [2]. Olfaction, which influences appetite and energy balance, is frequently altered in obesity [3,4]. Although weight loss may affect olfactory function, findings across studies are inconsistent [5,6]. For example, GLP-1 receptor agonist therapy has been reported to improve olfactory test scores, enhance brain activation in response to olfactory stimuli (right parahippocampus), and increase olfactory-evoked neural activation (left hippocampus) [6,7]. Given the complexity of olfactory processing and the absence of standardized testing protocols, comprehensive evaluation using both subjective and validated objective tools is required [8-10]. Accordingly, this study aimed to evaluate olfactory function in patients undergoing AOM therapy and to identify related clinical characteristics.

The study protocol was approved by the Institutional Review Board of Chung-Ang University Hospital (Seoul, Korea; No. 2404-013-598). Participants were recruited from a tertiary hospital outpatient clinic between July 2024 and April 2025. Individuals aged ≥ 19 years who received a full explanation of the study and provided written informed consent were enrolled. Exclusion criteria were as follows: (1) pregnant or lactating women; (2) individuals with impaired judgment due to cognitive impairment,

such as dementia; (3) individuals with systemic health conditions considered unsuitable for medical obesity treatment; (4) individuals with symptoms of rhinitis or sinusitis previously diagnosed as allergic rhinitis or chronic rhinosinusitis; and (5) individuals presenting with nasal symptoms (e.g., nasal obstruction, rhinorrhea) or with abnormal findings on sinus endoscopy.

Anthropometric and laboratory assessments were performed at baseline. Subjective chemosensory symptoms were evaluated using the Questionnaire of Olfactory Disorders (QOD), which included items on general chemosensory discomfort and parosmia. Subjective olfactory discomfort and parosmia were defined as affirmative responses to the corresponding QOD items [11]. Objective olfaction was measured with the Yonsei Olfactory Function Test (YOF) (RHICO Medical Co.), with hyposmia defined as a Threshold-Discrimination-Identification (TDI) score ≤ 21 [12]. Gustatory function was assessed using a gustatory function test with validated taste solutions developed for the Korean population [13].

Participants received one or more of the currently approved AOMs (phentermine/topiramate, naltrexone/bupropion, liraglutide, semaglutide, and orlistat) for at least 24 weeks. Medications could be prescribed in combination at the physician's discretion based on the patient's baseline condition. Phentermine/topiramate combines a sympathomimetic amine and an anticonvulsant [14], whereas naltrexone/bupropion combines an opioid receptor antagonist and an aminoketone antidepressant, both of which contribute to appetite suppression [15]. Liraglutide and semaglutide are GLP-1 receptor agonists that delay gastric emptying and enhance insulin secretion, thereby producing weight-reducing effects [16]. Orlistat, a gastric and pancreatic lipase inhibitor, facilitates the physical excretion of dietary fat [17]. The distribution of participants according to AOM type was: phentermine/topiramate (n=62), naltrexone/bupropion (n=3), liraglutide (n=3), semaglutide (n=6), and orlistat (n=79). Statistical

• Received July 30, 2025

Revised September 11, 2025

Accepted September 23, 2025

• Corresponding author: **Hyun Jin Min**

Department of Otorhinolaryngology-Head and Neck Surgery, Chung-Ang University College of Medicine, 84 Heukseok-ro, Dongjak-gu, Seoul 06974, Korea

Tel: +82-2-6299-1765, Fax: +82-2-825-1765

Email: jjinient@gmail.com

© 2025 by Korean Society of Otorhinolaryngology-Head and Neck Surgery.

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

analyses included the *t*-test, chi-square test, and multivariate linear regression, conducted using R version 4.5.1 (R Foundation). Forest plots were used to visualize the results.

Seventy participants were included in the current analyses (Supplementary Fig. 1). The hyposmia group had a significantly higher body mass index (BMI) ($P=0.042$) and high-sensitivity C-reactive protein levels ($P=0.017$) (Table 1). Multiple linear regression analysis demonstrated that younger age ($P<0.001$), lower BMI ($P=0.005$), higher body fat percentage ($P=0.001$), and higher basal metabolic rate ($P=0.004$) were factors significantly associated with higher TDI scores. In contrast, elevated white blood cell count ($P=0.046$) and alanine aminotransferase (ALT) ($P=0.018$) levels were factors significantly associated with lower TDI scores (Fig. 1).

With respect to subjective olfactory dysfunction, 36 participants reported normal function and 34 reported subjective discomfort. Only age significantly differed between the groups, with the abnormal group being older than the normal group ($P=0.028$) (Supplementary Table 1). Multiple linear regression analysis revealed that higher BMI ($P=0.004$) and elevated ALT levels ($P=0.012$) were significantly associated with lower visual analog scale (VAS) scores, indicating less perceived olfactory dysfunction. Conversely, greater waist circumference ($P=0.013$), higher thyroid-stimulating hormone ($P=0.007$), and the presence of parosmia ($P=0.012$) were significantly associated with higher VAS scores (Supplementary Fig. 2).

Regarding parosmia symptoms, 50 of the 70 participants (71.4%) reported experiencing parosmia. Participants with parosmia showed significantly lower olfactory threshold scores than those without parosmia ($P=0.016$), suggesting reduced olfactory sensitivity (Supplementary Table 2). Additionally, low-density lipoprotein (LDL) cholesterol levels were significantly lower in the parosmia group ($P=0.036$). Multiple linear regression analysis showed that higher BMI ($P=0.012$), basal metabolic rate ($P=0.031$), and sum VAS score ($P=0.020$) were factors significantly associated with higher parosmia scores, indicating more severe parosmia. In contrast, higher body weight ($P=0.019$), skeletal muscle mass ($P=0.045$), and uric acid levels ($P=0.036$) were associated with lower parosmia scores (Supplementary Fig. 3).

In this study, 7% of participants were diagnosed with hyposmia based on the YOF, whereas 48.6% reported subjective olfactory disturbances and 71.4% reported experiencing parosmia. These findings highlight the importance of the assessment method used to evaluate olfactory function. Because different methodologies can produce divergent results, they should be carefully considered when interpreting olfactory function in both clinical and research contexts.

The YOF results indicated that BMI was the only factor consistently associated with both hyposmia diagnosis and TDI score. Interestingly, higher body fat mass was significantly associated with better olfactory function, as reflected by higher TDI scores. Few previous studies have directly compared BMI and body fat

Table 1. Characteristics of participants in the hyposmia and normosmia groups based on the YOF test

Variable	Hyposmia (n=5)	Normosmia (n=65)	P-value
Age (yr)	46.60±15.82	41.25±10.50	0.293
Sex			0.290
Male	2 (40)	13 (20)	
Female	3 (60)	52 (80)	
Non-smoker	5 (100)	61 (93.85)	1.000
Anthropometry			
Height (cm)	158.76±10.53	163.30±7.77	0.223
Weight (kg)	89.64±19.24	80.29±19.99	0.316
Body mass index (kg/m ²)	35.36±5.47	29.89±5.70	0.042
Skeletal muscle (kg)	29.54±9.17	27.05±6.30	0.412
Body fat percent (%)	41.58±6.17	38.10±6.28	0.236
Basal metabolic rate (kcal)	1,506.00±310.92	1,430.58±225.81	0.485
Waist circumference (cm)	107.68±11.61	97.09±14.63	0.120
YOF test			
Threshold	2.80±0.91	4.02±1.01	0.011
Discrimination	6.80±1.30	9.14±1.27	<0.001
Identification	10.80±1.30	11.72±0.52	0.001
TDI	20.40±0.82	24.88±1.58	<0.001
QOD test			
Discomfort (yes:no)	4:1	32:33	0.358
Parosmia score	20.00±19.18	20.51±19.88	0.956
Chemical taste test			
Taste threshold	5.20±0.45	5.54±0.56	0.193
Taste identification	16.00±2.55	18.20±3.27	0.147
Laboratory test			
WBC (×10 ⁹ /L)	7.46±2.31	6.72±1.91	0.408
hs-CRP (mg/L)	7.22±10.12	2.68±3.26	0.017
Fasting glucose (mg/dL)	100.20±10.71	105.95±27.61	0.647
Uric acid (mg/dL)	6.48±2.10	5.15±1.46	0.062
AST (IU/L)	33.20±26.75	29.03±28.16	0.750
ALT (IU/L)	53.40±58.13	32.26±26.87	0.129
Insulin (mU/L)	42.42±62.24	14.51±34.57	0.107
Total cholesterol (mg/dL)	190.40±48.64	208.72±37.01	0.300
Triglyceride (mg/dL)	155.80±113.19	120.12±69.01	0.292
HDL cholesterol (mg/dL)	48.20±12.46	56.32±14.26	0.221
LDL cholesterol (mg/dL)	101.20±32.56	121.06±35.12	0.225
Cortisol (µg/dL)	6.44±4.69	9.14±2.97	0.064
Free T4 (ng/dL)	1.40±0.24	1.28±0.17	0.120
TSH (µIU/mL)	2.41±1.26	2.49±1.51	0.913

Values are presented as mean±standard deviation or number (%).

YOF, Yonsei Olfactory Function Test; TDI, threshold discrimination identification; QOD, Questionnaire of Olfactory Disorders; WBC, white blood cell; hs-CRP, high-sensitivity C-reactive protein; AST, aspartate aminotransferase; ALT, alanine aminotransferase; HDL, high-density lipoprotein; LDL, low-density lipoprotein; Free T4, free thyroxine; TSH, thyroid-stimulating hormone.

mass with objectively measured olfactory function or hyposmia. As demonstrated in this study, BMI and body fat may exert different health effects [18]. Further research is therefore needed to clarify their distinct roles in olfactory function and dysfunction.

The factors associated with YOF outcomes and those associat-




Fig. 1. Forest plot illustrating the associations between clinical variables and Threshold-Discrimination-Identification scores. WBC, white blood cell; hs-CRP, high-sensitivity C-reactive protein; ALT, alanine aminotransferase; AST, aspartate aminotransferase.

ed with the sum of VAS scores did not fully overlap. Notably, higher BMI was associated with hyposmia diagnosis and lower TDI scores, yet it showed an inverse correlation with sum VAS scores. This suggests that BMI influences subjective olfactory discomfort and objective TDI scores in opposite directions. Such divergence underscores the importance of incorporating culturally validated subjective olfactory assessment tools, such as the QOD, when evaluating olfactory function in patients with obesity. Previous studies have reported associations between BMI and both objective and subjective measures of olfactory function. For example, BMI has been negatively correlated with TDI scores in patients with obesity compared to normal-weight controls [13], and a meta-analysis demonstrated that increasing BMI was linked to reduced olfactory threshold sensitivity [5]. However, studies in adolescents have shown that higher BMI may be associated with greater olfactory sensitivity, suggesting possible differences related to age or developmental stage [19]. In addition, one study reported that patients with subjective olfactory dysfunction had higher BMI values [20]. These findings indicate that future prospective studies with larger cohorts are needed to clarify the complex relationships between BMI, age, and olfactory function.

We also found that LDL cholesterol was the only clinical factor significantly associated with parosmia. Cardiovascular disease history is known to be linked with a higher prevalence of olfactory disorders [21,22], and LDL cholesterol plays a central role in the development and progression of atherosclerosis. The observed association between LDL cholesterol and parosmia may partially explain the relationship between cardiovascular disease and impaired olfactory function. Furthermore, a pharmacovigilance study reported olfactory adverse events, including parosmia, among neurological side effects related to GLP-1 receptor agonist use [23]. Since no objective olfactory function

test currently exists for specifically assessing parosmia, its reported prevalence may be overestimated. Nonetheless, incorporating parosmia assessment, at least through subjective scoring, may be essential when evaluating olfactory function in patients with obesity.

This study has some limitations. First, the sample size was relatively small, and only a few participants were diagnosed with hyposmia. Larger, prospective studies with comprehensive olfactory evaluation are needed to confirm these findings. Second, this cohort included only patients receiving AOMs and lacked a comparable control group. Consequently, the mechanisms underlying the discrepancy between subjective and objective assessments could not be determined. Finally, although we attempted to exclude other conditions that may influence olfaction, such as chronic rhinosinusitis and allergic rhinitis, based on symptoms and medical history, it is not possible to rule out all confounding factors completely.

Despite these limitations, we identified clinical factors associated with different olfactory parameters and showed that these factors may differ or even exert opposite associations depending on the outcome measure. This novel finding underscores the complexity of olfactory evaluation in this patient population.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

ORCID

Hye Jun Lee <https://orcid.org/0000-0001-5810-9787>

Il-Youp Kwak <https://orcid.org/0000-0002-7117-7669>
 Hyun Jin Min <https://orcid.org/0000-0003-3075-1350>

AUTHOR CONTRIBUTIONS


Conceptualization: HJL, HJM. Data collection: HJL, HJM. Formal analysis: IYK. Writing—original draft: HJL, HJM. Writing—review & editing: HJL, HJM. All authors submitted the final version, and reviewed the manuscript.

SUPPLEMENTARY MATERIALS


Supplementary materials can be found online at <https://doi.org/10.21053/ceo.2025-00238>.

REFERENCES


- World Health Organization (WHO). Obesity and overweight. WHO; 2023.
- Wilding JP, Batterham RL, Calanna S, Davies M, Van Gaal LF, Lingvay I, et al. Once-weekly semaglutide in adults with overweight or obesity. *N Engl J Med*. 2021 Mar;384(11):989-1002.
- Croy I, Nordin S, Hummel T. Olfactory disorders and quality of life—an updated review. *Chem Senses*. 2014 Mar;39(3):185-94.
- Manoel D, Moussa EA, Al-Naama A, Saraiva LR. The nose knows: olfaction as a metabolic gatekeeper in health and disease. *Physiology*. 2025 Nov;40(6):509-17.
- Peng M, Coutts D, Wang T, Cakmak YO. Systematic review of olfactory shifts related to obesity. *Obes Rev*. 2019 Feb;20(2):325-38.
- Zhang Z, Zhang B, Wang X, Zhang X, Yang QX, Qing Z, et al. Olfactory dysfunction mediates adiposity in cognitive impairment of type 2 diabetes: insights from clinical and functional neuroimaging studies. *Diabetes Care*. 2019 Jul;42(7):1274-83.
- Cheng H, Zhang Z, Zhang B, Zhang W, Wang J, Ni W, et al. Enhancement of impaired olfactory neural activation and cognitive capacity by liraglutide, but not dapagliflozin or acarbose, in patients with type 2 diabetes: a 16-week randomized parallel comparative study. *Diabetes Care*. 2022 May;45(5):1201-10.
- Khan R, Doty RL. GLP-1 receptor agonists significantly impair taste function. *Physiol Behav*. 2025 Mar;291:114793.
- Hummel T, Podlessek D. Clinical assessment of olfactory function. *Chem Senses*. 2021 Jan;46:bjab053.
- Jensen MM, Larsen KD, Homøe AS, Simonsen AL, Arndal E, Koch A, et al. Subjective and psychophysical olfactory and gustatory dysfunction among COVID-19 outpatients; short- and long-term results. *PLoS One*. 2022 Oct;17(10):e0275518.
- Choi WR, Jeong HY, Kim JH. Reliability and validity of the Korean version of the Questionnaire of Olfactory Disorders. *Int Forum Allergy Rhinol*. 2018 Dec;8(12):1481-5.
- Ha JG, Kim J, Nam JS, Park JJ, Cho HJ, Yoon JH, et al. Development of a Korean culture-friendly olfactory function test and optimization of a diagnostic cutoff value. *Clin Exp Otorhinolaryngol*. 2020 Aug;13(3):274-84.
- Hwang CS, Kim JW, Al Sharhan SS, Kim JW, Cho HJ, Yoon JH, et al. Development of a gustatory function test for clinical application in Korean subjects. *Yonsei Med J*. 2018 Mar;59(2):325-30.
- Smith SM, Meyer M, Trinkley KE. Phentermine/topiramate for the treatment of obesity. *Ann Pharmacother*. 2013 Mar;47(3):340-9.
- Billes SK, Sinnayah P, Cowley MA. Naltrexone/bupropion for obesity: an investigational combination pharmacotherapy for weight loss. *Pharmacol Res*. 2014 Jun;84:1-11.
- Knudsen LB, Lau J. The discovery and development of liraglutide and semaglutide. *Front Endocrinol (Lausanne)*. 2019 Apr;10:155.
- Heck AM, Yanovski JA, Calis KA. Orlistat, a new lipase inhibitor for the management of obesity. *Pharmacotherapy*. 2000 Mar;20(3):270-9.
- Padwal R, Leslie WD, Lix LM, Majumdar SR. Relationship among body fat percentage, body mass index, and all-cause mortality: a cohort study. *Ann Intern Med*. 2016 Apr;164(8):532-41.
- Herz RS, Van Reen E, Gredvig-Ardito CA, Carskadon MA. Insights into smell and taste sensitivity in normal weight and overweight-obese adolescents. *Physiol Behav*. 2020 Jul;221:112897.
- Patel ZM, DelGaudio JM, Wise SK. Higher body mass index is associated with subjective olfactory dysfunction. *Behav Neurol*. 2015 Jun;2015:675635.
- Liu G, Zong G, Doty RL, Sun Q. Prevalence and risk factors of taste and smell impairment in a nationwide representative sample of the US population: a cross-sectional study. *BMJ Open*. 2016 Nov;6(11):e013246.
- Seubert J, Laukka EJ, Rizzuto D, Hummel T, Fratiglioni L, Bäckman L, et al. Prevalence and correlates of olfactory dysfunction in old age: a population-based study. *J Gerontol A Biol Sci Med Sci*. 2017 Aug;72(8):1072-9.
- Chen H, Liu S, Gao S, Shi H, Yan Y, Xu Y, et al. Pharmacovigilance analysis of neurological adverse events associated with GLP-1 receptor agonists based on the FDA Adverse Event Reporting System. *Sci Rep*. 2025 May;15(1):18063.



Supplementary Fig. 1. Schematic depiction of the study diagram.



Supplementary Fig. 2. Forest plot illustrating the associations between clinical variables and the sum of the visual analog scale scores. ALT, alanine aminotransferase; TSH, thyroid-stimulating hormone.



Supplementary Fig. 3. Forest plot illustrating the associations between clinical variables and the parosmia scores. TSH, thyroid-stimulating hormone; ALT, alanine aminotransferase; VAS, visual analog scale.

Supplementary Table 1. Participant characteristics according to subjective olfactory discomfort

Variable	Subjective nose		P-value
	Normal (n=36)	Abnormal (n=34)	
Age (yr)	38.86±10.55	44.56±10.62	0.028
Sex			0.647
Male	9 (25.00)	6 (17.65)	
Female	27 (75.00)	28 (82.35)	
Non-smoker	33 (91.67)	33 (97.06)	1.000
Anthropometry			
Height (cm)	164.01±7.53	161.89±8.42	0.272
Weight (kg)	82.38±22.33	79.45±17.29	0.543
Body mass index (kg/m ²)	30.31±6.02	30.25±5.69	0.964
Skeletal muscle (kg)	27.80±6.72	26.61±6.27	0.448
Body fat percent (%)	38.19±5.46	38.50±7.16	0.839
Basal metabolic rate (kcal)	1,455.25±240.37	1,415.56±222.03	0.476
Waist circumference (cm)	98.00±14.79	97.69±14.68	0.931
YOF test			
Threshold	4.14±1.20	3.71±0.81	0.083
Discrimination	8.81±1.33	9.15±1.48	0.312
Identification	11.67±0.68	11.65±0.60	0.898
TDI	24.61±2.14	24.50±1.71	0.811
QOD test			
Parosmia	16.44±17.19	24.75±21.47	0.077
Olfactory function			0.358
Hyposmia	4 (11.11)	1 (2.94)	
Normosmia	32 (88.89)	33 (97.06)	
Chemical taste test			
Taste threshold	5.53±0.56	5.50±0.56	0.837
Taste identification	18.25±3.00	17.82±3.55	0.588
Laboratory test			
WBC (×10 ⁹ /L)	6.99±2.17	6.54±1.64	0.339
hs-CRP (mg/L)	3.86±5.28	2.08±2.13	0.074
Fasting glucose (mg/dL)	104.33±28.67	106.82±24.94	0.700
Uric acid (mg/dL)	5.41±1.51	5.08±1.57	0.373
AST (IU/L)	33.56±38.05	24.85±7.20	0.194
ALT (IU/L)	39.67±38.41	27.53±15.16	0.090
Insulin (mU/L)	16.55±31.37	16.44±43.05	0.990
Total cholesterol (mg/dL)	215.11±36.21	199.26±38.31	0.080
Triglyceride (mg/dL)	128.89±76.58	116.09±68.27	0.464
HDL cholesterol (mg/dL)	56.28±14.21	55.18±14.41	0.749
LDL cholesterol (mg/dL)	126.28±33.84	112.62±35.55	0.104
Cortisol (μg/dL)	9.08±3.06	8.82±3.28	0.733
Free T4 (ng/dL)	1.31±0.19	1.26±0.15	0.263
TSH (μIU/mL)	2.46±1.30	2.51±1.67	0.890

Values are presented as mean±standard deviation or number (%).

YOF, Yonsei Olfactory Function Test; TDI, threshold discrimination identification; QOD, Questionnaire of Olfactory Disorders test; WBC, white blood cell; hs-CRP, high-sensitivity C-reactive protein; AST, aspartate aminotransferase; ALT, alanine aminotransferase; HDL, high-density lipoprotein; LDL, low-density lipoprotein; Free T4, free thyroxine; TSH, thyroid stimulating hormone.

Supplementary Table 2. Participant characteristics according to subjective parosmia presence

Variable	Absence of parosmia (n=20)	Presence of parosmia (n=50)	P-value
Age (yr)	40.45±12.43	42.10±10.31	0.571
Sex			1.000
Male	4 (20)	11 (22)	
Female	16 (80)	39 (78)	
Non-smoker	19 (95)	47 (94)	1.000
Anthropometry			
Height (cm)	162.85±7.68	163.03±8.18	0.935
Weight (kg)	82.47±19.29	80.35±20.37	0.691
Body mass index (kg/m ²)	30.89±5.62	30.04±5.93	0.583
Skeletal muscle (kg)	27.22±5.86	27.23±6.78	0.996
Body fat percent (%)	39.44±6.39	37.90±6.27	0.359
Basal metabolic rate (kcal)	1,435.05±208.48	1,436.34±241.20	0.983
Waist circumference (cm)	99.27±15.34	97.28±14.45	0.611
YOF test			
Threshold	4.40±1.13	3.74±0.95	0.016
Discrimination	8.75±1.41	9.06±1.41	0.408
Identification	11.70±0.47	11.64±0.69	0.724
TDI	24.85±2.08	24.44±1.87	0.425
Olfactory function			1.000
Hyposmia	1 (5)	4 (8)	
Normosmia	19 (95)	46 (92)	
Subjective nose function			0.241
Normal	13 (65)	23 (46)	
Abnormal	7 (35)	27 (54)	
Chemical taste test			
Taste threshold	5.70±0.47	5.44±0.58	0.078
Taste identification	19.05±2.48	17.64±3.46	0.102
Laboratory test			
WBC (×10 ⁹ /L)	6.65±1.91	6.82±1.95	0.746
hs-CRP (mg/L)	3.61±3.62	2.77±4.37	0.451
Fasting glucose (mg/dL)	105.25±35.04	105.66±23.07	0.954
Uric acid (mg/dL)	5.58±1.44	5.12±1.57	0.265
AST (IU/L)	36.75±48.54	26.36±12.29	0.161
ALT (IU/L)	35.00±32.70	33.28±29.08	0.830
Insulin (mU/L)	21.77±55.85	14.39±26.94	0.458
Total cholesterol (mg/dL)	215.50±39.34	204.18±37.11	0.261
Triglyceride (mg/dL)	107.90±48.71	128.58±79.63	0.284
HDL cholesterol (mg/dL)	53.15±13.30	56.78±14.56	0.338
LDL cholesterol (mg/dL)	133.50±39.48	114.10±31.96	0.036
Cortisol (µg/dL)	9.02±2.42	8.92±3.42	0.909
Free T4 (ng/dL)	1.26±0.18	1.30±0.17	0.450
TSH (µIU/mL)	2.53±1.48	2.46±1.50	0.870

YOF, Yonsei Olfactory Function Test; TDI, threshold discrimination identification; WBC, white blood cell; hs-CRP, high-sensitivity C-reactive protein; AST, aspartate aminotransferase; ALT, alanine aminotransferase; HDL, high-density lipoprotein; LDL, low-density lipoprotein; Free T4, free thyroxine; TSH, thyroid stimulating hormone.