

SYSTEMATIC REVIEW **OPEN ACCESS**

Practice Guidelines for the Diagnosis and Treatment of Subepithelial Lesion Observed in Upper Gastrointestinal Endoscopy

Beom Jin Kim¹  | Seung Joo Kang²  | Su Youn Nam^{3,4}  | Chang Seok Bang⁵  | Byung-Hoon Min⁶  |
 Sung Eun Kim⁷  | Seung Young Kim⁸  | Hyunchul Lim⁹  | Chung Hyun Tae¹⁰  | Moon Won Lee¹¹  |
 Seung Han Kim¹²  | Sang Hoon Kim¹  | Jong Yeul Lee¹³  | Byung-Wook Kim¹⁴  | Jeong Seop Moon¹⁵  |
 Jong-Jae Park¹²  | Hwoon-Yong Jung¹⁶  | Korean College of Helicobacter and Upper Gastrointestinal Research

¹Department of Internal Medicine, Chung-Ang University College of Medicine, Seoul, South Korea | ²Department of Internal Medicine, Seoul National University Hospital Healthcare System Gangnam Center, Seoul, South Korea | ³Department of Internal Medicine, School of Medicine, Kyungpook National University, Daegu, South Korea | ⁴Division of Gastroenterology, Department of Internal Medicine, Kyungpook National University Chilgok Hospital, Daegu, South Korea | ⁵Department of Internal Medicine, Hallym University College of Medicine, Chuncheon, South Korea | ⁶Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, South Korea | ⁷Department of Internal Medicine, Kosin University College of Medicine, Busan, South Korea | ⁸Division of Gastroenterology and Hepatology, Department of Internal Medicine, Korea University Ansan Hospital, Korea University College of Medicine, Seoul, Republic of Korea | ⁹Department of Internal Medicine, Yongin Severance Hospital, Yonsei University College of Medicine, Yongin, South Korea | ¹⁰Department of Internal Medicine, Ewha Womans University College of Medicine, Seoul, South Korea | ¹¹Department of Internal Medicine, Pusan National University School of Medicine, Busan, South Korea | ¹²Division of Gastroenterology, Department of Internal Medicine, Korea University Guro Hospital, Korea University College of Medicine, Seoul, South Korea | ¹³Center for Gastric Cancer, National Cancer Center, Goyang, South Korea | ¹⁴Division of Gastroenterology, Department of Internal Medicine, Incheon St. Mary's Hospital, College of Medicine, The Catholic University of Korea, Seoul, South Korea | ¹⁵Department of Internal Medicine, Inje University College of Medicine, Seoul, South Korea | ¹⁶Department of Gastroenterology, University of Ulsan College of Medicine, Asan Medical Center, Asan Digestive Disease Research Institute, Seoul, South Korea

Correspondence: Hwoon-Yong Jung (hyjung@amc.seoul.kr)

Received: 9 June 2025 | **Revised:** 17 November 2025 | **Accepted:** 20 December 2025

Keywords: endoscopic treatment | guidelines | subepithelial lesion | systematic review

ABSTRACT

Subepithelial lesions (SELs) of the upper gastrointestinal tract are commonly detected during endoscopic examinations and encompass a broad spectrum of benign, potentially malignant, and malignant tumors. While most SELs are asymptomatic and incidentally found, accurate diagnosis remains challenging due to their subepithelial location, necessitating advanced imaging and tissue acquisition techniques. Endoscopic ultrasound (EUS) plays a crucial role in differentiating SELs, but its diagnostic accuracy remains limited. In response to the need for standardized clinical management, the Korean College of Helicobacter and Upper Gastrointestinal Research has developed evidence-based practice guidelines for the diagnosis and endoscopic treatment of SELs. These guidelines were established through a systematic review of existing literature and expert consensus, resulting in 11 key recommendations addressing diagnostic strategies, surveillance intervals, biopsy techniques, and indications for endoscopic or surgical resection.

Beom Jin Kim and Seung Joo Kang, both authors equally contributed as co-first authors.

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2026 The Author(s). *Journal of Gastroenterology and Hepatology* published by Journal of Gastroenterology and Hepatology Foundation and John Wiley & Sons Australia, Ltd.

1 | Introduction

A subepithelial lesion (SELs) refers to a raised lesion arising from the gastrointestinal wall between the lamina propria and the serosa, located beneath the epithelium and covered by mucosa identical to adjacent tissue [1]. These lesions are classified as benign or potentially malignant/malignant. Common benign types include leiomyomas, granular cell tumors, lipomas, and fibromas; common malignant or potentially malignant types include gastrointestinal stromal tumors (GIST), neuroendocrine tumors (NET), glomus tumors, and lymphomas [2, 3].

In a study analyzing approximately 80 000 patients who underwent screening endoscopy, SELs were found in 1.94% of the upper gastrointestinal tract, and gastric SELs were found in 0.74% of patients [4]. The mean size of gastric SELs varies across studies, typically ranging from 8 to 15 mm [4–7], with screening cohort reporting mean sizes of 8.7 mm [4] and studies with histological confirmation showing mean sizes of 13.6 mm [7]. Most gastric SELs are asymptomatic and incidentally discovered, although they can occasionally cause dysphagia, pain, obstruction, or bleeding [6].

Their endoscopic appearance is typically uniform, making visual differentiation difficult. Due to their subepithelial location, histological diagnosis by forceps biopsy is often challenging. Although EUS helps in differential diagnosis, its accuracy is limited (43%–50%) [7].

To improve diagnostic yield, EUS-guided fine-needle biopsy (EUS-FNB) and mucosal incision-assisted biopsy (MIAB) have been developed. These guidelines aim to provide evidence-based recommendations for the diagnosis, surveillance, and treatment of upper gastrointestinal SELs. Where evidence is limited, expert consensus supports the guidance. To our knowledge, this represents the first comprehensive clinical practice guideline for gastric SELs developed by a Korean medical society and published in English. Developed in the context of Korea's widely implemented gastric cancer screening program, these guidelines address the increasing detection of incidental SELs and provide standardized recommendations for their diagnosis, surveillance, and treatment.

2 | Development Methods for the Guideline

1. Clinical Practice Guideline Committee and multidisciplinary approach
2. Clinical practice guideline development process
 1. Selection of key questions
Key questions (KQs) were selected through review of international guidelines [8–11] and discussion among all committee members, based on the PICO framework. Eleven KQs were finalized.
 2. Literature search and selection
A comprehensive literature search was conducted in MEDLINE, Embase, Cochrane Library, and KoreaMed in September 2022, without restriction by year, language, or publication status. Duplicates were excluded. Two reviewers per KQ screened and selected relevant studies following PRISMA guidelines [12].

3. Literature quality assessment and meta-analysis
Quality was assessed using validated tools: RoB 2.0 for RCTs [13], Revised Risk of Bias Assessment tool for Non-Randomized Studies (ROBANS) for nonrandomized studies [14], Quality Assessment of Diagnostic Accuracy Studies-2 (QUADAS-2) for diagnostic studies [15], and A Measurement Tool to Assess Systematic Reviews-2 (AMSTAR-2) for systematic review literature evaluations [16]. Meta-analyses were conducted using random-effects models when appropriate, and heterogeneity assessed by I^2 . The most recent or largest study was selected when data duplication was suspected.
4. Derivation of recommendations and determination of recommendation strength and evidence level
Recommendations were formulated based on the GRADE approach. Strength (strong vs. conditional) reflected the balance of benefits and harms, while evidence certainty was rated from high to very low [17]. For insufficient evidence, expert consensus was used.
5. Agreement of recommendations and adoption
A modified Delphi consensus process was conducted with two rounds of voting among multidisciplinary experts. In the first round, 43 experts evaluated 11 key recommendations using a 5-point Likert scale. Consensus was defined as $\geq 70\%$ agreement, where agreement included “strongly agree” (1 point) or “agree with reservations” (2 points). Recommendations achieving consensus were accepted, while those with $<70\%$ agreement underwent revision based on expert feedback and were subjected to a second round of voting. In the second round, 28 experts participated to evaluate the revised recommendations. Panel consistency between rounds was assessed to ensure reliability of the consensus process.
6. Dissemination of clinical practice guidelines and revision plan
The final guidelines will be published domestically and internationally and posted on the society's website. Future revisions will be considered by the KCHUGR guideline committee if significant new evidence emerges.

The Clinical Practice Guideline Committee comprised a steering, working, and internal/external review group. The steering committee, composed of society leadership, oversaw strategy, budget approval, stakeholder coordination, and ensured independence. The working group included 14 members from the Korean College of Helicobacter and Upper Gastrointestinal Research and experts recommended by the Korean Physicians' Association. Methodologists also participated. Internal and external review committees had two members each. Meetings began in May 2022, with ongoing discussions and internal votes until December 2023, followed by a public hearing with field experts in March 2024.

3 | Key Questions and Recommendations

3.1 | Delphi Consensus Process Results

The first round of voting involved 43 experts evaluating 11 key recommendations. Seven recommendations (Questions 1, 2, 4, 8, 9, 10, and 11) achieved immediate consensus with agreement

rates ranging from 73.5% to 87.8%. Four recommendations did not reach the 70% threshold: Question 3 (69.4%), Question 5 (67.3%), Question 6 (69.4%), and Question 7 (63.3%), requiring revision and a second round of voting. Question 5 was subsequently divided into two sub-questions (5-1 and 5-2) for clarity.

In the second round, 28 experts participated, with 96.4% (27/28) having also participated in the first round, demonstrating high panel consistency. All five revised recommendations achieved consensus in the second round. Question 3 improved from 69.4% to 82.4% agreement, Questions 5-1 and 5-2 achieved 70.6% and 73.5%, respectively, Question 6 improved from 69.4% to 79.4%, and Question 7 showed the most substantial improvement from 63.3% to 79.4%.

The high retention rate of experts between rounds and the improved agreement rates in the second round indicate a robust consensus-building process. Final recommendations were adopted based on these voting results.

3.2 | KQ1: Is Endoscopic Ultrasonography Useful for the Diagnosis of Subepithelial Lesions Found on Endoscopy?

Statement 1. *EUS imaging is useful for the diagnosis of SELs because it can characterize the features of SELs (level of evidence: moderate; strength of recommendation: strong).*

EUS is a useful test for the diagnosis of SELs in the gastrointestinal tract because it can characterize the size, layer of origin, and presence of extrinsic compression. However, the accuracy of EUS varies depending on the size and type of SELs. Therefore, we conducted a literature search and analyzed studies that performed EUS on gastric SELs with confirmed histological findings.

In a review of 21 studies with histologically confirmed gastric SELs, EUS imaging accuracy (without tissue acquisition) ranged from 48% to 90.9% (Supplementary Figure 1) [7, 18–37]. All studies were prospective or retrospective observational studies, and quality was assessed using QUADAS-2 (Supplementary Table 1). Histologically confirmed SELs included GISTs, leiomyomas, ectopic pancreas, NETs, lipomas, schwannomas, lymphomas, duplication cysts, inflammatory fibroid polyps, MALT lymphomas, and others.

The sizes of SELs in the 21 studies ranged from 3.3 mm (mean) to 40 mm (median). In 14 studies that evaluated the accuracy of EUS imaging alone, EUS accuracy for all SELs ranged from 48% to 90.9% [7, 18–30]. This variation appears to be related to lesion size among other factors. One study reported 48% accuracy with a mean size of 3.3 mm [30], whereas a Korean study with a mean size of 25.2 mm (range 10–45 mm) reported 74.6% accuracy for GISTs [31]. While these studies suggest that EUS accuracy may improve with larger lesion size, the optimal size threshold for EUS evaluation requires further investigation. Current international guidelines generally recommend EUS evaluation for SELs ≥ 10 mm based on clinical experience and expert consensus [8–11].

The main clinical concern with SELs is their malignant potential. Thus, we assessed EUS diagnostic performance for GISTs, the most common potentially malignant SELs (Supplementary Table 2). In six studies ($N=381$), EUS showed sensitivity of 71%–95% and specificity of 33%–88% [23, 27, 31–34]. Two studies specifically on malignant GISTs reported sensitivities of 83.3% and 62.5%, specificities of 76.5% and 83.3%, and accuracies of 78.2% and 78.9% [32, 35].

Studies have also evaluated EUS accuracy by lesion type. Among studies with ≥ 5 cases per lesion type, accuracy ranged from 50% to 100% for NETs, 17% to 85.3% for leiomyomas, 7.7% to 92% for ectopic pancreas, 0% to 80% for lipomas, and 0% for schwannomas. Other results included 100% for inflammatory fibrous polyps, 62.5% for adenoid tumors, 11% for gastric/duplicate cysts, and 0% for vascular lesions. One prospective study ($n=48$) reported EUS sensitivity and specificity of 92% and 100% for extrinsic compression [26].

While some SELs can show surface or color changes, most are indistinguishable endoscopically. EUS helps identify the origin layer (e.g., muscularis mucosa, submucosa, or propria) and provides information on lesion echogenicity and homogeneity (Table 1). Malignant GISTs are predicted with 80%–92% sensitivity when EUS reveals >3 cm size, heterogeneity, and irregular margins [34, 36].

Although not included in our analysis, previous studies have evaluated contrast-enhanced EUS. A meta-analysis reported its sensitivity of 0.87 (95% CI: 0.82–0.91), specificity of 0.82 (95% CI: 0.74–0.89), and positive likelihood ratio of 3.55 (95% CI: 2.39–5.27) for predicting GISTs [37]. It also showed 0.93 sensitivity and 0.81 specificity for differentiating low- and high-risk GISTs [37].

EUS elastography has been rarely studied for SELs. A pilot study showed GISTs were stiffer than other SELs based on stiffness and color scoring. In another study using strain ratio, sensitivity and specificity were 100% and 94% for distinguishing GISTs (strain ratio: 51.1) from leiomyomas (strain ratio: 6.0), but differentiation from schwannomas (strain ratio: 62.0) remained difficult [38].

In summary, EUS imaging alone (without tissue acquisition) is useful for characterizing SELs, although its sensitivity and specificity are not sufficient for all types. Many included studies were retrospective with small samples, yet over 1500 SELs were analyzed. Diagnostic EUS is very safe, with a complication rate of 0.001% [39]. Given its clinical utility and minimal risk, EUS is strongly recommended for diagnosing gastric SELs.

3.3 | KQ2: Is It Recommended to Follow-Up With Endoscopy for Gastric Subepithelial Lesions Smaller Than 1 cm?

Statement 2. *For gastric SELs less than 1 cm in size that are not suspected to be malignant, endoscopic follow-up without further investigation may be considered (level of evidence: expert consensus; strength of recommendation: conditional).*

TABLE 1 | EUS characteristics of subepithelial tumors.

Subepithelial tumors	Layer of origin	EUS findings
Benign		
GIST-low risk	4th (rarely 2nd, 3rd)	Hypoechoic, homogenous, round, smooth margin, <3 cm
Leiomyoma	2nd, 4th	Hypoechoic, well-circumscribed
Ectopic pancreas	2nd, 3rd, 4th	Hypoechoic or mixed echogenicity (heterogeneous = acinous tissue, anechoic = ductal structures), indistinct margin
Schwannoma	4th	Hypoechoic
Lipoma	3rd	Intensely hyperechoic, homogeneous, smooth margin
Granular cell tumor	2nd, 3rd	Hypoechoic, heterogeneous
Inflammatory fibroid polyp	2nd, 3rd	Hypo- to hyperechoic, homogeneous, indistinct margin
Duplication cyst	Any or extramural	Anechoic, three to five layer wall, round or oval, absent Doppler signal
Lymphangioma	3rd	Anechoic with internal septa
Varices	3rd	Anechoic, serpiginous, Doppler positive
Malignant potential		
GIST-high risk	4th (rarely 2nd or 3rd)	Hypoechoic, heterogeneous, irregular extraluminal margin, cystic spaces, echogenic foci, >3 cm,
Neuroendocrine tumor	2nd, 3rd	Mildly hypoechoic, homogeneous, oval or round, smooth margin
Lymphoma	2nd, 3rd, 4th	Hypoechoic
Metastasis	Any or all	Hypoechoic, heterogeneous
Glomus tumor	3rd, 4th	Hypoechoic

Abbreviations: EUS, endoscopic ultrasonography; GIST, gastrointestinal stromal tumor.

Our literature search found no study directly comparing follow-up by endoscopy alone versus after EUS. Fifteen observational studies describing the natural course and endoscopic follow-up of gastric SELs were identified (Supplementary Figure 2) [40–54]. Study quality was assessed using ROBANS 2.0 (Supplementary Table 3).

Eight studies evaluated tumor growth and GIST proportion during follow-up of gastric SELs ≤ 3 –4 cm after EUS [42–44, 48–52]. Tumor enlargement occurred in 10%–28% of cases; overall, 22% increased in size. Among resected lesions, 51% were GISTs, with 13% classified as intermediate or high risk. In tumors <1 cm, only 17% showed size change.

Two studies followed gastric SELs <3 cm by endoscopy alone [45, 47]. With mean follow-up of 24–59 months, only 4.0%–6.4% of lesions <1 cm increased in size; among these, 1.5% were intermediate- or high-risk GISTs. Larger initial size was associated with more rapid and frequent growth. In two other studies, all SELs ≤ 2 cm were resected, and only 0.3% of ≤ 1 cm lesions were intermediate- or high-risk GISTs [53, 54].

In gastric SELs <1 cm with rolling sign and no high-risk features on EUS or biopsy (if performed), follow-up showed low growth rate (~5%) and slow enlargement. As the proportion of intermediate- or high-risk GISTs is very low, endoscopic follow-up alone is unlikely to impact prognosis (Supplementary Table 4).

3.4 | KQ3: Is Biopsy Recommended for Lesions Larger Than 2 cm Suspected to Be GISTs on Endoscopic Ultrasonography?

Statement 3. *Among the lesions suspected of being GISTs, if there are high-risk findings or if pathological diagnosis is required for treatment decision, a tissue biopsy may be considered (level of evidence: expert consensus; strength of recommendation: conditional).*

High-risk findings included size >3–5 cm, heterogeneous echotexture, irregular margins, and echogenic

foci.

The diagnostic accuracy of gastric SELs by EUS alone is limited (50%–64%) [7, 30]. Features like heterogeneity, large size (>3–5 cm), irregular margins, or internal changes may suggest GISTs [34, 55, 56], but interobserver variability limits reliability. Biopsy can enhance diagnostic accuracy.

A review of 16 studies on EUS-guided biopsy for gastric SELs (Supplementary Figure 3) [57–72] showed a diagnostic rate of 66.3% (95% CI: 62.3–69.7) with FNA and 76.2% (95% CI: 70.1–81.6) with FNB. Given these rates exceed those of EUS alone, histologic diagnosis is recommended when pathology informs

treatment decisions, such as for chemotherapy or surgery of cardia SELs (Supplementary Table 5).

3.5 | KQ4: Are Endoscopic Ultrasonography–Guided Fine-Needle Biopsy or Mucosal Incision–Assisted Biopsy Recommended for the Histological Diagnosis of SEL?

Statement 4. *EUS-FNB and MIAB are equally recommended for tissue acquisition from SEL, and MIAB is preferred if the size of the SEL is less than 2 cm (level of evidence: moderate; strength of recommendation: conditional).*

Standard forceps biopsy is often inadequate for SELs due to limited mucosal access. Bite-on-bite biopsies using jumbo forceps can improve yield, especially for submucosal (third layer) lesions [73, 74]. However, effectiveness is limited for muscularis propria lesions and may cause bleeding.

EUS-FNA and FNB are widely used for SELs, with diagnostic accuracy ranging from 46% to 93% [8, 75]. A meta-analysis comparing both methods showed FNB had higher accuracy and required fewer passes [76]. Adequate sample rates were 94.9% (95% CI: 92.3%–97.5%) for FNB and 80.6% (95% CI: 71.4%–89.7%) for FNA. MIAB (unroofing technique) involves mucosal incision and surface biopsy. In a meta-analysis of seven studies, its diagnostic yield was 89% (95% CI: 82.7%–93.5%) with a bleeding rate of 5.03% (95% CI: 0.4%–12.9%) [77]. Our meta-analysis of five studies comparing MIAB with EUS-FNA/FNB (Supplementary Figures 4, 5) [78–82] found diagnostic rates of 91% (95% CI: 86.0%–95.0%) for MIAB vs. 76.7% (95% CI: 69.7%–83.0%) for EUS-FNA/FNB, favoring MIAB (risk ratio 1.13, 95% CI: 1.04–1.24).

A subgroup analysis of three studies reported diagnostic rates by lesion size [78–80]. For lesions ≥ 2 cm, MIAB and EUS had similar yields: 91.2% (95% CI: 81.9%–97.4%) vs. 92.5% (95% CI: 83.5%–98.1%). For lesions < 2 cm, MIAB showed a higher rate of 90.1% (95% CI: 81.0%–96.4%) compared to 56.0% (95% CI: 43.3%–68.3%) for EUS, with a risk ratio of 1.59 (95% CI: 1.25–2.01; Supplementary Table 6).

3.6 | KQ5: When Should Follow-Up Examinations Be Conducted for Subepithelial Lesions Greater Than 1 cm?

Statement 5-1. *For SELs measuring 1–2 cm identified through endoscopy, if the diagnosis on endoscopic ultrasound is unclear, a follow-up endoscopy should be performed after 6 months, and if the size remains < 2 cm, a follow-up examination should be conducted after 12 months (level of evidence: expert consensus; strength of recommendation: conditional).*

Statement 5-2. *For SELs measuring 2 cm or larger identified through endoscopy, if there are no high-risk findings on endoscopic ultrasound, a follow-up endoscopy should be performed after 6 months, and if there is no significant size increase, another*

follow-up examination should be conducted after 12 months (level of evidence: expert consensus; strength of recommendation: conditional).

Significant size increase is defined as an increase of 25% or more in the maximum diameter, based on natural history studies of gastric SELs. Song et al. reported that among 954 upper gastrointestinal SELs followed for a mean of 47.3 months, only 34 (3.6%) demonstrated $\geq 25\%$ diameter increase [4]. This threshold represents clinically meaningful growth that exceeds typical measurement variability. Growth kinetics vary significantly by initial lesion size: lesions < 10 mm show annual growth of 1.31–1.68 mm, those measuring 10–20 mm grow 2.09–3.83 mm per year, while lesions > 20 mm demonstrate 2.28–7.13 mm annual growth [45]. Studies confirm that growth rates correlate with initial size, with larger lesions showing more rapid progression [45, 49]. The 25% criterion provides an objective measure to distinguish true tumor progression from measurement error while guiding clinical decision-making regarding surveillance intervals and intervention timing.

SELs such as GISTs and NETs may show malignant behavior, requiring resection. Differentiating benign tumors (e.g., lipomas, heterotopic pancreas, granular cell tumors, schwannomas, leiomyomas) from potentially malignant tumors is critical. While initial detection often occurs during screening endoscopy with visual size estimation, endoscopic ultrasound should be performed when SELs appear to be 1 cm or larger for accurate size measurement, as visual estimation may be inaccurate, particularly for lesions with extraluminal components.

No specific prospective study on follow-up for gastric SELs ≥ 10 mm was found, but 15 studies reported their natural course and endoscopic follow-up (Supplementary Figure 6; Supplementary Table 7). For GISTs, tumor doubling time was 24 months in low-risk, 17.1 in intermediate, and 3.9 in high-risk groups [40]. Other studies showed annual growth rates of 1.57–1.67 mm (< 10 mm), 1.68–3.83 mm (10–20 mm), and 3.72–7.13 mm (> 20 mm), with faster growth in larger lesions (Table 2) [45, 49].

EUS aids in differentiating SELs, with sensitivity for GISTs of 62.5%–83.3%, specificity 76.5%–83.3%, and accuracy 78.2%–78.9%, although definitive diagnosis is not always possible. If GIST is suspected but unclear, biopsy is preferred; otherwise, endoscopic follow-up may be needed. For SELs < 20 mm, even if identified as GISTs, high-risk potential is low (0%–2.5%) [53]. If EUS findings are inconclusive, follow-up at 6 months is recommended to assess size, and if stable, re-evaluation after 12 months can be considered [10].

For SELs > 20 mm found during initial endoscopy, biopsy may be considered if GIST differentiation is needed. If high-risk GIST features are absent on EUS, international guidelines suggest follow-up endoscopy after 6 months to assess growth, and if stable, another exam at 12 months may follow [10]. Clearly benign lesions (lipomas, heterotopic pancreas, granular cell tumors, schwannomas, leiomyomas, glomus tumors) on EUS do not require regular follow-up, regardless of size [10].

TABLE 2 | Growth rates (mm/year) of subepithelial tumors according to their initial size.

	<1 cm	≥1 cm to >2 cm	≥2 cm	p
Kim et al. (2011) n = 84	1.68	2.64	3.72	0.003
Song et al. (2015) n = 34	1.31 ± 1.38	2.09 ± 1.47	2.28 ± 1.90	0.059
Kim et al. (2022) n = 138	1.57 ± 1.23	3.83 ± 4.24	7.13 ± 9.71	<0.001

3.7 | KQ6: Is Endoscopic Resection More Effective Than Follow-Up for Proven or Suspected GISTs Less Than 2 cm?

Statement 6. *Endoscopic resection is recommended for proven or suspected gastric GISTs less than 2 cm to establish definitive diagnosis and risk stratification. Close follow-up may be considered as an alternative after multidisciplinary discussion (level of evidence: very low; strength of recommendation: conditional).*

Management recommendations differ significantly based on diagnostic certainty. For proven GISTs <2 cm (histologically confirmed), endoscopic or surgical resection is recommended as the preferred approach to establish definitive risk stratification, as even small GISTs may harbor intermediate or high-risk features (2.4% in our analysis). Surveillance may be considered only after multidisciplinary discussion in selected patients with significant comorbidities or limited life expectancy. For SELs suspected but not proven to be GISTs, tissue acquisition should be attempted first (Statement 4). If GIST is confirmed, proceed with resection. If tissue diagnosis remains inconclusive despite adequate sampling, management should be individualized based on EUS features, patient factors, and shared decision-making. For undiagnosed SELs <2 cm without high-risk features, surveillance according to Statement 5-1 is appropriate, as the majority will be benign lesions not requiring intervention.

Size and mitotic index are key factors in determining GIST malignancy. According to the 2008 modified NIH criteria, gastric GISTs <2 cm with ≤5 mitoses are very low risk, while those with >5 are high risk [83]. Thus, even small GISTs require risk stratification through pathology.

Tissue acquisition techniques have improved, but EUS-FNA/B diagnostic yield remains size-dependent, at ~70% for 1–2 cm lesions [84, 85]. One study reported 100% accuracy with onsite pathology, although mitotic index correlation with surgical specimens was unclear [86]. EUS-FNA and bite-on-bite biopsy cannot reliably assess mitotic rate due to tumor heterogeneity [87], and biopsy-confirmed GISTs may still carry malignancy risk despite low-risk classification.

No RCTs have compared follow-up and resection for gastric GISTs <2 cm. A prospective study of 972 SETs <2 cm resected endoscopically found 42.3% were GISTs [53]. Of these, 97.6% were very low or low risk, but 2.4% were intermediate (1.9%) or high risk (0.5%). In a retrospective study, only 10.5% of 66 gastric SETs <2 cm were malignant GISTs [88]. While malignancy was suspected by EUS in 22 of 26 cases, most were benign. Three malignancies were missed by EUS. EUS sensitivity, specificity, and accuracy were 50%, 62%, and 60%. Additionally, 73% of FNAs were noncontributive. Thus, EUS and FNA/B are unreliable for predicting malignancy, and endoscopic resection should be considered for definitive diagnosis. Even for <2 cm GISTs, resection requires multidisciplinary discussion.

3.8 | KQ7: Is Endoscopic Resection More Effective Than Surgery for Gastric GIST Larger Than 2 cm?

Statement 7. *Surgical resection is recommended for gastric GIST larger than 2 cm. However, endoscopic resection may be considered depending on the experience of the endoscopist (level of evidence: low; strength of recommendation: conditional).*

Laparoscopic resection is standard for GISTs, offering efficacy, safety, and short hospital stays. With improved endoscopic techniques, resection is now a viable option for smaller tumors. ESD and submucosal excavation are feasible for submucosal tumors, while EFTR is used for those in the muscularis propria.

We reviewed 2026 articles and included 13 retrospective studies comparing endoscopic resection and surgery for gastric GISTs >2 cm, focusing on efficacy, complications, and recurrence (Supplementary Figures 7, 8) [89–101]. Most surgeries were performed laparoscopically, except for one patient who underwent both open and laparoscopic surgeries. Regarding endoscopic resection, there were five articles on endoscopic resection [89–91, 93, 101], four articles on ESD [95, 96, 99, 100], and others on EFTR [92, 94, 97, 98]. In the meta-analyses, surgery was superior to endoscopic resection in terms of complete resection (OR: 8.31; 95% CI: 2.80–24.69, $I^2 = 39\%$; Supplementary Figure 9). However, recurrence (OR: 0.52; 95% CI: 0.21–1.31; $I^2 = 0\%$) and complications (OR: 0.91; 95% CI: 0.58–1.45; $I^2 = 0\%$) were similar between groups (Supplementary Figures 10, 11). Endoscopic resection took less time than surgery despite high heterogeneity (mean difference: −39.32 min; 95% CI: −60.67 to −17.96; $I^2 = 95\%$; Supplementary Figure 12). Thus, both approaches appear feasible for gastric GISTs >2 cm.

In endoscopic resection, lesion relation to the muscularis propria is key. ESD is suitable when GISTs are narrowly connected (subtype 1) or broadly attached at an obtuse angle (subtype 2) to the muscularis propria and protrude into the lumen. For lesions within the wall (subtype 3) or protruding toward the serosa (subtype 4), EFTR or surgery is recommended [102].

In addition, because most GISTs invade the muscularis propria, a high level of skill is important for the success of the

procedure. R0 resection is not easy; there is a risk of perforation, and many cases are located in the fundus and cardia [103]. Therefore, a medical environment should be established where smooth treatment can be achieved in cases where endoscopic resection is initiated but collaboration with surgery is required. Treatment of gastric type 1 NETs <1 cm is controversial and also for “very small” gastric type 1 NETs. “Very small” or “small than 5 mm”. There are patients with multiple gastric type 1 NETs smaller than 1 cm; how to manage in this clinical scenario?

3.9 | KQ8: What Is the Optimal Treatment Approach for Gastric NETs <2 cm Based on Type and Tumor Characteristics?

Statement 8. *Endoscopic resection or surgery can be considered for gastric neuroendocrine tumors less than 2 cm, depending on the NET type (type 1, 2, or 3), size, invasion depth, and tumor grade (level of evidence: very low; strength of recommendation: conditional).*

Gastric NETs are classified into three types: type 1 (70%–80%) with chronic atrophic gastritis and hypergastrinemia; type 2 (5%–10%) with Zollinger–Ellison syndrome; and type 3 (10%–15%) with normal gastrin and no associated disease [104]. Treatment options include endoscopic resection or surgery, depending on size, grade, and depth. The management approach for gastric NETs <2 cm varies significantly by type:

- Type 1 (<1 cm): Surveillance is the preferred approach, with excellent prognosis and virtually no risk of metastasis. Surgical resection is not indicated.
- Type 1 (1–2 cm): Endoscopic resection may be considered based on grade and invasion depth.
- Types 2 and 3: More aggressive management with endoscopic resection or surgery based on individual factors.

We reviewed six retrospective studies on gastric NETs <2 cm, focusing on outcomes like lymph node metastasis, recurrence, and death (Supplementary Figures 13, 14) [105–110].

No prospective studies have compared surgery and endoscopic treatment for gastric NETs, and most focus on oncologic outcomes. For type 1 NETs <1 cm, studies consistently show excellent prognosis with surveillance alone, with no reported lymph node metastases or disease-specific mortality, making surgical resection unnecessary in virtually all cases. However, outcomes vary for 1–2 cm lesions. In a study of 123 patients, lymph node metastasis was 0%, 8.3%, and 100% depending on invasion depth, and 0% in grades 1–2 vs. 50% in grade 3 [110]. In a Taiwanese study of 187 patients with type 1 gastric NETs, no lymph node or distant organ metastasis occurred in lesions <1 cm [106]. Yet, for 1–2 cm grade 1 NETs with muscularis propria invasion, metastasis was reported. Among four grade 2 NETs (1–2 cm), half invaded mucosa/submucosa, and half invaded the muscularis propria. Thus, while <1 cm lesions show similar outcomes across treatments, 1–2 cm NETs require caution based on depth and grade.

No RCTs have compared endoscopic resection and surgery for type 3 gastric NETs. In a multicenter retrospective study ($n=176$), lymph node metastases occurred in 8.4%, but no recurrence was noted in 53 patients with G1 submucosal lesions <1 cm [111]. Korean retrospective studies have also reported no recurrence in G1 lesions confined to the submucosal layer of less than 1.5 cm [112, 113].

In summary, for type 1 gastric NETs <1 cm, endoscopic resection is preferred if submucosal; for 1–2 cm lesions, treatment depends on depth and grade. Type 3 NETs <2 cm require individualized decisions based on lesion size, invasion depth, and grade.

3.10 | KQ9: Is Follow-Up Observation Possible for Gastric Type 1 Neuroendocrine Tumors Smaller Than 1 cm?

Statement 9. *We suggest endoscopic resection for type 1 NETs measuring <1 cm. However, regular endoscopic surveillance may also be considered for very small lesions (level of evidence: very low; strength of recommendation: conditional).*

Gastric NETs account for approximately 7% of all digestive NETs [114] and are subdivided into three main types [115]. Type 1 gastric NETs have a low risk of metastasis, which is directly correlated with tumor size [110]. Endoscopic treatment is often performed based on the type, size, and grade of the tumor [10].

To assess the feasibility of follow-up for gastric NETs <1 cm, we identified 1017 studies. Of 843 initially screened, five original articles (since 2019) and five review articles (since 2017) were selected by title/abstract. Full texts were reviewed, and three original articles from 2019, four from reviews, and one via manual search, were included for final quality assessment (Supplementary Figures 15, 16; Supplementary Table 8).

Seven studies assessed death or disease progression without treatment, and one analyzed lymphovascular invasion by tumor size after endoscopic resection. Most were retrospective; one was prospective. In untreated patients ($n=8-87$), no NET-related deaths occurred. In 11 patients with type 1 gastric NETs followed for a median of 54 months (range 9–136), four showed increased lesion numbers, but no lesions >1 cm or metastases were observed [116]. In a Japanese multicenter study ($n=82$), including endoscopic surveillance ($n=25$), endoscopic resection ($n=41$), and surgery ($n=16$), no untreated patients developed metastasis, progression, or died. RFS was 97.6%, and disease-specific survival 100% [109]. In a single-center study of 26 patients, eight were monitored without treatment (2–13 mm); one died, but none from NETs [117]. Among 97 gastric NET patients, 3 underwent surgery, 45 received curative endoscopic resection, 13 had observation only, and 36 received somatostatin analogs. All remained alive and metastasis-free at follow-up [118].

A Korean retrospective study identified 104 type 1 gastric NETs among 223 cases diagnosed from 1996 to 2011 [119]. Of these, 67 patients received endoscopic treatment, and 27 were observed

without intervention. Complete resection was achieved in 53 patients (79.1%) and recurrence occurred in 14 (20.9%). Among those observed, 19 (70.4%) had stable disease and 8 (29.6%) showed progression. Mean tumor size did not differ by progression status, but four patients had tumors >10mm, and tumor grade was not reported. Mean follow-up was 49 months (range 31–210), with no deaths in either group.

In a recent retrospective study, 87 of 115 patients with grade 1 gastric NETs (75 with tumors ≤ 1 cm) underwent endoscopic surveillance without initial treatment [120]. Over a median 62-month follow-up, 91% ($n = 79$) showed no clinically significant changes in tumor size or grade. Among ≤ 1 cm tumors ($n = 75$), no size increase occurred. Three patients progressed from G1 to G2 but received no further treatment due to comorbidities: two regressed to G1, and one remained G2. Among 1–2 cm tumors ($n = 12$), three had >0.5 cm growth. Overall, G1 NETs <1 cm showed no meaningful progression.

A recent prospective study followed 127 gastric NETs (G1, $n = 80$) with endoscopy [105]. Tumors <1 cm were resected with forceps or cold snare; larger ones by EMR or ESD. Among 127 lesions: 87.4% were <0.5 cm, 8.7% were 0.6–1 cm, 3.1% were 1.1–2 cm, and 0.8% were >2 cm. Of the G1 NETs <0.5 cm, 15 patients (18.7%) were safely observed without resection. In a study of 155 cases (165 lesions) of G1 NETs, 24.8% had lymphovascular invasion [121]. Even in G1 tumors, the rate was 22.8%, with 3.4% for <0.5 cm and 8.7% for 0.5–1 cm. To keep invasion $\leq 10\%$, tumor size needed to be ≤ 0.8 cm; for $\leq 5\%$ invasion, ≤ 0.6 cm. A 0.6-cm cutoff yielded 79% sensitivity and 63% specificity.

The management of type 1 gastric NETs <1 cm remains controversial internationally. While some guidelines increasingly favor surveillance, our data showing 3.4%–8.7% lymphovascular invasion rates even in small lesions cannot be overlooked. The optimal approach may vary based on healthcare system resources and accessibility. In Korea's context with excellent endoscopic access and technical expertise, both resection and surveillance are valid options, with the choice individualized based on patient factors and preferences.

Based on the aforementioned evidence, in terms of lesion progression, in the case of very small grade 1 NETs measuring less than 0.5 cm, follow-up without resection may be an option. However, lymphovascular invasion occurs in 3.4% of patients with tumors less than 0.5 cm; therefore, endoscopic observation without complete resection can be considered only in cases where regular endoscopic examination is possible. There was very little evidence regarding survival rate, disease progression, and lymphovascular invasion (Supplementary Tables 9–11).

For patients with multiple type 1 gastric NETs <1 cm, management should be individualized. Options include: (1) selective resection of the largest lesions combined with surveillance of smaller ones, (2) complete resection of all visible lesions if technically feasible, or (3) surveillance alone if lesions are all <5 mm with regular endoscopic monitoring. The decision should consider the total number of lesions, patient age, comorbidities, and ability to comply with surveillance protocols. Evidence for managing multiple small type 1 NETs remains limited and represents an area requiring future research.

In summary, for gastric grade 1 neuroendocrine tumors (NETs) smaller than 0.5 cm, active surveillance without resection may be considered, although the condition itself may be of concern. Both the anticipated benefits and harms of this approach are minimal, the supporting evidence is of very low certainty, and there is considerable uncertainty regarding patient values and preferences. Therefore, no clear superiority can be determined between resection and surveillance. In the context of Korea, where biennial national cancer screening and high accessibility to healthcare services are available, both acceptability and feasibility of surveillance are considered generally high. Taking these factors into account, a conditional recommendation is made in favor of surveillance over resection in such cases. In the voting of related professional societies, 88.4% (38/43) agreed with the recommendation; however, at the public hearing, there were concerns about the follow-up observations of gastric NETs smaller than 0.5 cm. In some cases, small gastric NETs are almost completely removed or degenerated by biopsy and are not visible when endoscopy is performed again for resection. Progression of small, grade I, and type I gastric NET is extremely rare. Considering the conditions in Korea, where access to gastroscopy is very high, small, grade I and type I gastric NETs can be observed without the need for resection. However, small grade I and type I gastric NETs can show an aggressive course, although this is very rare. In Korea, criminal prosecutions for medical negligence are 750 times higher than those in the UK (750 cases per year in Korea and 1–2 cases per year in the UK). Legal prosecution is possible even in unusual circumstances. Due to the high risk of exposure to legal risks, it was recommended to revise the size to “very small” rather than specifying it. In the final recommendation, “less than 0.5 cm” was revised to “very small,” which generally refers to lesions ≤ 5 mm, though this threshold requires clinical discretion based on individual patient factors.

3.11 | KQ10: Which Endoscopic Resection Technique Should Be Used to Resect Subepithelial Tumors?

Statement 10. *Endoscopic resection methods for gastric SELs include endoscopic submucosal tunnel resection, EFTR, and ESD, depending on the experience of the endoscopist and the circumstances of the institution (level of evidence: low; strength of recommendation: conditional).*

Advances in endoscopic instruments and techniques now allow endoscopic resection to be considered as an alternative to surgery for gastric SELs, depending on clinical context. Thirty-six studies reported outcomes of endoscopic resection, although most were retrospective, with only seven directly comparing procedures. The selection and quality assessment process is shown in Supplementary Figure 17. In one study comparing ESD and EFTR, en bloc resection rates were 96.2% and 95%, respectively. Another comparing submucosal tunnel resection and EFTR showed rates of 93.3%–95% and 96.4%–96.8% [122–124].

For gastric SELs in the muscularis propria, ESD showed en bloc resection rates of 90.2%–100% and complete resection rates of 74.2%–100% [125–135]. EFTR achieved 92.8%–100% and

97.3%–100%, respectively, while submucosal tunnel resection yielded 84.7%–100% en bloc and 96.8%–100% complete resection rates [122–124, 136–146]. A meta-analysis of nine ESD studies ($n = 288$) reported a pooled complete resection rate of 86.2% (95% CI: 78.9–91.3); for muscularis propria–origin tumors, the rate was 84.4% (95% CI: 78.7–88.8). Thus, EFTR and submucosal tunnel resection may be more effective than ESD for tumors in the muscularis propria. However, as long-term recurrence rates are similar, choice of procedure can depend on lesion characteristics and operator preference. In tumors within the submucosal layer, mostly gastric NETs, four studies reported complete resection rates of 69%–100% for EMR and 66.6%–92.8% for ESD [119, 147, 148].

Considering the above findings, when SELs are located in the muscularis propria of the stomach, the efficacy of EFTR and endoscopic submucosal tunnel resection is high, considering the en bloc and complete resection rates of these procedures. However, because few institutions have much experience with the above two procedures, ESD can also be considered an option in terms of oncological results. ESD is superior to EMR for gastric SELs in the submucosal layer.

3.12 | KQ11: What Is the Follow-Up Strategy After Endoscopic Resection of Subepithelial Lesions?

Statement 11. The follow-up strategy after endoscopic resection of SELs should be determined based on the type of SELs and whether complete resection was achieved (level of evidence: expert consensus; strength of recommendation: conditional).

1. Completely resected benign SELs are not recommended for follow-up examinations.
2. Completely resected GISTs and NETs are recommended for regular endoscopic and radiological follow-up examinations.
3. In cases of incomplete resection, follow-up observation, additional endoscopic resection, or surgical resection is considered depending on the type of resection.

The KQ addresses the follow-up strategy after endoscopic resection of SELs. The PICO framework was as follows: patient, those who underwent endoscopic resection for SELs; intervention, regular follow-up examinations performed after endoscopic resection; comparator, no regular follow-up examinations; and outcome, benefits and risks, including recurrence-free survival and costs. The number of articles retrieved from the systematic literature review is shown in Supplementary Table 12.

One article titled “How should patients with completely resected gastrointestinal stromal tumors (GIST) be followed-up?” analyzed 33 patients post-resection to assess recurrence [149]. Although limited and indirect, the study suggests endoscopic follow-up every 6 months for 2 years, then annually for 3 years. However, the small sample size limits generalizability. The 2022 ESGE guidelines state that SEL follow-up depends on lesion type and completeness of resection. No follow-up is needed for

completely resected benign SELs, except type I gastric NETs, which require endoscopy every 1–2 years. According to the US NCCN guidelines, type I gastric NETs should be followed with endoscopy every 6–12 months for the first 3 years, then annually. Type 2 and 3 gastric NETs warrant annual surveillance, while G3 or >20 mm NETs require endoscopy every 4–6 months during year one, then annually [10, 150].

For GISTs, the ESGE guidelines recommend only general imaging examinations, such as CT, because local recurrence is rare. However, owing to the limited research on the natural course after endoscopic resection, endoscopic examinations are performed every 6–12 months for the first 3 years after resection. In cases of incomplete GIST resection, reoperation or re-endoscopic procedures are necessary. If this is not feasible, the tumor should be considered locally advanced, and the treatment plan should be determined through multidisciplinary consultation [10, 151, 152].

Funding

The authors have nothing to report.

Ethics Statement

Given the nature of this study as a systematic review and meta-analysis of previously published research, and the absence of direct involvement with human subjects or data collection, Institutional Review Board (IRB) approval was not required.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Data Transparency

All data support the published claims and comply with field standards.

References

1. M. W. Lee and B. E. Lee, “Approach to Small Gastric Subepithelial Lesions,” *Korean Journal of Helicobacter and Upper Gastrointestinal Research* 23, no. 1 (2023): 28–33.
2. S. G. Kim, “Incidental Gastrointestinal Subepithelial Mass,” *Korean Journal of Gastroenterology* 56, no. 6 (2010): 341–345.
3. S. E. Kim and M. I. Park, “Natural Course of Gastric Subepithelial Tumor,” *Korean Journal of Helicobacter and Upper Gastrointestinal Research* 15, no. 1 (2015): 1–8.
4. J. H. Song, S. G. Kim, S. J. Chung, H. Y. Kang, S. Y. Yang, and Y. S. Kim, “Risk of Progression for Incidental Small Subepithelial Tumors in the Upper Gastrointestinal Tract,” *Endoscopy* 47, no. 8 (2015 Aug): 675–679.
5. J. H. Lee, H. L. Lee, Y. W. Ahn, et al., “Prevalence of Gastric Subepithelial Tumors in Korea: A Single Center Experience,” *Korean Journal of Gastroenterology* 66, no. 5 (2015 Nov): 274–276.
6. J. L. Humphris and D. B. Jones, “Subepithelial Mass Lesions in the Upper Gastrointestinal Tract,” *Journal of Gastroenterology and Hepatology* 23, no. 4 (2008 Apr): 556–566.

7. C. Karaca, B. G. Turner, S. Cizginer, D. Forcione, and W. Brugge, "Accuracy of EUS in the Evaluation of Small Gastric Subepithelial Lesions," *Gastrointestinal Endoscopy* 71, no. 4 (2010 Apr): 722–727.
8. Standards of Practice Committee, A. L. Faulx, S. Kothari, et al., "The Role of Endoscopy in Subepithelial Lesions of the GI Tract," *Gastrointestinal Endoscopy* 85, no. 6 (2017): 1117–1132.
9. K. Sharzei, A. Sethi, and T. Savides, "AGA Clinical Practice Update on Management of Subepithelial Lesions Encountered During Routine Endoscopy: Expert Review," *Clinical Gastroenterology and Hepatology* 20, no. 11 (2022): 2435–2443.e4.
10. P. H. Deprez, L. M. G. Moons, O. T. D., et al., "Endoscopic Management of Subepithelial Lesions Including Neuroendocrine Neoplasms: European Society of Gastrointestinal Endoscopy (ESGE) Guideline," *Endoscopy* 54, no. 4 (2022): 412–429.
11. B. C. Jacobson, A. Bhatt, K. B. Greer, et al., "ACG Clinical Guideline: Diagnosis and Management of Gastrointestinal Subepithelial Lesions," *American Journal of Gastroenterology* 118, no. 1 (2023): 46–58.
12. D. Moher, A. Liberati, J. Tetzlaff, D. G. Altman, and PRISMA Group, "Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement," *PLoS Medicine* 6, no. 7 (2009): e1000097.
13. J. A. C. Sterne, J. Savović, M. J. Page, et al., "RoB 2: A Revised Tool for Assessing Risk of Bias in Randomised Trials," *BMJ* 366 (2019): 14898.
14. I. Hinneburg, "ROBINS-1: A Tool for Assessing Risk of Bias in Non-Randomised Studies of Interventions," *Medizinische Monatsschrift für Pharmazeuten* 40, no. 4 (2017 Apr): 175–177.
15. P. F. Whiting, A. W. S. Rutjes, M. E. Westwood, et al., "QUADAS-2: A Revised Tool for the Quality Assessment of Diagnostic Accuracy Studies," *Annals of Internal Medicine* 155, no. 8 (2011): 529–536.
16. B. J. Shea, B. C. Reeves, G. Wells, et al., "AMSTAR 2: A Critical Appraisal Tool for Systematic Reviews That Include Randomised or Non-Randomised Studies of Healthcare Interventions, or Both," *BMJ* 358 (2017): j4008.
17. S. Y. Kim, M. Y. Choi, H. J. Kim, et al., *NECA'S Handbook for Clinical Practice Guideline Developer Version 2.0* (National Evidence Based Healthcare Collaborating Agency, 2022).
18. S. Y. Kim, K. N. Shim, J. H. Lee, et al., "Comparison of the Diagnostic Ability of Endoscopic Ultrasonography and Abdominopelvic Computed Tomography in the Diagnosis of Gastric Subepithelial Tumors," *Clinical Endoscopy* 52, no. 6 (2019): 565–573.
19. G. R. Lu, L. Miao, Y. H. Zhou, Z. Z. Cai, C. L. Xu, and Z. X. Xue, "The Value and Misdiagnosis Analysis of Endoscopic Ultrasonography in Diagnosing Subepithelial Lesions of the Upper Digestive Tract," *International Journal of Clinical and Experimental Medicine* 11, no. 11 (2018): 12181–12188.
20. R. S. Okten, S. Kacar, F. Kucukay, N. Sasmaz, and T. Cumhur, "Gastric Subepithelial Masses: Evaluation of Multidetector CT (Multiplanar Reconstruction and Virtual Gastroscopy) Versus Endoscopic Ultrasonography," *Abdominal Imaging* 37, no. 4 (2012 Aug): 519–530.
21. S. C. Reddymasu, M. Oropeza-Vail, K. Pakseresht, et al., "Are Endoscopic Ultrasonography Imaging Characteristics Reliable for the Diagnosis of Small Upper Gastrointestinal Subepithelial Lesions?," *Journal of Clinical Gastroenterology* 46, no. 1 (2012 Jan): 42–45.
22. Q. Su, J. Peng, X. Chen, Z. Xiao, R. Liu, and F. Wang, "Role of Endoscopic Ultrasonography for Differential Diagnosis of Upper Gastrointestinal Submucosal Lesions," *BMC Gastroenterology* 21, no. 1 (2021): 365.
23. Y. Minoda, E. Ihara, K. Komori, et al., "Efficacy of Endoscopic Ultrasound With Artificial Intelligence for the Diagnosis of Gastrointestinal Stromal Tumors," *Journal of Gastroenterology* 55, no. 12 (2020): 1119–1126.
24. T. W. Lim, C. W. Choi, D. H. Kang, H. W. Kim, S. B. Park, and S. J. Kim, "Endoscopic Ultrasound Without Tissue Acquisition Has Poor Accuracy for Diagnosing Gastric Subepithelial Tumors," *Medicine (Baltimore)* 95, no. 44 (2016): e5246.
25. Y. Chu, X. Qiao, X. Gao, et al., "Combined EUS and CT for Evaluating Gastrointestinal Submucosal Tumors Before Endoscopic Resection," *European Journal of Gastroenterology & Hepatology* 26, no. 8 (2014 Aug): 933–936.
26. T. Rösch, B. Kapfer, U. Will, et al., "Accuracy of Endoscopic Ultrasonography in Upper Gastrointestinal Submucosal Lesions: A Prospective Multicenter Study," *Scandinavian Journal of Gastroenterology* 37, no. 7 (2002 Jul): 856–862.
27. K. Hirai, T. Kuwahara, K. Furukawa, et al., "Artificial Intelligence-Based Diagnosis of Upper Gastrointestinal Subepithelial Lesions on Endoscopic Ultrasonography Images," *Gastric Cancer* 25, no. 2 (2022 Mar): 382–391.
28. Y. H. Yu and S. L. Gao, "Diagnosis of Gastrointestinal Mucosal and Submucosal Protuberant Lesions by Endoscopic Ultrasonography: An Analysis of 432 Cases," *Shijie Huaren Xiaohua Zazhi* 19 (2011): 1745.
29. F. Ji, Z. W. Wang, L. J. Wang, J. W. Ning, and G. Q. Xu, "Clinicopathological Characteristics of Gastrointestinal Mesenchymal Tumors and Diagnostic Value of Endoscopic Ultrasonography," *Journal of Gastroenterology and Hepatology* 23, no. 8 Pt 2 (2008 Aug): e318–e324.
30. J. H. Hwang, M. D. Saunders, S. J. Rulyak, S. Shaw, H. Nietsch, and M. B. Kimmey, "A Prospective Study Comparing Endoscopy and EUS in the Evaluation of GI Subepithelial Masses," *Gastrointestinal Endoscopy* 62, no. 2 (2005 Aug): 202–208.
31. C. W. Huh, D. H. Jung, J. S. Kim, Y. R. Shin, S. H. Choi, and B. W. Kim, "CT Versus Endoscopic Ultrasound for Differentiating Small (2–5 cm) Gastrointestinal Stromal Tumors From Leiomyomas," *AJR. American Journal of Roentgenology* 213, no. 3 (2019 Sep): 586–591.
32. C. Lefort, V. Gupta, A. Lisotti, et al., "Diagnosis of Gastric Submucosal Tumors and Estimation of Malignant Risk of GIST by Endoscopic Ultrasound. Comparison Between B Mode and Contrast-Harmonic Mode," *Digestive and Liver Disease* 53, no. 11 (2021): 1486–1491.
33. A. Săftoiu, P. Vilmann, H. Hassan, and J. G. Krag, "Utility of Colour Doppler Endoscopic Ultrasound Evaluation and Guided Therapy of Submucosal Tumours of the Upper Gastrointestinal Tract," *Ultraschall in der Medizin - European Journal of Ultrasound* 26, no. 6 (2005): 487–495.
34. B. Brand, L. Oesterhelweg, K. F. Binmoeller, et al., "Impact of Endoscopic Ultrasound for Evaluation of Submucosal Lesions in Gastrointestinal Tract," *Digestive and Liver Disease* 34, no. 4 (2002 Apr): 290–297.
35. N. Ando, H. Goto, Y. Niwa, et al., "The Diagnosis of GI Stromal Tumors With EUS-Guided Fine Needle Aspiration With Immunohistochemical Analysis," *Gastrointestinal Endoscopy* 55, no. 1 (2002 Jan): 37–43.
36. G. H. Kim, D. Y. Park, S. Kim, et al., "Is It Possible to Differentiate Gastric GISTs From Gastric Leiomyomas by EUS?," *World Journal of Gastroenterology* 15, no. 27 (2009): 3376–3381.
37. Y. T. Yang, N. Shen, F. Ao, and W. Q. Chen, "Diagnostic Value of Contrast-Enhanced Harmonic Endoscopic Ultrasonography in Predicting the Malignancy Potential of Submucosal Tumors: A Systematic Review and Meta-Analysis," *Surgical Endoscopy* 34, no. 9 (2020 Sep): 3754–3765.
38. S. H. Kim, I. K. Yoo, C. I. Kwon, S. P. Hong, and J. Y. Cho, "Utility of EUS Elastography in the Diagnosis of Gastric Subepithelial Tumors: A Pilot Study (With Video)," *Gastrointestinal Endoscopy* 91, no. 1 (2020 Jan): 172–177.e2.
39. C. Jenssen, S. Faiss, and D. Nürnberg, "Complications of Endoscopic Ultrasound and Endoscopic Ultrasound-Guided Interventions—Results

- of a Survey Among German Centers,” *Zeitschrift für Gastroenterologie* 46, no. 10 (2008): 1177–1184.
40. S. Koizumi, M. Kida, H. Yamauchi, et al., “Clinical Implications of Doubling Time of Gastrointestinal Submucosal Tumors,” *World Journal of Gastroenterology* 22, no. 45 (2016): 10015–10023.
41. S. Hata, M. Arai, T. Suzuki, et al., “Clinical Significance of Endoscopic Ultrasound for Gastric Submucosal Tumors,” *Clinics and Research in Hepatology and Gastroenterology* 37, no. 2 (2013): 207–212.
42. V. M. Kushnir, R. N. Keswani, T. G. Hollander, et al., “Compliance With Surveillance Recommendations for Foregut Subepithelial Tumors Is Poor: Results of a Prospective Multicenter Study,” *Gastrointestinal Endoscopy* 81, no. 6 (2015): 1378–1384.
43. M. L. Hu, K. L. Wu, C. S. Changchien, S. K. Chuah, and Y. C. Chiu, “Endosonographic Surveillance of 1–3 cm Gastric Submucosal Tumors Originating From Muscularis Propria,” *World Journal of Gastroenterology* 23, no. 12 (2017): 2194–2200.
44. K. H. Lok, L. Lai, H. L. Yiu, M. L. Szeto, and S. K. Leung, “Endosonographic Surveillance of Small Gastrointestinal Tumors Originating From Muscularis Propria,” *Journal of Gastrointestinal and Liver Diseases* 18, no. 2 (2009): 177–180.
45. B. Kim, S. Kang, E. Lee, et al., “Gastric Subepithelial Tumor: Long-Term Natural History and Risk Factors for Progression,” *Surgical Endoscopy* 36, no. 7 (2022): 5232–5242.
46. A. Sawaki, N. Mizuno, K. Takahashi, et al., “Long-Term Follow up of Patients With Small Gastrointestinal Stromal Tumors in the Stomach Using Endoscopic Ultrasonography–Guided Fine-Needle Aspiration Biopsy,” *Digestive Endoscopy* 18, no. 1 (2006): 40–44.
47. M. Y. Kim, H. Y. Jung, K. D. Choi, et al., “Natural History of Asymptomatic Small Gastric Subepithelial Tumors,” *Journal of Clinical Gastroenterology* 45, no. 4 (2011): 330–336.
48. Y. J. Fang, T. Y. Cheng, M. S. Sun, et al., “Suggested Cutoff Tumor Size for Management of Small EUS-Suspected Gastric Gastrointestinal Stromal Tumors,” *Journal of the Formosan Medical Association* 111, no. 2 (2012 Feb): 88–93.
49. Z. Gao, C. Wang, Q. Xue, et al., “The Cut-Off Value of Tumor Size and Appropriate Timing of Follow-Up for Management of Minimal EUS-Suspected Gastric Gastrointestinal Stromal Tumors,” *BMC Gastroenterology* 17, no. 1 (2017): 8.
50. E. Melzer and H. Fidder, “The Natural Course of Upper Gastrointestinal Submucosal Tumors: An Endoscopic Ultrasound Survey,” *Israel Medical Association Journal* 2, no. 6 (2000): 430–432.
51. M. Bruno, P. Carucci, A. Repici, et al., “The Natural History of Gastrointestinal Subepithelial Tumors Arising From Muscularis Propria: An Endoscopic Ultrasound Survey,” *Journal of Clinical Gastroenterology* 43, no. 9 (2009): 821–825.
52. K. R. S. Gill, L. Camellini, R. Conigliaro, et al., “The Natural History of Upper Gastrointestinal Subepithelial Tumors: A Multicenter Endoscopic Ultrasound Survey,” *Journal of Clinical Gastroenterology* 43, no. 8 (2009): 723–726.
53. J. Hu, X. Sun, N. Ge, et al., “The Necessity of Treatment for Small Gastric Subepithelial Tumors (1–2 cm) Originating From Muscularis Propria: An Analysis of 972 Tumors,” *BMC Gastroenterology* 22, no. 1 (2022): 182.
54. L. Zhu, S. Khan, Y. Hui, et al., “Treatment Recommendations for Small Gastric Gastrointestinal Stromal Tumors: Positive Endoscopic Resection,” *Scandinavian Journal of Gastroenterology* 54, no. 3 (2019): 297–302.
55. D. O. Faigel and S. Abulhawa, “Gastrointestinal Stromal Tumors: The Role of the Gastroenterologist in Diagnosis and Risk Stratification,” *Journal of Clinical Gastroenterology* 46, no. 8 (2012): 629–636.
56. A. Chak, M. I. Canto, T. Rösch, et al., “Endosonographic Differentiation of Benign and Malignant Stromal Cell Tumors,” *Gastrointestinal Endoscopy* 45, no. 6 (1997): 468–473.
57. K. Akahoshi, M. Oya, T. Koga, et al., “Clinical Usefulness of Endoscopic Ultrasound–Guided Fine Needle Aspiration for Gastric Subepithelial Lesions Smaller Than 2 cm,” *Journal of Gastrointestinal and Liver Diseases* 23, no. 4 (2014): 405–412.
58. H. H. Okasha, S. Elkholy, M. Sayed, et al., “Endoscopic Ultrasound and Endoscopic Ultrasound–Guided Fine Needle Aspiration in the Diagnosis of Diffuse Gastrointestinal Lesions With Inconclusive Endoscopic Biopsies,” *Turkish Journal of Gastroenterology* 28, no. 5 (2017): 370–376.
59. D. C. Joo, G. H. Kim, M. W. Lee, B. E. Lee, D. H. Beak, and G. A. Song, “Diagnostic Performance of Endoscopic Ultrasonography–Guided Fine-Needle Biopsy in Upper Gastrointestinal Subepithelial Tumors Measuring 2–5 cm in Size,” *Surgical Endoscopy* 36, no. 11 (2022): 8060–8066, <https://pubmed.ncbi.nlm.nih.gov/35441867/>.
60. V. Arantes, R. Logroño, S. Faruqi, I. Ahmed, I. Waxman, and M. S. Bhutani, “Endoscopic Sonographically Guided Fine-Needle Aspiration Yield in Submucosal Tumors of the Gastrointestinal Tract,” *Journal of Ultrasound in Medicine* 23, no. 9 (2004): 1141–1150.
61. K. M. Hoda, S. A. Rodriguez, and D. O. Faigel, “EUS-Guided Sampling of Suspected GI Stromal Tumors,” *Gastrointestinal Endoscopy* 69, no. 7 (2009): 1218–1223.
62. M. Polkowski, W. Gerke, D. Jarosz, et al., “Diagnostic Yield and Safety of Endoscopic Ultrasound–Guided Trucut Biopsy in Patients With Gastric Submucosal Tumors: A Prospective Study,” *Endoscopy* 41, no. 4 (2009): 329–334.
63. M. A. Mekky, K. Yamao, A. Sawaki, et al., “Diagnostic Utility of EUS-Guided FNA in Patients With Gastric Submucosal Tumors,” *Gastrointestinal Endoscopy* 71, no. 6 (2010): 913–919.
64. M. Philipper, S. Hollerbach, H. E. Gabbert, et al., “Prospective Comparison of Endoscopic Ultrasound–Guided Fine-Needle Aspiration and Surgical Histology in Upper Gastrointestinal Submucosal Tumors,” *Endoscopy* 42, no. 4 (2010): 300–305.
65. G. Fernández-Esparrach, O. Sendino, M. Solé, et al., “Endoscopic Ultrasound–Guided Fine-Needle Aspiration and Trucut Biopsy in the Diagnosis of Gastric Stromal Tumors: A Randomized Crossover Study,” *Endoscopy* 42, no. 4 (2010): 292–299.
66. J. DeWitt, R. E. Emerson, S. Sherman, et al., “Endoscopic Ultrasound–Guided Trucut Biopsy of Gastrointestinal Mesenchymal Tumor,” *Surgical Endoscopy* 25, no. 7 (2011): 2192–2202.
67. T. Suzuki, M. Arai, T. Matsumura, et al., “Factors Associated With Inadequate Tissue Yield in EUS-FNA for Gastric SMT,” *ISRN Gastroenterology* 2011 (2011): 619128.
68. J. H. Lee, K. D. Choi, M. Y. Kim, et al., “Clinical Impact of EUS-Guided Trucut Biopsy Results on Decision Making for Patients With Gastric Subepithelial Tumors ≥ 2 cm in Diameter,” *Gastrointestinal Endoscopy* 74, no. 5 (2011): 1010–1018.
69. R. R. Watson, K. F. Binmoeller, C. M. Hamerski, et al., “Yield and Performance Characteristics of Endoscopic Ultrasound–Guided Fine Needle Aspiration for Diagnosing Upper GI Tract Stromal Tumors,” *Digestive Diseases and Sciences* 56, no. 6 (2011): 1757–1762.
70. A. J. Eckardt, A. Adler, E. M. Gomes, et al., “Endosonographic Large-Bore Biopsy of Gastric Subepithelial Tumors: A Prospective Multicenter Study,” *European Journal of Gastroenterology & Hepatology* 24, no. 10 (2012): 1135–1144.
71. E. Çağlar, I. Hatemi, D. Atasoy, G. Sişman, and H. Sentürk, “Concordance of Endoscopic Ultrasonography–Guided Fine Needle Aspiration Diagnosis With the Final Diagnosis in Subepithelial Lesions,” *Clinical Endoscopy* 46, no. 4 (2013): 379–383.

72. G. H. Kim, Y. K. Cho, E. Y. Kim, et al., "Comparison of 22-Gauge Aspiration Needle With 22-Gauge Biopsy Needle in Endoscopic Ultrasonography-Guided Subepithelial Tumor Sampling," *Scandinavian Journal of Gastroenterology* 49, no. 3 (2014): 347–354.
73. S. Komanduri, L. Keefer, and S. Jakate, "Diagnostic Yield of a Novel Jumbo Biopsy "Unroofing" Technique for Tissue Acquisition of Gastric Submucosal Masses," *Endoscopy* 43, no. 10 (2011): 849–855.
74. J. M. Buscaglia, S. Nagula, V. Jayaraman, et al., "Diagnostic Yield and Safety of Jumbo Biopsy Forceps in Patients With Subepithelial Lesions of the Upper and Lower GI Tract," *Gastrointestinal Endoscopy* 75, no. 6 (2012): 1147–1152.
75. G. Y. Pih and D. H. Kim, "Endoscopic Ultrasound-Guided Fine Needle Aspiration and Biopsy in Gastrointestinal Subepithelial Tumors," *Clinical Endoscopy* 52, no. 4 (2019): 314–320.
76. A. Facciorusso, S. P. Sunny, V. Del Prete, M. Antonino, and N. Muscatiello, "Comparison Between Fine-Needle Biopsy and Fine-Needle Aspiration for EUS-Guided Sampling of Subepithelial Lesions: A Meta-Analysis," *Gastrointestinal Endoscopy* 91, no. 1 (2020): 14–22.e2.
77. A. Dhaliwal, S. Kolli, B. S. Dhindsa, et al., "Clinical Efficacy and Safety of Mucosal Incision-Assisted Biopsy for the Diagnosis of Upper Gastrointestinal Subepithelial Tumors: A Systematic Review and Meta-Analysis," *Annals of Gastroenterology* 33, no. 2 (2020): 155–161.
78. Y. Minoda, T. Chinen, T. Osoegawa, et al., "Superiority of Mucosal Incision-Assisted Biopsy Over Ultrasound-Guided Fine Needle Aspiration Biopsy in Diagnosing Small Gastric Subepithelial Lesions: A Propensity Score Matching Analysis," *BMC Gastroenterology* 20, no. 1 (2020): 19.
79. T. Osoegawa, Y. Minoda, E. Ihara, et al., "Mucosal Incision-Assisted Biopsy Versus Endoscopic Ultrasound-Guided Fine-Needle Aspiration With a Rapid On-Site Evaluation for Gastric Subepithelial Lesions: A Randomized Cross-Over Study," *Digestive Endoscopy* 31, no. 4 (2019): 413–421.
80. J. Park, J. C. Park, J. H. Jo, et al., "Prospective Comparative Study of Endoscopic Ultrasonography-Guided Fine-Needle Biopsy and Unroofing Biopsy," *Digestive and Liver Disease* 51, no. 6 (2019): 831–836.
81. O. Sanaei, G. Fernández-Esparrach, C. De La Serna-Higuera, et al., "EUS-Guided 22-Gauge Fine Needle Biopsy Versus Single-Incision With Needle Knife for the Diagnosis of Upper Gastrointestinal Subepithelial Lesions: A Randomized Controlled Trial," *Endoscopy International Open* 8, no. 3 (2020): E266–E273.
82. V. Zoundjiekpon, P. Falt, P. Fojtik, et al., "Endosonography-Guided Fine-Needle Aspiration Versus "Key-Hole Biopsy" in the Diagnostics of Upper Gastrointestinal Subepithelial Tumors. A Prospective Randomized Interventional Study," *Biomedical Papers of the Medical Faculty of the University Palacky, Olomouc, Czech Republic* 164, no. 1 (2020): 63–70.
83. H. Joensuu, "Risk Stratification of Patients Diagnosed With Gastrointestinal Stromal Tumor," *Human Pathology* 39, no. 10 (2008): 1411–1419.
84. K. Akahoshi, Y. Sumida, N. Matsui, et al., "Preoperative Diagnosis of Gastrointestinal Stromal Tumor by Endoscopic Ultrasound-Guided Fine Needle Aspiration," *World Journal of Gastroenterology* 13, no. 14 (2007): 2077–2082.
85. T. Attila and Ö. Aydın, "Lesion Size Determines Diagnostic Yield of EUS-FNA With Onsite Cytopathologic Evaluation for Upper Gastrointestinal Subepithelial Lesions," *Turkish Journal of Gastroenterology* 29, no. 4 (2018): 436–441.
86. T. Tamura, Y. Yamashita, K. Ueda, et al., "Rapid On-Site Evaluation by Endosonographers During Endoscopic Ultrasonography-Guided Fine-Needle Aspiration for Diagnosis of Gastrointestinal Stromal Tumors," *Clinical Endoscopy* 50, no. 4 (2017): 372–378.
87. S. Kersting, M. S. Janot-Matuschek, C. Schnitzler, D. E. Chourio Barboza, W. Uhl, and U. Mittelkötter, "GIST: Correlation of Risk Classifications and Outcome," *Journal of Medicine and Life* 15, no. 8 (2022): 932–943.
88. F. Caillol, E. Meunier, C. Zemmour, et al., "Efficiency of an Endoscopic Resection Strategy for Management of Submucosal Tumors <20 mm in the Upper Gastrointestinal Tract," *Endoscopy International Open* 10, no. 4 (2022): E347–E353.
89. M. K. Joo, J. J. Park, H. Kim, et al., "Endoscopic Versus Surgical Resection of GI Stromal Tumors in the Upper GI Tract," *Gastrointestinal Endoscopy* 83, no. 2 (2016 Feb): 318–326.
90. I. H. Jeong, J. H. Kim, S. R. Lee, et al., "Minimally Invasive Treatment of Gastric Gastrointestinal Stromal Tumors: Laparoscopic and Endoscopic Approach," *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques* 22, no. 3 (2012): 244–250.
91. X. Ren, M. Ni, T. Zhu, et al., "Comparison of Safety and Outcomes Between Endoscopic and Surgical Resections of Intermediate-Risk Primary Gastric Gastrointestinal Stromal Tumors," *Digestive Diseases* 41, no. 2 (2023): 187–197.
92. Y. Zhao, T. Pang, B. Zhang, et al., "Retrospective Comparison of Endoscopic Full-Thickness Versus Laparoscopic or Surgical Resection of Small (≤ 5 cm) Gastric Gastrointestinal Stromal Tumors," *Journal of Gastrointestinal Surgery* 24, no. 12 (2020): 2714–2721.
93. T. Pang, Y. Zhao, T. Fan, et al., "Comparison of Safety and Outcomes Between Endoscopic and Surgical Resections of Small (≤ 5 cm) Primary Gastric Gastrointestinal Stromal Tumors," *Journal of Cancer* 10, no. 17 (2019): 4132–4141.
94. L. Y. Huang, J. Cui, C. R. Wu, et al., "Endoscopic Full-Thickness Resection and Laparoscopic Surgery for Treatment of Gastric Stromal Tumors," *World Journal of Gastroenterology* 20, no. 25 (2014): 8253–8259.
95. Y. Meng, W. Li, L. Han, et al., "Long-Term Outcomes of Endoscopic Submucosal Dissection Versus Laparoscopic Resection for Gastric Stromal Tumors Less Than 2 cm," *Journal of Gastroenterology and Hepatology* 32, no. 10 (2017): 1693–1697.
96. X. Yin, Y. Yin, H. Chen, et al., "Comparison Analysis of Three Different Types of Minimally Invasive Procedures for Gastrointestinal Stromal Tumors ≤ 5 cm," *Journal of Laparoendoscopic & Advanced Surgical Techniques. Part A* 28, no. 1 (2018): 58–64.
97. C. R. Wu, L. Y. Huang, J. Guo, et al., "Clinical Control Study of Endoscopic Full-Thickness Resection and Laparoscopic Surgery in the Treatment of Gastric Tumors Arising From the Muscularis Propria," *Chinese Medical Journal* 128, no. 11 (2015): 1455–1459.
98. L. Wang, W. Ren, C. Fan, et al., "Full-Thickness Endoscopic Resection of Nonintracavitary Gastric Stromal Tumors: A Novel Approach," *Surgical Endoscopy* 25, no. 2 (2011): 641–647.
99. B. He, S. Yan, R. Li, H. Qiu, and J. Tu, "A Comparative Study of Treatment of Gastrointestinal Stromal Tumors With Laparoscopic Surgery: A Retrospective Study," *Journal of BUON* 23, no. 3 (2018): 820–825.
100. W. J. Dai, G. Liu, M. Wang, et al., "Endoscopic Versus Laparoscopic Resection of Gastric Gastrointestinal Stromal Tumors: A Multicenter Study," *Oncotarget* 8, no. 7 (2017): 11259–11267.
101. L. Chen, Q. Zhang, F. Y. Li, et al., "Comparison of Treatment Outcomes Between Laparoscopic and Endoscopic Surgeries for Relatively Small Gastric Gastrointestinal Stromal Tumors," *Surgical Oncology* 27, no. 4 (2018): 737–742.
102. H. H. Kim, "Endoscopic Treatment for Gastrointestinal Stromal Tumor: Advantages and Hurdles," *World Journal of Gastrointestinal Endoscopy* 7, no. 3 (2015): 192–205.
103. H. Y. Huang, C. F. Li, W. W. Huang, et al., "A Modification of NIH Consensus Criteria to Better Distinguish the Highly Lethal Subset of Primary Localized Gastrointestinal Stromal Tumors: A Subdivision of

- the Original High-Risk Group on the Basis of Outcome,” *Surgery* 141, no. 6 (2007): 748–756.
104. Y. Sato, S. Hashimoto, K. I. Mizuno, M. Takeuchi, and S. Terai, “Management of Gastric and Duodenal Neuroendocrine Tumors,” *World Journal of Gastroenterology* 22, no. 30 (2016): 6817–6828.
105. G. Esposito, M. Cazzato, M. Rinzivillo, et al., “Management of Type-I Gastric Neuroendocrine Neoplasms: A 10-Years Prospective Single Centre Study,” *Digestive and Liver Disease* 54, no. 7 (2022): 890–895.
106. C. S. Chung, C. L. Tsai, Y. Y. Chu, et al., “Clinical Features and Outcomes of Gastric Neuroendocrine Tumors After Endoscopic Diagnosis and Treatment: A Digestive Endoscopy Society of Taiwan (DEST),” *Medicine (Baltimore)* 97, no. 38 (2018): e12101.
107. R. A. Gladdy, V. E. Strong, D. Coit, et al., “Defining Surgical Indications for Type I Gastric Carcinoid Tumor,” *Annals of Surgical Oncology* 16, no. 11 (2009): 3154–3160.
108. S. Manfredi, T. Walter, E. Baudin, et al., “Management of Gastric Neuro-Endocrine Tumours in a Large French National Cohort (GTE),” *Endocrine* 57, no. 3 (2017): 504–511.
109. Y. Sato, H. Imamura, Y. Kaizaki, et al., “Management and Clinical Outcomes of Type I Gastric Carcinoid Patients: Retrospective, Multicenter Study in Japan,” *Digestive Endoscopy* 26, no. 3 (2014): 377–384.
110. A. Vanoli, S. La Rosa, E. Miceli, et al., “Prognostic Evaluations Tailored to Specific Gastric Neuroendocrine Neoplasms: Analysis of 200 Cases With Extended Follow-Up,” *Neuroendocrinology* 107, no. 2 (2018): 114–126.
111. T. Hirasawa, N. Yamamoto, and T. Sano, “Is Endoscopic Resection Appropriate for Type 3 Gastric Neuroendocrine Tumors? Retrospective Multicenter Study,” *Digestive Endoscopy* 33, no. 3 (2021): 408–417.
112. K. J. Lee, H. M. Kim, S. K. Lee, et al., “Type 3 Gastric Neuroendocrine Neoplasm Clinical Features: A Multicenter Study in Korea,” *Korean Society of Gastrointestinal Cancer* 5, no. 2 (2017): 86–90.
113. B. H. Min, M. Hong, J. H. Lee, et al., “Clinicopathological Features and Outcome of Type 3 Gastric Neuroendocrine Tumours,” *British Journal of Surgery* 105, no. 11 (2018 Oct): 1480–1486.
114. A. Dasari, C. Shen, D. Halperin, et al., “Trends in the Incidence, Prevalence, and Survival Outcomes in Patients With Neuroendocrine Tumors in the United States,” *JAMA Oncology* 3, no. 10 (2017): 1335–1342.
115. G. Delle Fave, D. O’Toole, A. Sundin, et al., “ENETS Consensus Guidelines Update for Gastroduodenal Neuroendocrine Neoplasms,” *Neuroendocrinology* 103, no. 2 (2016): 119–124.
116. D. Ravizza, G. Fiori, C. Trovato, et al., “Long-Term Endoscopic and Clinical Follow-Up of Untreated Type 1 Gastric Neuroendocrine Tumours,” *Digestive and Liver Disease* 39, no. 6 (2007): 537–543.
117. L. Sagatun, R. Fossmark, C. S. Jianu, et al., “Follow-Up of Patients With ECL Cell-Derived Tumours,” *Scandinavian Journal of Gastroenterology* 51, no. 11 (2016): 1398–1405.
118. D. Campana, D. Ravizza, P. Ferolla, et al., “Risk Factors of Type 1 Gastric Neuroendocrine Neoplasia in Patients With Chronic Atrophic Gastritis. A Retrospective, Multicentre Study,” *Endocrine* 56, no. 3 (2017): 633–638.
119. H. S. Lee, S. W. Jeon, G. H. Kim, et al., “Is Endoscopic Resection for Type 1 Gastric Neuroendocrine Tumors Essential for Treatment?: Multicenter, Retrospective Long-Term Follow-Up Results,” *Korean Journal of Helicobacter and Upper Gastrointestinal Research* 16, no. 1 (2016): 13.
120. K. Exarchou, H. Hu, N. A. Stephens, et al., “Endoscopic Surveillance Alone Is Feasible and Safe in Type I Gastric Neuroendocrine Neoplasms Less Than 10 mm in Diameter,” *Endocrine* 78, no. 1 (2022): 186–196.
121. H. Miyashita, T. Yamasaki, Y. Akita, et al., “Small Neuroendocrine Tumors of the Whole Gastrointestinal Tract Performed Endoscopic or Surgical Resections Also Show Positive for Lymphovascular Invasion,” *Digestion* 102, no. 6 (2021): 921–928.
122. D. M. Li, L. L. Ren, and Y. P. Jiang, “Long-Term Outcomes of Endoscopic Resection for Gastric Subepithelial Tumors,” *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques* 30, no. 2 (2020 Apr): 187–191.
123. T. Y. Duan, Y. Y. Tan, X. H. Wang, L. Lv, and D. L. Liu, “A Comparison of Submucosal Tunneling Endoscopic Resection and Endoscopic Full-Thickness Resection for Gastric Fundus Submucosal Tumors,” *Revista Española de Enfermedades Digestivas* 110, no. 3 (2018 Mar): 160–165.
124. Y. Tan, X. Tang, T. Guo, et al., “Comparison Between Submucosal Tunneling Endoscopic Resection and Endoscopic Full-Thickness Resection for Gastric Stromal Tumors Originating From the Muscularis Propria Layer,” *Surgical Endoscopy* 31, no. 8 (2017 Aug): 3376–3382.
125. J. Liu, Y. Tan, D. Liu, C. Li, M. Le, and H. Zhou, “Factors Predicting Technical Difficulties During Endoscopic Submucosal Excavation for Gastric Submucosal Tumor,” *Journal of International Medical Research* 49, no. 9 (2021): 3000605211029808.
126. C. Zeng, Y. Zhu, X. Shu, N. Lv, Q. Cai, and Y. Chen, “Endoscopic Resection of Gastric Submucosal Masses by a Dental Floss Traction Method,” *Canadian Journal of Gastroenterology & Hepatology* 2019 (2019): 1083053.
127. W. An, P. B. Sun, J. Gao, et al., “Endoscopic Submucosal Dissection for Gastric Gastrointestinal Stromal Tumors: A Retrospective Cohort Study,” *Surgical Endoscopy* 31, no. 11 (2017): 4522–4531.
128. S. Wang and L. Shen, “Efficacy of Endoscopic Submucosal Excavation for Gastrointestinal Stromal Tumors in the Cardia,” *Surgical Laparoscopy, Endoscopy & Percutaneous Techniques* 26, no. 6 (2016 Dec): 493–496.
129. F. S. Meng, Z. H. Zhang, G. D. Shan, Y. P. Chen, and F. Ji, “Endoscopic Submucosal Dissection for the Treatment of Large Gastric Submucosal Tumors Originating From the Muscularis Propria Layer: A Single Center Study,” *Zeitschrift für Gastroenterologie* 53, no. 7 (2015): 655–659.
130. Z. He, C. Sun, J. Wang, et al., “Efficacy and Safety of Endoscopic Submucosal Dissection in Treating Gastric Subepithelial Tumors Originating in the Muscularis Propria Layer: A Single-Center Study of 144 Cases,” *Scandinavian Journal of Gastroenterology* 48, no. 12 (2013): 1466–1473.
131. S. Zhang, G. Q. Chao, M. Li, G. B. Ni, and B. Lv, “Endoscopic Submucosal Dissection for Treatment of Gastric Submucosal Tumors Originating From the Muscularis Propria Layer,” *Digestive Diseases and Sciences* 58, no. 6 (2013 Jun): 1710–1716.
132. S. Y. Chun, K. O. Kim, D. S. Park, et al., “Endoscopic Submucosal Dissection as a Treatment for Gastric Subepithelial Tumors That Originate From the Muscularis Propria Layer: A Preliminary Analysis of Appropriate Indications,” *Surgical Endoscopy* 27, no. 9 (2013): 3271–3279.
133. I. D. Jeong, S. W. Jung, S. J. Bang, J. W. Shin, N. H. Park, and D. H. Kim, “Endoscopic Enucleation for Gastric Subepithelial Tumors Originating in the Muscularis Propria Layer,” *Surgical Endoscopy* 25, no. 2 (2011): 468–474.
134. I. L. Lee, P. Y. Lin, S. Y. Tung, C. H. Shen, K. L. Wei, and C. S. Wu, “Endoscopic Submucosal Dissection for the Treatment of Intraluminal Gastric Subepithelial Tumors Originating From the Muscularis Propria Layer,” *Endoscopy* 38, no. 10 (2006): 1024–1028.
135. L. Li, F. Wang, B. Wu, Q. Wang, C. Wang, and J. Liu, “Endoscopic Submucosal Dissection of Gastric Fundus Subepithelial Tumors Originating From the Muscularis Propria,” *Experimental and Therapeutic Medicine* 6, no. 2 (2013): 391–395.

136. G. Jian, L. Tan, H. Wang, et al., "Factors That Predict the Technical Difficulty During Endoscopic Full-Thickness Resection of a Gastric Submucosal Tumor," *Revista Española de Enfermedades Digestivas* 113, no. 1 (2021): 35–40.
137. Y. Li, Z. Cui, J. Yu, X. Bao, and S. Wang, "Do We Need to Conduct Full-Thickness Closure After Endoscopic Full-Thickness Resection of Gastric Submucosal Tumors?," *Turkish Journal of Gastroenterology* 31, no. 12 (2020): 942–947.
138. J. Hu, N. Ge, S. Wang, et al., "Direct Endoscopic Full-Thickness Resection for Submucosal Tumors With an Intraluminal Growth Pattern Originating From the Muscularis Propria Layer in the Gastric Fundus," *BMC Gastroenterology* 20, no. 1 (2020): 70.
139. B. Li, Q. Shi, Z. P. Qi, et al., "The Efficacy of Dental Floss and a Hemoclip as a Traction Method for the Endoscopic Full-Thickness Resection of Submucosal Tumors in the Gastric Fundus," *Surgical Endoscopy* 33, no. 11 (2019): 3864–3873.
140. J. Li, Y. Meng, S. Ye, P. Wang, and F. Liu, "Usefulness of the Thread-Traction Method in Endoscopic Full-Thickness Resection for Gastric Submucosal Tumor: A Comparative Study," *Surgical Endoscopy* 33, no. 9 (2019 Sep): 2880–2885.
141. J. Lu, T. Jiao, Y. Li, M. Zheng, and X. Lu, "Facilitating Retroflexed Endoscopic Full-Thickness Resection Through Loop-Mediated or Rope-Mediated Countertraction (With Videos)," *Gastrointestinal Endoscopy* 83, no. 1 (2016): 223–228.
142. F. Yang, S. Wang, S. Sun, et al., "Factors Associated With Endoscopic Full-Thickness Resection of Gastric Submucosal Tumors," *Surgical Endoscopy* 29, no. 12 (2015): 3588–3593.
143. L. P. Ye, Z. Yu, X. L. Mao, L. Zhu, and X. B. Zhou, "Endoscopic Full-Thickness Resection With Defect Closure Using Clips and an Endoloop for Gastric Subepithelial Tumors Arising From the Muscularis Propria," *Surgical Endoscopy* 28, no. 6 (2014): 1978–1983.
144. Y. Tan, B. Zhou, S. Zhang, et al., "Submucosal Tunneling Endoscopic Resection for Gastric Submucosal Tumors: A Comparison Between Cardia and Non-Cardia Location," *Journal of Gastrointestinal Surgery* 23, no. 11 (2019): 2129–2135.
145. Q. L. Li, W. F. Chen, C. Zhang, et al., "Clinical Impact of Submucosal Tunneling Endoscopic Resection for the Treatment of Gastric Submucosal Tumors Originating From the Muscularis Propria Layer (With Video)," *Surgical Endoscopy* 29, no. 12 (2015): 3640–3646.
146. J. Lu, T. Jiao, Y. Li, et al., "Heading Toward the Right Direction—Solution Package for Endoscopic Submucosal Tunneling Resection in the Stomach," *PLoS ONE* 10, no. 3 (2015): e0119870.
147. J. H. Noh, D. H. Kim, H. Yoon, et al., "Clinical Outcomes of Endoscopic Treatment for Type 1 Gastric Neuroendocrine Tumor," *Journal of Gastrointestinal Surgery* 25, no. 10 (2021): 2495–2502.
148. Y. H. Kwon, S. W. Jeon, G. H. Kim, et al., "Long-Term Follow Up of Endoscopic Resection for Type 3 Gastric NET," *World Journal of Gastroenterology* 19, no. 46 (2013): 8703–8708.
149. S. Stelzner, C. Antoni, G. Haroske, L. Jacobasch, J. U. Erk, and K. Ludwig, "How Should Patients With Completely Resected Gastrointestinal Stromal Tumours (GIST) Be Followed Up?," *Zentralblatt für Chirurgie* 130, no. 6 (2005 Dec): 554–561.
150. O. H. Clark, A. B. Benson, J. D. Berlin, et al., "NCCN Clinical Practice Guidelines in Oncology: Neuroendocrine Tumors," *Journal of the National Comprehensive Cancer Network* 7, no. 7 (2009): 712–747.
151. B. Landi, J. Y. Blay, S. Bonvalot, et al., "Gastrointestinal Stromal Tumours (GISTs): French Intergroup Clinical Practice Guidelines for Diagnosis, Treatments and Follow-Up (SNFGE, FFCD, GERCOR, UNICANCER, SFCD, SFED, SFRO)," *Digestive and Liver Disease* 51, no. 9 (2019): 1223–1231.
152. T. Nishida, S. Hirota, A. Yanagisawa, et al., "Clinical Practice Guidelines for Gastrointestinal Stromal Tumor (GIST) in Japan: English

Version," *International Journal of Clinical Oncology* 13, no. 5 (2008 Oct): 416–430.

Supporting Information

Additional supporting information can be found online in the Supporting Information section. **Figure S1:** PRISMA flow diagram for KQ1. **Figure S2:** PRISMA flow diagram for KQ2. **Figure S3:** PRISMA flow diagram for KQ3. **Figure S4:** PRISMA flow diagram for KQ4. **Figure S5:** Quality assessment of included studies for KQ4. **Figure S6:** PRISMA flow diagram for KQ5. **Figure S7:** PRISMA flow diagram for KQ7. **Figure S8:** Quality assessment of included studies for KQ7. **Figure S9:** Forest plot of studies comparing incomplete resection in 13 observational studies. **Figure S10:** Forest plot of studies comparing tumor recurrence in 13 observational studies. **Figure S11:** Forest plot of studies comparing postoperative complication in 13 observational studies. **Figure S12:** Forest plot of studies comparing operation time in 13 observational studies. **Figure S13:** PRISMA flow diagram for KQ8. **Figure S14:** Quality assessment of studies for KQ8. **Figure S15:** PRISMA flow diagram for KQ9. **Figure S16:** Risk of bias for KQ9. **Figure S17:** Quality assessment of included studies for KQ10. **Table S1:** Quality assessment of included studies by QUADAS-2 for KQ1. **Table S2:** Summary of findings for KQ1. **Table S3:** Quality assessment of included studies for KQ2. **Table S4:** Summary of findings for KQ2. **Table S5:** Quality assessment of included studies in KQ3. **Table S6:** Summary of findings for KQ4. **Table S7:** Quality assessment of included studies for KQ5. **Table S8:** Quality assessment of included studies for KQ9. **Table S9:** Summary of findings regarding endoscopic follow-up compared to resection for carcinoid tumor less than 10 mm. **Table S10:** Summary of findings regarding endoscopic follow-up compared to resection for carcinoid tumor less than 10 mm. **Table S11:** Summary of judgment regarding endoscopic follow-up compared to resection for carcinoid tumor less than 10 mm. **Table S12:** Quality assessment of included studies for KQ11.