



The Dynamic Brain in Overactive Bladder: Temporal Evolution of Periaqueductal Gray Connectivity Reveals Neural Rigidity

Jin Wook Kim^{1,2}

 <https://orcid.org/0000-0003-4157-9365>

¹Department of Medical Informatics, Chung-Ang University, Seoul, Korea

²Department of Urology, Chung-Ang University Gwangmyeong Hospital, Gwangmyeong, Korea

Email: jinwook@cau.ac.kr

The periaqueductal gray (PAG) stands at the crossroads of conscious control and autonomic reflexes in micturition, yet we have only begun to understand its dynamic functional architecture in health and disease. In this issue of the *International Neurourology Journal*, Chadily et al. [1] present compelling evidence that overactive bladder (OAB) syndrome involves not merely altered connectivity patterns, but a fundamental loss of neural flexibility in PAG organization. Their work using ultra-high-field 7 Tesla functional magnetic resonance imaging (MRI) reveals that while healthy individuals demonstrate dynamic reorganization of PAG connectivity over time—potentially reflecting adaptive sensory processing—OAB patients exhibit a remarkably static pattern, locked in a state of functional rigidity.

The Evolving Landscape of Neural Control

Our understanding of lower urinary tract neural control has undergone remarkable transformation over the past decade. As we previously discussed in our comprehensive review, the brain-bladder axis operates through complex, hierarchically organized circuits where the PAG serves as the critical integration center, receiving sensory information from the bladder and coordinating with higher cortical regions to determine appropriate responses [2]. The PAG exchanges bladder-related signals with multiple brain regions, functioning not as a simple relay but as an active processing hub that interprets ascending sensory information and modulates descending motor commands [3].

The current study by Chadily et al. [1] advances this understanding by examining not just static connectivity patterns but their temporal evolution. Using sophisticated clustering analysis with the Louvain module detection algorithm, they demonstrate that healthy controls show progressive changes in PAG functional organization during resting-state scanning (correlation coefficients increasing from 0.30 to 0.48), while OAB patients maintain consistently high correlations (~0.58) throughout the scanning period. This difference ($P=0.017$) suggests that the pathophysiology of OAB may involve not just altered connectivity strength but impaired dynamic flexibility.

Methodological Innovation and Technical Considerations

The use of 7 Tesla MRI represents a significant methodological advance in neurourology research. As we have emphasized in previous editorials, technological innovation drives scientific discovery, and ultra-high-field imaging provides unprecedented spatial resolution for examining small brainstem structures [4]. The PAG, with its compact anatomy and columnar organization, particularly benefits from this enhanced resolution. The authors' approach of dividing the 420-volume scan into 6 temporal blocks of 70 volumes each provides a novel window into the temporal dynamics of PAG organization, revealing patterns invisible to conventional static connectivity analyses.

However, we must acknowledge important limitations that temper our interpretation. The small sample size—only 4 OAB patients and 6 controls in the final analysis—and the significant age mismatch between groups (mean age, 61 years vs. 31



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

years) raise questions about generalizability. Age-related changes in brain connectivity are well-documented, and we cannot exclude the possibility that the observed differences reflect aging rather than OAB pathophysiology per se. Furthermore, the absence of urodynamic confirmation prevents us from definitively linking the observed neural patterns to detrusor overactivity, the presumed peripheral substrate of OAB symptoms.

Clinical Implications and the Promise of Biomarkers

Despite these limitations, the implications of this work extend beyond academic interest. The identification of reduced dynamic flexibility in PAG organization offers a potential neuroimaging biomarker for OAB—something our field has long sought. As we have noted in our discussions of future directions for neurourological research, objective biomarkers could revolutionize patient stratification and treatment selection [5]. If validated in larger cohorts, PAG connectivity patterns might predict treatment response to anticholinergics, beta-3 agonists, or neuromodulation therapies.

The concept of neural rigidity in OAB aligns with emerging understanding of the condition as a disorder of sensory processing rather than purely motor dysfunction. The static PAG organization observed in patients may reflect a system unable to adaptively process changing bladder sensations, potentially explaining the characteristic urgency that defines OAB. This perspective shifts our focus from peripheral bladder abnormalities to central neural mechanisms, opening new therapeutic avenues targeting brain plasticity rather than bladder contractility.

Integration with Existing Knowledge

These findings complement our evolving understanding of PAG function in micturition control. Previous work has established that the PAG contains distinct functional columns corresponding to different aspects of autonomic control [6]. The current study suggests these columnar organizations may not be fixed but rather exhibit dynamic reorganization in response to changing physiological states. The loss of this flexibility in OAB represents a novel pathophysiological mechanism that bridges molecular, cellular, and systems-level dysfunction.

We must also consider how these findings relate to other brain regions implicated in OAB. The PAG does not operate in isolation but forms part of an extensive network including the prefrontal cortex, anterior cingulate cortex, and insular cortex [7]. Future studies should examine whether the observed rigidity in PAG organization extends to these connected regions,

potentially revealing a broader pattern of neural inflexibility in OAB.

Future Directions and Research Priorities

Moving forward, several research priorities emerge from this work. First, we need larger, age-matched cohorts with comprehensive phenotyping including urodynamic studies, symptom questionnaires, and bladder diaries. Second, longitudinal studies examining PAG connectivity before and after successful treatment could establish whether restoration of dynamic flexibility accompanies clinical improvement. Third, investigation of PAG dynamics during actual bladder filling—not just resting state—would provide more direct insight into sensory processing abnormalities.

The technical challenges of such studies should not be underestimated. Scanning during natural bladder filling requires extended protocols that may be uncomfortable for OAB patients experiencing urgency. The use of catheterization, while ensuring standardized bladder states, introduces an artificial element that may influence neural activity. Nevertheless, these challenges must be overcome to fully understand the dynamic neural processes underlying normal and pathological bladder control.

Conclusion: Toward Precision Neurourology

The work of Chadily et al. [1] represents an important step toward precision medicine in neurourology. By revealing temporal dynamics of PAG organization previously hidden from view, they have identified a novel mechanism—neural rigidity—that may contribute to OAB pathophysiology. While replication in larger cohorts is essential, this finding opens exciting possibilities for developing objective biomarkers and targeted therapies.

As we continue to unravel the complex neurobiology of bladder control, we must embrace both technological innovation and conceptual evolution. The integration of advanced neuroimaging with systems neuroscience approaches promises to transform our understanding of lower urinary tract dysfunction. The *International Neurourology Journal* remains committed to publishing such transformative research, fostering the interdisciplinary collaboration necessary to translate these discoveries into clinical benefit for our patients.

The journey from identifying altered PAG connectivity patterns to developing targeted therapeutics will require sustained effort across multiple disciplines. Yet the potential rewards—objective diagnosis, personalized treatment selection, and novel

therapeutic targets — justify this investment. As we stand at the threshold of a new era in neurourology, studies like this remind us that the brain, in all its dynamic complexity, holds the key to understanding and treating lower urinary tract dysfunction.

• **Conflict of Interest:** No potential conflict of interest relevant to this article was reported.

REFERENCES

1. Chadily SF, van Houtum NR, van Klaveren ME, Knops A, van Koeveeringe GA, de Rijk MM, et al. Altered periaqueductal gray functional connectivity in overactive bladder patients: a resting-state functional magnetic resonance imaging study using clustering analysis. *Int Neurourol J* 2025;29:215-22.
2. Kim JW, Kim SJ, Park JM, Na YG, Kim KH. Past, present, and future in the study of neural control of the lower urinary tract. *Int Neurourol J* 2020;24:191-9.
3. Lee CL, Lee J, Park JM, Na HS, Shin JH, Na YG, et al. Sophisticated regulation of micturition: review of basic neurourology. *J Exerc Rehabil* 2021;17:295-307.
4. Kim JW. Diffusion tensor imaging: the high-resolution image of functionality in the central nervous system. *Int Neurourol J* 2022; 26:171-2.
5. Cho YS. Future directions for neurourological research. *Int Neurourol J* 2020;24:189-90.
6. Fernández Chadily S, de Rijk MM, Janssen JM, van den Hurk J, van Koeveeringe GA. Assessment of brainstem functional organization in healthy adults and overactive bladder patients using ultra-high field fMRI. *Biomedicines* 2023;11:403.
7. de Rijk MM, van den Hurk J, Rahnama'i MS, van Koeveeringe GA. Parcellation of human periaqueductal gray at 7-T fMRI in full and empty bladder state: the foundation to study dynamic connectivity changes related to lower urinary tract functioning. *Neurourol Urodyn* 2021;40:616-23.