



Erratum

Erratum to: Phytochemical analysis and antioxidant properties of mixtures containing *Platycodonis Radix*, *Codonopsis Radix*, *Paeoniae Radix*, and *Lonicerae Flos*

Hak-Dong Lee^{1,2} · Neil Patrick Uy¹ · Wang Kwon Kim³ · Sanghyun Lee^{1,2} 

Published Online: 17 July 2024

© The Korean Society for Applied Biological Chemistry 2024

Erratum to: J Appl Biol Chem (2024) 67:23, 165–173
<https://doi.org/10.3839/jabc.2024.023>

Add the following to the original.
p. 166

Sample preparation and steam treatment

Two sample types, one subjected to steam treatment (*Platycodonis Radix* and *Codonopsis Radix*) and the other without steam treatment (*Paeoniae Radix* and *Lonicerae Flos*), were prepared by Korea Best One Co., Republic of Korea (Fig. 1). The steamed *Platycodonis Radix* (SPR), steamed *Codonopsis Radix* (SCR), non-steamed *Paeoniae Radix* (NPR), and non-steamed *Lonicerae Flos* (NLF) were ground and extracted with water using a reflux system. The first extraction was performed with water at 15 times the weight of the raw materials (100 °C, 8 h). After filtering and vacuum concentrating the first extract, water at 10 times the weight of the initially extracted raw materials was added for the second extraction (100 °C, 8 h). After filtering and concentrating the second extract, the final Brix degree was approximately 40.

Four mixtures were prepared according to the ratios of the raw materials used for extraction: mixtures of SPR, SCR, and NPR (at ratios of 1:1:1 and 2:2:1) and mixtures of SPR, SCR, and NLF (at ratios of 1:1:1 and 2:2:1) were provided by Korea Best One Co., Republic of Korea. The extracts were dissolved in 80% MeOH (50 mg/mL) and filtered through a syringe filter (PVDF, 0.22 µm). Stock solutions of the standards were prepared by dissolution in 80% MeOH. Working solutions were prepared by diluting the stock solution to the desired concentrations for the standard calibration curves. The condition of steam treatment of SPR and SCR is shown at Table 1.

Hak-Dong Lee and Neil Patrick Uy equally contributed to this article.

Sanghyun Lee (✉)
E-mail: slee@cau.ac.kr

¹Department of Plant Science and Technology, Chung-Ang University, Anseong 17546, Republic of Korea

²Natural Product Institute of Science and Technology, Anseong 17546, Republic of Korea

³Korea Best One Corp., Chuncheon 24341, Republic of Korea

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.