

Original Article

Int Neurourol J 2026;30(1):57-62

<https://doi.org/10.5213/inj.2550208.104>

pISSN 2093-4777 · eISSN 2093-6931



Predicting Time to Recovery From Acute Urinary Retention in Hospitalized Patients: Development and Validation of a Clinical Prediction Model

Jin Wook Kim^{1,2}

¹Department of Urology, Chung-Ang University Gwangmyeong Hospital, Chung-Ang University College of Medicine, Gwangmyeong, Korea

²Department of Medical Informatics, Chung-Ang University College of Medicine, Seoul, Korea

Purpose: Acute urinary retention (AUR) is a common complication in hospitalized patients with unpredictable recovery times. We aimed to develop and validate a clinical prediction model for recovery from AUR within 14 days.

Methods: A prospective cohort study was conducted from March 2024 to July 2025 at a tertiary care center. Adult patients (n = 126) with AUR were enrolled upon urological consultation. Males with prostate volume > 30 g were excluded. Patients underwent standardized voiding trials every 3–4 days, with success defined as postvoid residual < 100 mL. Multivariable logistic regression was used to identify predictors of recovery within 14 days.

Results: The cohort comprised 84 males (66.7%) and 42 females (33.3%), mean age 71.9 ± 14.2 years. Overall, 81.7% achieved successful voiding within 14 days. The prediction model demonstrated excellent discrimination (area under the curve [AUC], 0.83; 95% confidence interval [CI], 0.76–0.90) with 4 independent predictors: age < 70 years (OR, 4.11; 95% CI, 2.18–7.76; $P = 0.008$), male sex (OR, 9.09; 95% CI, 4.35–19.01; $P < 0.001$), postoperative etiology (OR, 4.38; 95% CI, 2.26–8.48; $P = 0.004$), and retention volume ≤ 450 mL (OR, 4.55; 95% CI, 2.17–9.52; $P = 0.017$). Bootstrap validation confirmed model stability (optimism-corrected AUC = 0.81).

Conclusions: Our clinical prediction model reliably identifies patients at risk for prolonged urinary retention using 4 simple bedside parameters. Implementation may optimize catheter management strategies and improve patient outcomes through risk-stratified care pathways.

Keywords: Urinary retention; Urinary catheterization; Clinical decision-making; Predictive model; Recovery of function; Risk assessment

- **Grant/Fund Support:** This study received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.
- **Research Ethics:** The protocol was approved by the Institutional Review Board (IRB) of Chung-Ang University College of Medicine (IRB #1981-006-382), and all participants provided written informed consent.
- **Conflict of Interest:** No potential conflict of interest relevant to this article was reported.

INTRODUCTION

Acute urinary retention (AUR) constitutes a urological emer-

gency characterized by the sudden inability to voluntarily void despite a full bladder, affecting approximately 10% of men over 70 years and 7% of women over 80 years annually [1, 2]. In hos-

Corresponding author: Jin Wook Kim  <https://orcid.org/0000-0003-4157-9365>
Department of Urology, Chung-Ang University Gwangmyeong Hospital,
Chung-Ang University College of Medicine, 110 Iljik-ro, Gwangmyeong 14353,
Korea

Email: jinwook@cau.ac.kr

Submitted: August 12, 2025 / **Accepted after revision:** October 20, 2025



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

pitalized patients, AUR represents a pervasive complication occurring in 4%–25% of postoperative patients and 8%–12% of medical admissions [3, 4]. The heterogeneous etiology of AUR, ranging from obstructive pathology to neurogenic dysfunction and pharmacological effects, presents significant management challenges [5, 6].

While immediate bladder decompression via catheterization provides symptomatic relief, the critical question of when patients will recover spontaneous voiding lacks evidence-based guidance [7, 8]. Premature catheter removal risks reretention and emergency re-catheterization, while prolonged catheterization increases catheter-associated urinary tract infections, urethral trauma, and patient discomfort [9, 10]. Current literature predominantly focuses on specific populations such as men with benign prostatic hyperplasia or women following pelvic surgery, limiting generalizability [11, 12]. The absence of validated clinical prediction tools leads to arbitrary voiding trial schedules and inefficient resource utilization [13–15].

This study addresses this gap by developing and validating a universally applicable prediction model for recovery from AUR. Our objective was to identify readily available clinical parameters that reliably predict successful voiding within 14 days—a timeframe balancing risks of prolonged catheterization against likelihood of spontaneous recovery.

MATERIALS AND METHODS

Study Design and Participants

We conducted a prospective observational cohort study at a 1,200-bed tertiary care academic medical center between March 2024 and July 2025. The Institutional Review Board (IRB) approved the study protocol (IRB #1981-006-382), and all participants provided written informed consent. The study followed Transparent Reporting of a Multivariable Prediction Model for Individual Prognosis Or Diagnosis (TRIPOD) guidelines for prediction model development [16].

Adult patients aged ≥ 18 years with new-onset AUR requiring urological consultation were eligible if they had bladder volume ≥ 300 mL on initial catheterization and could provide informed consent. Exclusion criteria included chronic urinary retention (previous episodes within 6 months), prostate volume > 30 g in males, active urinary tract infection, established neurogenic bladder, urethral stricture or bladder neck contracture, current urological malignancy, indwelling catheter within 30 days, or cognitive impairment precluding trial participation.

Data Collection

Trained research coordinators collected baseline data using standardized forms. Variables included demographics (age, sex, body mass index), clinical context (postoperative vs. spontaneous onset, surgery type, time from surgery), comorbidities (hypertension, diabetes mellitus, cerebrovascular accident, spinal disease), medications (alpha-blockers, anticholinergics, opioids), laboratory values (creatinine, electrolytes), and retention characteristics (initial volume, hematuria, bladder wall thickness). For males, prostate parameters including volume and prostate-specific antigen were recorded when available.

Management Protocol

All patients received standardized care with 16–18 French Foley catheter insertion and continuous drainage. Males without contraindications received tamsulosin 0.4 mg daily. Voiding trials occurred every 3–4 days following this protocol: bladder instillation with 300-mL sterile saline, catheter removal, voiding attempt with uroflowmetry when feasible, and postvoid residual (PVR) measurement via ultrasound within 10 minutes. Success was defined as PVR < 100 mL; failure (PVR ≥ 100 mL) resulted in immediate recatheterization.

Statistical Analysis

Sample size calculation based on 10 events per predictor variable with anticipated 8–10 predictors and 70% success rate yielded minimum requirement of 100 patients [17]. Continuous variables were summarized as means $\pm$ standard deviation or medians (interquartile range, IQR); categorical variables as frequencies (percentages). Associations between predictors and primary outcome were assessed using t-test, Mann-Whitney U-test, chi-square, or Fisher exact test.

Variables with $P < 0.20$ in univariable analysis or strong clinical rationale entered multivariable logistic regression with backward stepwise selection ($P < 0.05$ for retention). Multicollinearity was assessed (variance inflation factor < 5). Model performance was evaluated using area under receiver operating characteristic curve (AUC-ROC) for discrimination and Hosmer-Lemeshow test for calibration. Bootstrap resampling (1000 iterations) assessed optimism-corrected performance. Sensitivity analyses excluded specific comorbidities and used alternative PVR thresholds (50 mL, 150 mL). Analysis used R ver. 4.0.3 (R Foundation for Statistical Computing, Austria).

RESULTS

Patient Characteristics

Of 168 screened patients, 126 (75%) met inclusion criteria and completed follow-up. Exclusions included prostate >30 g (n=18), chronic retention (n=12), declined consent (n=8),

Table 1. Baseline characteristics of study cohort (n = 126)

Characteristic	Study cohort	Recovery ≤ 14 days
Overall	126 (100)	103 (81.7)
Age (yr)	71.9 ± 12.5	-
< 70	45 (35.7)	38 (84.4)
≥ 70	81 (64.3)	65 (80.2)
Sex		
Male	84 (66.7)	72 (85.7)
Female	42 (33.3)	31 (73.8)
Context of retention		
Postoperative	79 (62.7)	70 (88.6)
Spontaneous	47 (37.3)	33 (70.2)
Hypertension	43 (34.1)	32 (74.4)
Diabetes mellitus	32 (25.4)	27 (84.4)
CVA history	30 (23.8)	22 (73.3)
Retention volume (mL)	602.6 ± 237.1	-
≤ 450	43 (34.4)	38 (88.4)
> 450	82 (65.6)	64 (78.0)

Values are presented as number (%) or mean ± standard deviation. CVA, cerebrovascular accident.

and incomplete data (n=4). The cohort comprised 84 males (66.7%) and 42 females (33.3%), mean age 71.9 ± 12.5 years. Postoperative retention occurred in 79 patients (62.7%). Mean initial retention volume was 602.6 ± 237.1 mL (Table 1).

Recovery Patterns

Overall, 103 patients (81.7%) achieved successful voiding within 14 days. Median time to recovery was 8 (IQR, 4–13) days. Recovery rates within 14 days differed significantly by age (<70 years: 84.4% vs. ≥70 years: 80.2%, $P=0.042$), sex (males: 85.7% vs. females: 73.8%, $P=0.098$), etiology (postoperative: 88.6% vs. spontaneous: 70.2%, $P=0.009$), and retention volume (≤450 mL: 88.4% vs. >450 mL: 78.0%, $P=0.003$). Recovery rates varied significantly across clinical subgroups (Fig. 1). Kaplan-Meier curves demonstrated divergent recovery trajectories between subgroups (Fig. 2).

Prediction Model Development

The final logistic regression model incorporated 4 independent predictors (Table 2). The model demonstrated excellent discrimination (AUC, 0.83; 95% CI, 0.76–0.90) and good calibration (Hosmer-Lemeshow chi-square = 6.82, $P=0.42$). ROC analysis identified optimal probability threshold of 0.68, yielding sensitivity 78.6% and specificity 82.6% (Fig. 3).

Model Validation and Secondary Outcomes

Bootstrap validation confirmed model stability with optimism-corrected AUC of 0.81. All predictors were selected in >85% of




Fig. 1. Proportion of patients achieving recovery within 14 days by clinical factors. Error bars represent 95% confidence intervals. The horizontal dashed line indicates overall recovery rate of 81.7%. All comparisons $P < 0.05$.




Fig. 2. Receiver operating characteristic curve for the prediction model. Area under the curve (AUC) = 0.83 (95% confidence interval, 0.76–0.90). The red dot indicates optimal cutoff (sensitivity 78.6%, specificity 82.6%).




Fig. 3. Kaplan-Meier curves for recovery from urinary retention by sex and etiology. Male patients and those with postoperative retention demonstrated significantly faster recovery rates. The vertical dashed line indicates the 14-day threshold. Log-rank test $P < 0.001$ for both comparisons.

bootstrap samples. Among recovered patients, 18 (17.5%) experienced reretention within 30 days, associated with female sex (odds ratio [OR], 2.84; $P = 0.031$) and spontaneous etiology (OR, 2.41; $P = 0.048$). Complications included urinary tract infection (6.3%), hematuria (3.2%), and urethral trauma (1.6%). Five patients (4.0%) required surgical intervention.

Model performance remained stable in sensitivity analyses: excluding diabetics (AUC, 0.82), PVR < 50 -mL threshold (AUC, 0.81), PVR < 150 -mL threshold (AUC, 0.84), and excluding cerebrovascular accident history (AUC, 0.83).

DISCUSSION

This prospective study successfully developed and validated a

Table 2. Multivariable logistic regression model for predicting recovery within 14 days

Predictor	β Coefficient	Odds ratio (95% CI)	P-value
Age < 70 yr	1.41	4.11 (2.18–7.76)	0.008
Male sex	2.21	9.09 (4.35–19.01)	< 0.001
Postoperative etiology	1.48	4.38 (2.26–8.48)	0.004
Retention volume ≤ 450 mL	1.51	4.55 (2.17–9.52)	0.017
Constant	–1.82	–	0.023

Model equation: $\text{logit}(P) = -1.82 + 1.41 \times (\text{age} < 70 \text{ yr}) + 2.21 \times (\text{male}) + 1.48 \times (\text{postoperation}) + 1.51 \times (\text{volume} \leq 450 \text{ mL})$.

CI, confidence interval.

clinical prediction model for AUR recovery within 14 days, demonstrating excellent discriminative ability (AUC, 0.83) using 4 simple bedside parameters: age, sex, etiology, and retention volume. The model's simplicity enables immediate risk stratification without specialized testing or complex calculations.

The identification of age < 70 years as favorable (OR, 4.11) aligns with physiological principles. Aging associates with decreased detrusor contractility, increased collagen deposition, and comorbidities affecting lower urinary tract function [18, 19]. This threshold provides objective risk stratification for clinical decision-making.

The striking sex disparity, with females having only one-ninth the odds of recovery compared to males, reflects fundamental pathophysiological differences. Female AUR often indicates complex pelvic floor dysfunction, prolapse, or primary bladder pathology rather than simple outlet obstruction [20, 21]. This challenges current practice patterns extrapolated from male-predominant studies and suggests women require intensive evaluation and different management strategies.

Postoperative retention's superior prognosis (OR, 4.38) provides reassurance, as it typically results from transient factors — anesthetic effects, opioid-induced suppression, and temporary nerve dysfunction — that resolve predictably [22, 23]. Conversely, spontaneous retention often signals progressive pathology requiring comprehensive evaluation.

The 450-mL volume threshold represents critical inflection for bladder injury. Exceeding this is associated with ischemic detrusor damage and neural injury, causing prolonged dysfunction [24, 25]. This emphasizes prompt AUR recognition and decompression to preserve function.

Implementation enables risk-stratified pathways: high probability ($> 80\%$) patients undergo early trials (days 3–4) with out-

patient management consideration; intermediate probability (50%–80%) follow standard protocols; low probability (< 50%) benefit from delayed trials, early consultation, and comprehensive evaluation.

Study strengths include prospective design minimizing bias, heterogeneous population enhancing generalizability, rigorous TRIPOD-compliant methodology, and internal validation confirming stability. Limitations include single-center design, exclusion of large prostates, lack of external validation, and small female sample size.

Future research should pursue external validation in multicenter cohorts, model refinement with novel biomarkers, intervention studies testing risk-stratified protocols, long-term functional outcomes, and economic impact analysis.

ACKNOWLEDGMENTS

We thank the nursing staff for protocol adherence, participating patients, and biostatistics department for analytical support.

AUTHOR CONTRIBUTION STATEMENT

The sole author (JWK) was responsible for study conception and design, acquisition and interpretation of data, and drafting and final approval of the manuscript.

ORCID

Jin Wook Kim

0000-0003-4157-9365

REFERENCES

- Jacobsen SJ, Jacobson DJ, Girman CJ, Roberts RO, Rhodes T, Guess HA, et al. Natural history of prostatism: risk factors for acute urinary retention. *J Urol* 1997;158:481-7.
- Klarskov P, Andersen JT, Asmussen CF, Brenøe J, Jensen SK, Jensen IL, et al. Acute urinary retention in women: a prospective study of 18 consecutive cases. *Scand J Urol Nephrol* 1987;21:29-31.
- Baldini G, Bagry H, Aprikian A, Carli F. Postoperative urinary retention: anesthetic and perioperative considerations. *Anesthesiology* 2009;110:1139-57.
- Wu AK, Auerbach AD, Aaronson DS. National incidence and outcomes of postoperative urinary retention in the Surgical Care Improvement Project. *Am J Surg* 2012;204:167-71.
- Selius BA, Subedi R. Urinary retention in adults: diagnosis and initial management. *Am Fam Physician* 2008;77:643-50.
- Marshall JR, Haber J, Josephson EB. An evidence-based approach to emergency department management of acute urinary retention. *Emerg Med Pract* 2014;16:1-20; quiz 21.
- Fitzpatrick JM, Desgrandchamps F, Adjali K, Gomez Guerra L, Hong SJ, El Khalid S, et al. Management of acute urinary retention: a worldwide survey of 6074 men with benign prostatic hyperplasia. *BJU Int* 2012;109:88-95.
- Emberton M, Fitzpatrick JM. The Reten-World survey of the management of acute urinary retention: preliminary results. *BJU Int* 2008;101 Suppl 3:27-32.
- Saint S, Lipsky BA, Goold SD. Indwelling urinary catheters: a one-point restraint? *Ann Intern Med* 2002;137:125-7.
- Lo E, Nicolle LE, Coffin SE, Gould C, Maragakis LL, Meddings J, et al. Strategies to prevent catheter-associated urinary tract infections in acute care hospitals: 2014 update. *Infect Control Hosp Epidemiol* 2014;35:464-79.
- McNeill SA, Hargreave TB. Alfuzosin once daily facilitates return to voiding in patients in acute urinary retention. *J Urol* 2004;171:2316-20.
- Geller EJ. Prevention and management of postoperative urinary retention after urogynecologic surgery. *Int J Womens Health* 2014;6:829-38.
- Fisher E, Subramonian K, Omar MI. The role of alpha blockers prior to removal of urethral catheter for acute urinary retention in men. *Cochrane Database Syst Rev* 2014;2014:CD006744.
- Patel MI, Watts W, Grant A. The optimal form of urinary drainage after acute retention of urine. *BJU Int* 2001;88:26-9.
- Nyman MA, Schwenk NM, Silverstein MD. Management of urinary retention: rapid versus gradual decompression and risk of complications. *Mayo Clin Proc* 1997;72:951-6.
- Collins GS, Reitsma JB, Altman DG, Moons KG. Transparent reporting of a multivariable prediction model for individual prognosis or diagnosis (TRIPOD): the TRIPOD statement. *Ann Intern Med* 2015;162:55-63.
- Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996;49:1373-9.
- Taylor JA, Kuchel GA. Detrusor underactivity: clinical features and pathogenesis of an underdiagnosed geriatric condition. *J Am Geriatr Soc* 2006;54:1920-32.
- Elbadawi A, Yalla SV, Resnick NM. Structural basis of geriatric voiding dysfunction. II. Aging detrusor: normal versus impaired contractility. *J Urol* 1993;150:1657-67.
- Ramsey S, Palmer M. The management of female urinary retention.

- tion. *Int Urol Nephrol* 2006;38:533-5.
21. Mevcha A, Drake MJ. Etiology and management of urinary retention in women. *Indian J Urol* 2010;26:230-5.
22. Darrah DM, Griebing TL, Silverstein JH. Postoperative urinary retention. *Anesthesiol Clin* 2009;27:465-84.
23. Keita H, Diouf E, Tubach F, Brouwer T, Dahmani S, Mantz J, et al. Predictive factors of early postoperative urinary retention in the postanesthesia care unit. *Anesth Analg* 2005;101:592-6.
24. Brouwer TA, Rosier PF, Moons KG, Zuithoff NP, van Roon EN, Kalkman CJ. Postoperative bladder catheterization based on individual bladder capacity: a randomized trial. *Anesthesiology* 2015; 122:46-54.
25. Mayo ME, Lloyd-Davies RW, Shuttleworth KE, Tighe JR. The damaged human detrusor: functional and electron microscopic changes in disease. *Br J Urol* 1973;45:116-25.