Journal of Animal Science and Biotechnology

RESEARCH

Open Access

Tissue-specific epigenetic regulation of fat metabolism in pigs through integrated analysis of DNA methylation and gene expression networks

Do-Young Kim[1], Byeonghwi Lim[2], Rajesh Kumar Pathak[3], Woncheoul Park[4], Jong-Eun Park[5*] and Jun-Mo Kim[1*]

Abstract

Background Fat metabolism in pigs is controlled by tissue-specific molecular mechanisms that ultimately affect growth performance and meat quality. Understanding how epigenetic modifications interact with gene expression across key metabolic and fat-depositing tissues is essential for identifying regulatory processes and potential biomarkers to improve pork quality traits. Therefore, this study aimed to elucidate tissue specific epigenetic regulation of fat metabolism by integrating DNA methylation and gene expression profiles from liver, backfat, and loin (longissimus dorsi) tissues at two physiologically developmental stages (10 and 26 weeks), representing the early post-weaning growth phase and near-market weight, respectively. By explicitly comparing these ages and tissues, the study was designed to capture the transition from muscle-dominated growth to increased lipid deposition and to identify tissue- and stage-specific regulatory signatures that may serve as biomarkers for pork quality.

Results Genome-wide DNA methylation exhibited weak clustering by tissue, whereas gene expression showed clear tissue separation. The liver harbored fewer genes with differential methylation across stage and tissue but a greater number of genes with differential expression than backfat and loin, suggesting distinct regulatory modes. Integrative analysis of the overlap genes between methylation and expression signals highlighted epigenetically mediated regulation of extracellular matrix organization, lipid metabolism, and muscle development pathways. Furthermore, weighted gene co-expression network analysis revealed distinct tissue-specific correlations between co-methylated and co-expressed modules, with enrichment in cholesterol biosynthesis, muscle contractility, and extracellular matrix remodeling. Together, these findings suggest that methylation changes are more subtle than transcriptional shifts, yet they are aligned with key functional pathways, consistent with a role for methylation as a fine-tuning mechanism that shapes tissue-specific transcriptional networks during growth.

Conclusions Across liver, backfat, and loin, DNA methylation modulates transcriptional programs in a tissue-dependent manner, prioritizing pathways central to lipid handling, extracellular matrix remodeling, and muscle function. This integrated multi-omics framework highlights candidate epigenetic markers and regulatory modules with potential utility for improving pork quality traits through selection or management strategies.

*Correspondence:
Jong-Eun Park
jepark@jejunu.ac.kr
Jun-Mo Kim
junmokim@cau.ac.kr
Full list of author information is available at the end of the article

© The Author(s) 2026. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativecommons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Keywords DNA methylation, Epigenetic, Gene expression, Multi-omics integration analysis, Porcine

Background

The pig industry plays a vital role in global livestock production, providing a reliable source of animal protein and making a significant contribution to agricultural economies. Modern pig production aims to strike a balance between rapid growth performance and consistent meat quality, thereby ensuring economic efficiency and consumer satisfaction [1]. Genetic improvement through selective breeding and crossbreeding has resulted in substantial advancements in growth rate, feed efficiency, and carcass traits in commercial pigs [2, 3]. However, meat quality is influenced by more than genetics alone; environmental factors, such as feeding strategies, nutritional content, and rearing conditions, also play essential roles in shaping pork characteristics by affecting intramuscular fat (IMF) deposition and fat distribution [4, 5]. These factors play crucial roles in shaping meat quality by influencing body fat distribution and intramuscular fat content, highlighting the importance of both genetic and environmental factors in determining meat quality.

In pigs, the liver, backfat, and loin (longissimus dorsi) tissues are intimately involved in lipid metabolism and play crucial roles in meat quality development [6–9]. De novo fatty acid synthesis occurs primarily in adipose tissue, especially in backfat, which serves as a major site for lipid storage and acts as an endocrine organ through the secretion of adipokines and other bioactive molecules [10, 11]. The liver plays a central regulatory role in whole-body lipid metabolism by coordinating fatty acid uptake, triglyceride synthesis, and lipoprotein (VLDL) secretion, thereby influencing fat deposition in peripheral tissues such as backfat and muscle [12–14]. The loin (longissimus muscle) contributes to meat quality traits such as tenderness, marbling, and flavor, due to its intramuscular fat content and muscle fiber composition [4, 15, 16]. Understanding how these tissues interact to regulate fat metabolism at different developmental stages is therefore crucial for elucidating the mechanisms underlying growth and pork quality [17–20].

In addition to transcriptomic analysis, the study of epigenetic mechanisms, such as DNA methylation, has become a crucial area of research for understanding the regulation of various metabolic diseases [21–24]. Epigenetic modifications are linked to major lipid and fat metabolic pathways that control fat accumulation and influence the expression of genes involved in lipogenesis, lipolysis, and energy homeostasis [25–29]. Integrating epigenetic research with multi-omics data, such as the epigenome and transcriptome, provides a comprehensive understanding of the mechanisms regulating gene expression and how these modifications influence metabolic processes [30, 31]. For example, a study investigating adipocyte fractions from retroperitoneal adipose tissue in obese pigs identified differentially methylated regions linked to metabolic signaling pathways, including mitogen-activated protein kinase (MAPK) and insulin signaling, shedding light on the epigenetic regulation of fat metabolism [32]. These studies have demonstrated the utility of integrating epigenetics with multi-omics approaches to deepen our understanding of metabolic regulation in livestock.

Here, we tested the hypothesis that tissue-specific DNA methylation functions as a fine-tuning mechanism coordinating transcriptional networks underlying lipid metabolism in the liver, backfat, and loin during growth. To capture the dynamics of this regulatory layer, two critical developmental stages were selected based on physiological relevance: 10 weeks, representing the early post-weaning growth phase characterized by rapid skeletal muscle development and establishment of adipose tissue depots [33–35], and 26 weeks corresponding to near market weight when fat deposition patterns are well established [36–40]. This developmental window encompasses the transition from muscle-dominated growth to enhanced lipid accumulation, providing an opportunity to identify both conserved epigenetic signatures and developmentally dynamic regulatory changes [4, 29]. To address this hypothesis, we integrated methylated DNA-binding domain sequencing (MBD-seq) and RNA sequencing (RNA-seq) data from crossbred pigs (Korean native pig × Yorkshire) at 10 and 26 weeks. Specifically, we aimed to (i) compare tissue- and stage-specific methylation and gene expression patterns; (ii) identify genes showing both differential methylation and expression (DM–DE overlap); and (iii) construct co-methylation and co-expression modules to uncover pathways associated with fat metabolism and pork quality. This integrative approach enhances our understanding of context-dependent epigenetic regulation and identifies candidate regulatory networks and markers relevant to meat quality improvement.

Methods

Animal materials and ethics statement

This study included a total of 16 crossbred pigs (Korean native pig × Yorkshire, F1 generation), as described in previous studies [41–43]. All experimental procedures

were approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science, South Korea (NIAS2016-848). Pigs had unrestricted access to feed and water, and the diet composition is detailed in previous studies [41]. The pigs were randomly assigned to two groups: 10 weeks ($n=8$; 4 males, 4 females) representing the early post-weaning period, and 26 weeks ($n=8$; 4 males, 4 females) representing an average body weight at slaughter of approximately 90 kg. The animals were sacrificed at the end of the growth stage (10 and 26 weeks) and samples were collected. All tissues were collected by an experienced technician to minimize variation. For each animal, samples of liver, backfat, and loin (longissimus dorsi) were obtained from anatomically consistent locations as follows: liver from the left lateral lobe, backfat from the dorsal midline directly above the eighth rib, and loin muscle from the central part of the longissimus dorsi located between the tenth and eleventh thoracic vertebrae. Tissue samples were consistently collected across all animals, cut into 1 cm^3 pieces, snap-frozen in liquid nitrogen, and stored at −80 °C.

Sample extraction, library construction, and sequencing

Whole-genome methylation analysis was conducted using MBD-seq, whereas RNA-seq was used for transcriptomic analysis. For MBD-seq, genomic DNA was quantified from tissue samples using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA, USA). Then, genomic DNA (1 μg) was fragmented and enriched for methylated regions using MBD proteins with a MethylMiner Methylated DNA Enrichment Kit (Invitrogen, Carlsbad, CA, USA). The enriched methylated DNA was processed into libraries using a TruSeq Nano DNA Library Prep kit (Illumina, San Diego, CA, USA). Total RNA was extracted from tissue samples using TRIzol reagent (Invitrogen), and libraries were prepared for RNA-seq. Sequencing was performed on an Illumina HiSeq 4000 platform, with thorough library quantification and quality control prior to sequencing.

Sequencing data processing and analysis

For both MBD-seq and RNA-seq analyses, the quality of raw sequencing data was initially assessed using FastQC (v0.12.1) [44]. Subsequent data cleaning involved removing low-quality reads and adapter sequences using TrimGalore (v0.6.6) [45] for MBD-seq data and Trimmomatic (v0.38) [46] for RNA-seq data. After quality control, both datasets were aligned to the Sscrofa11.1 reference genome. Bowtie2 (v3.5.1) was used for MBD-seq alignment [47], and Hisat2 (v2.1.0) was used for RNA-seq alignment [48]. Samtools (v1.9) was used for file format conversions for both datasets, including duplicate removal from the MBD-seq data using the markup function [49]. For RNA-seq analysis, gene quantification was performed using feature counts (v2.0.0) [50].

For MBD-seq, MEDIPS and edgeR were used to identify the differentially methylated regions (DMRs) [51, 52]. The genome was divided into 500 bp windows, and differential methylation between the two stages was computed using fold change (FC) values and *P*-values derived from edgeR. Regions meeting the absolute $\log_2FC \geq 1$ and *P*-value < 0.001 criteria were considered significant DMRs. These DMRs were annotated to gene features such as exons, introns, and promoters, with promoters defined as 2,000 bp upstream of the transcription start site. Differentially methylated genes (DMGs) were identified based on the proportion of DMR coverage across features. For RNA-seq, gene expression data were normalized using the trimmed mean of M values (TMM) method in EdgeR. Differentially expressed genes (DEGs) were identified using the absolute $\log_2FC \geq 1$ and a false discovery rate < 0.05 criteria, based on the Benjamini–Hochberg multiple testing correction. Samples with significant discrepancies in library size, despite TMM normalization, were excluded from the final DMG and DEG analyses to ensure the robustness of the results.

To compare the DMGs and DEGs among the three tissues, we used Venn diagrams and upset plots in the R package to visualize the overlap between each omics gene set. For functional enrichment, we used the database annotation visualization integrated discovery (DAVID) web tool to perform Gene Ontology (GO) and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway analyses to identify biological processes, molecular functions, cellular components, and enriched pathways [53–55]. A *P*-value threshold of 0.01 was used to indicate the significance of functional annotation selection criteria.

Integrated DNA methylome and transcriptome analysis

We performed integrated analysis of DNA methylation and gene expression data to investigate epigenetic regulation of gene expression. The shared gene sets between DMGs and DEGs were clustered based on their regulatory status. Four shared clusters (Hypo-Up, Hyper-Up, Hypo-Down, and Hyper-Down) were identified based on the methylation and expression patterns.

Next, we conducted a weighted gene co-expression network analysis (WGCNA) to examine the correlation between the two omics datasets [56]. This approach enabled us to identify co-methylation and co-expression modules across tissues, which helped clarify the regulatory relationships. WGCNA was chosen because of its ability to reveal complex module relationships between large-scale datasets, making it well-suited for integrative multi-omics analysis. We then profiled the co-expression

and co-methylation modules from each omics dataset and analyzed the correlations between these modules using the Pearson method. Modules with high correlations were further investigated by identifying the functional roles of the gene sets within each correlated module. Finally, Cytoscape was used to construct gene interaction networks for highly correlated modules [57].

Results

DNA methylation and expression clustering

RNA-seq and MBD-seq data were generated from backfat, liver, and loin tissues at 10 and 26 weeks of age. The average number of raw reads was 36,983,389 for transcriptomic data and 11,351,613 for DNA methylation data. After quality trimming, the average reads were 36,706,627 and 11,242,492, respectively. The unique mapping rates averaged 75.50% for RNA-seq and 56.90% for MBD-seq, while the overall mapping rates were 92.88% and 93.82%, respectively (Table S1).

Clustering of methylation profiles (Fig. S1A) showed weak tissue and time points separation, whereas expression profiles (Fig. S1B) showed clear tissue-specific clustering, with liver forming a distinct cluster across stages. Different development stage related differences were also evident within liver and loin expression profiles, while methylation profiles were comparatively conserved. While DNA methylation profiles remained relatively conserved, the gene expression data suggested more dynamic changes.

Identification of differential methylation patterns during growth

A comparative analysis of DMGs across backfat, liver, and loin revealed tissue-specific and shared methylation patterns (Fig. 1A). Backfat and loin shared more DMGs than liver, which shared fewer with backfat and some with loin. Loin displayed 7,534 DMGs (6,772 unique), backfat 1,425 (695 unique), and liver 98 (58 unique) (Table S2, Fig. S2A). Hypermethylated genes predominated in backfat and loin, whereas liver DMG counts were limited, precluding inference of a predominant direction. The DMG counts are reported alongside the number unique to each tissue, and Venn partitions are provided in Fig. 1B with exact set sizes.

DMGs function annotations based on GO biological processes and KEGG pathway databases revealed tissue-specific and common biological functions (Fig. 1C, Fig. S3, Table S3). Backfat samples exhibited structural integrity (cell adhesion, DNA repair, and extracellular matrix), signaling pathways (GTPase, adhesion, integrin, and phosphorylation), and immune modulation (T-cell polarity and vesicle).

In the loin, hypermethylated genes were associated with the cell cycle, protein phosphorylation, and intracellular transport, which support the fundamental biological functions required to maintain muscle function and structure. Conversely, hypomethylated genes were associated with protein localization, actin filament dynamics, mitochondrial functions, and autophagy. Upon examining the comprehensive gene set, processes related to cytoskeletal organization, protein processing, and intracellular signaling pathways were found to be significantly enriched. Furthermore, MAPK signaling and insulin pathways were identified as playing a role in muscle development, energy balance, and cellular responses.

In contrast, methylation in liver tissue was more selective, with minimal widespread trends, and was primarily related to transcriptional regulation and hemostasis. The results of these analyses highlighted that backfat, loin, and liver tissues exhibit distinct methylation-associated biological functions and physiological roles. Backfat is mainly involved in immune regulation and cellular transport functions, while loin is more involved in muscle signaling and structure maintenance. In contrast, the functional contribution of DNA methylation in the liver is limited.

Profiling of gene expression dynamics during growth

Gene expression profiling identified DEGs in all three tissues (Fig. 2A). The liver displayed the highest numbers of unique DEGs at down- and up-regulation (4,131 and 3,535, respectively), with fewer in backfat (1,310 and 732) and loin (874 and 497) (Table S4, Fig. S2B). UpSet plots partitioned DEGs into up- and down-regulated sets, highlighting tissue-specific patterns: liver showed a larger proportion of up-regulated genes, while backfat and loin exhibited more down-regulated genes (Fig. 2B).

This analysis highlighted the distinct physiological roles of backfat, loin, and liver tissues, as evidenced by their unique functional enrichment of gene expression (Fig. 2C, Fig. S4, Table S5). Backfat was significantly enriched in lipid metabolism and energy regulation pathways, particularly fatty acid beta-oxidation and the positive regulation of cholesterol efflux. The upregulation of pathways such as retinol metabolism and hormone metabolic processes further emphasized the role of backfat in metabolic and endocrine functions, aligning with its primary function in fat storage and signaling.

The loin showed enrichment of pathways such as proteasome-mediated ubiquitin-dependent protein catabolic processes and intracellular signal transduction, which are broadly involved in muscle protein turnover and signaling regulation. In addition, pathways related to energy homeostasis, calcium signaling, and vascular smooth

muscle contraction were enriched in the loin transcriptome, reflecting general mechanisms of muscle contraction, calcium handling, and energy balance that support muscle functionality and growth.

Although the liver showed fewer enriched pathways than backfat and loin, it displayed notable upregulation in the regulation of RNA splicing and metabolic pathways, reinforcing its central role in systemic metabolism. Pathways such as calcium ion transmembrane transport and calcium ion import across the plasma membrane were also enriched, indicating that the liver contributes to calcium homeostasis. Taken together, these results illustrated both shared and unique physiological roles across tissues. Shared processes, such as metabolic pathways, reflect common functions, while specialized enrichment underscores tissue-specific roles, such as lipid metabolism in backfat, muscle function in the loin, and metabolic regulation in the liver.

Unified functional enrichment through common genes among DMGs and DEGs

Based on shared genes between DMGs and DEGs in each tissue, we aimed to analyze the direct regulatory interaction between DNA methylation and gene expression. The number of shared DMG–DEG genes identified were 71 (backfat), 230 (liver), and 18 (loin) (Fig. 3A–C, Fig. S5). Shared genes were grouped into four regulatory clusters by methylation–expression pairing: C1/C4 showed inverse relationships (hyper ↔ low, hypo ↔ high), whereas C2/C3 showed positive relationships (hyper ↔ high, hypo ↔ low) (Fig. 3D–F). Loin was enriched for C2 and C4 cluster genes, while backfat and liver were more evenly distributed across clusters. To maintain directionality clarity, cluster labels are aligned across panels (C1–C4), and colors are consistent between heatmaps and summary bar charts. Heatmaps (Fig. 3G) showed tissue-specific patterns; annotations distinguished promoter versus gene-body locations and regulatory roles (transcription factors/cofactors).

Functional enrichment analysis of shared genes offered a significant biological interpretation of the regulatory interactions (Table S6). Backfat was notably enriched in genes related to the structural integrity of the extracellular matrix (ECM), with significant terms associated with cell adhesion and integrin signaling. The liver was linked to the regulation of cytokinesis and mitotic spindle assembly, indicating its role in cell cycle control. The loin was enriched in cholesterol metabolism, protein glycosylation, and ECM organization, suggesting important metabolic functions.

Phenotype-module associations through WGCNA

Methylation modules (Fig. 4A) correlated primarily with stage, with green (liver $r = 0.73$; loin $r = -0.48$) and grey (backfat $r = -0.33$) showing additional tissue associations. The turquoise module lacked trait specificity and was excluded from downstream interpretation. Module–trait heatmaps display Pearson's r with P-values. Expression modules (Fig. 4B) showed distinct tissue/growth stage associations: blue/yellow positively with backfat and negatively with liver; magenta/turquoise positively with backfat and negatively with loin; black/red/green associated with liver–loin; greenyellow/grey/pink/purple associated with stage. These module characteristics indicated how gene methylation and expression patterns varied with the growth stage and across different tissues, revealing unique signatures in modules for tissue-specific roles and growth stage specific expression dynamics.

Integrating DNA methylation and gene expression data provided insights into the epigenetic interactions that may underlie tissue-specific and growth-stage-dependent regulation (Fig. 4C). Integration of module eigengenes revealed associations between expression (black/green/red/blue/pink/yellow) and methylation (green/grey) modules. In particular, the green methylation module showed a high correlation with liver and loin traits, and the expression levels of components of the black, red, and green modules indicated a similar tendency. In addition, the yellow and blue expression modules showed substantial associations with backfat and liver traits. Common traits among these modules suggested that epigenetic interactions in the liver play a complementary role. Furthermore, a negative correlation was identified between the grey module of backfat-related methylation and the blue and pink modules of expression. The overall module for DNA methylation and the green-yellow, grey, and purple modules for expression showed comprehensive correlations with the growth stage; therefore, the relationships between these modules were excluded from further analyses to focus on tissue specificity according to growth.

(See figure on next page.)

Fig. 1 Comparative analysis of differentially methylated genes (DMGs) and gene ontology enrichment across three tissues. **A** Venn diagram of DMGs. **B** Upset plot of shared DMGs based on hyper- and hypomethylation patterns. The bar graph represents the number of shared combinations between tissues, and the violin plot represents the $\log_2$ fold change (FC). **C** Gene Ontology biological process treemaps for DMGs, categorized by hypermethylated (Hyper), hypomethylated (Hypo), and total genes. Each box represents a biological process term, with the box size corresponding to the $-\log_{10}$(P-value) of the terms enrichment




Fig. 1 (See legend on previous page.)

Module based data integration across multi-omics layers

The functional gene enrichment network and associated regulatory mechanisms of each WGCNA module, integrating expression and DNA methylation data, present unique functional and core gene profiles (Fig. 5, Table S7). The yellow–green module played a minor role, with a limited number of genes associated with primary metabolic functions and stress responses (Fig. 5A). The blue–green integration module was significantly enriched in pathways related to extracellular matrix organization, collagen biosynthetic processes, and cell signaling. It included *COL1A1*, *COL5A1*, and ENSSSCG00000032761, which are essential for tissue structure and cell communication (Fig. 5B). The integration module (expression–methylation) indicated an association with backfat (positive–non-significant), liver (negative–positive), and loin (non-significant–negative), suggesting a targeted effect involved in connective tissue and extracellular matrix dynamics in pig growth. The black–green module functions related to oxygen transport and hydrogen peroxide catabolic processes were prominent (Fig. 5C). This module contained three to four genes with specific functions in the stress response and metabolic regulation. The primary functional process within the blue–grey integration module involved the ECM and collagen fibril organization, which are essential for tissue growth and repair (Fig. 5D). Genes such as *COL1A1*, *COL26A1*, and *TNXB* are involved in connective tissue function. Notably, the integration module showed associations with backfat (positive–negative) and the liver (negative–non-significant), suggesting a potential inhibitory mechanism in the liver tissue while promoting backfat development, with no significant relation to loin tissue. This pattern implied that adipose tissue development may involve demethylation processes that activate genes associated with tissue formation. Gene function analysis of the green–green integrated module indicated that it regulates various physiological processes, including muscle development and intercellular signaling (*MYH7*, *ACTN2*, *PLEC*, *MYOZ1*, *ALPK2*, and *PDLIM3*) and lipid and cholesterol metabolism (*ABCA4*, *ABCA5*, *ABCA8*, *ABCA9*, *VPS13B*, ENSSSCG00000015069; Fig. 5E). The negative correlation between gene expression and methylation levels in the liver (negative–positive) and loin (positive–negative) mediated tissue-specific functional regulation. The pink–grey module contributed to translation (ribosomal protein large/small subunit gene families), cell division (*CDC20*, *CCNB1*, and *CCNB3*), DNA replication (minichromosome maintenance gene families), and DNA repair mechanisms, indicating its high level of involvement in fundamental cellular processes (Fig. 5F). The red–green module focused on lipid metabolism, cellular proliferation, and viral responses (Fig. 5G). It included pathways such as cholesterol biosynthesis, homeostasis, metabolic processes, and the regulation of cell population proliferation, underscoring its role in maintaining the lipid balance, which is essential for energy storage and cellular function in the growth stage to the completion of development.

Discussion

Tissue-specific epigenetic regulation of fat metabolism

In this study, we analyzed DNA methylation and transcriptome data to clarify the regulatory mechanisms underlying fat metabolism in the liver, backfat, and loin muscle tissues during pig development. Our results indicated distinct tissue-specific methylation and gene expression patterns, highlighting the complex regulatory interactions essential for lipid metabolism and pork quality traits. These findings are consistent with those of previous studies, suggesting that epigenetic modifications play a significant role in metabolic regulation in pigs [26, 29, 32]. DNA methylation clustering analysis revealed relatively weak tissue specificity compared with gene expression clustering. This suggested that, while DNA methylation serves as a regulatory mechanism, transcriptional regulation may be more dominant in defining tissue-specific metabolic functions. Furthermore, DNA methylation was not evenly distributed across tissues. While backfat and loin muscle exhibited more extensive methylation changes, the liver demonstrated a limited range of DMGs, indicating significant epigenetic plasticity in tissues directly associated with fat storage and muscle growth, reinforcing its primary role as a metabolic hub that coordinates systemic lipid metabolism [12, 14].

Gene expression profiles and functional specialization

Although the liver had fewer DMRs, it had the highest number of DEGs. This suggested that hepatic gene regulation during development is less dependent on

(See figure on next page.)

Fig. 2 Comparative analysis of differentially expressed genes (DEGs) and enriched biological processes across three tissues. **A** Venn diagram of DEGs. **B** Upset plot of DEGs based on up- and downregulation patterns. The bar graph represents the number of shared combinations between tissues, and the violin plot represents the $\log_2$ fold change (FC). **C** Gene Ontology biological process treemaps for DEGs, highlighting upregulated (Up), downregulated (Down), and total gene sets. Each box represents a biological process term, with the box size corresponding to the $-\log_{10}$(*P*-value) of the terms enrichment




Fig. 2 (See legend on previous page.)




Fig. 3 Integrative analysis of shared genes between differential methylation and differential expression. Venn diagrams summarizing the intersection between differentially methylated genes (DMGs) and differentially expressed genes (DEGs) in (**A**) backfat, (**B**) loin, and (**C**) liver. **D**–**F** Regulation categorized clusters between methylation and expression levels. (**G**) Heatmap of common DEGs and DMGs in multiple tissues. Annotations include gene characteristics (promoter, gene body), regulatory roles (transcription factors, cofactors), and tissue

methylation dynamics and more influenced by trans-acting factors and signaling pathways. This finding is consistent with the observations of Wang et al. [25] who demonstrated that stable, yet functionally relevant methylation plays a role in regulating lipid metabolism in the liver [25]. Liver DEGs were primarily enriched in metabolic pathways, RNA splicing, and calcium homeostasis. This finding highlighted the importance of the liver in regulating overall lipid and energy metabolism [12, 14, 27]. In contrast, DEGs in backfat were significantly involved in lipid metabolism and endocrine signaling, which supports their recognized role in fat accumulation




Fig. 4 Correlation heatmap of DNA methylation and gene expression modules with phenotypic traits and inter-modular relationships. **A**–**B** Heatmaps showing the relationships between weighted gene co-expression network analysis modules, based on DNA methylation (**A**) and gene expression (**B**), and various phenotypic traits. **C** Heatmap of correlations reflecting the degree of association between methylation and expression modules. The correlations are represented by color intensity, with positive correlations in red and negative correlations in blue. The correlation coefficients and corresponding *P*-values are shown in each cell

and metabolic regulation [7, 8, 20]. The loin exhibited DEGs enriched in muscle functionality, protein turnover, and energy balance, emphasizing its essential function in pork tenderness and marbling [6, 58]. These results are consistent with those of previous studies, indicating that muscle and adipose tissues display tissue-specific transcriptomic signatures linked to their physiological functions [59]. Overall, the broader diversity of enriched methylation and expression signatures in backfat and loin suggests more extensive developmental and functional remodeling in these tissues compared with the liver.

Functional characteristics of genes with common differential methylation and expression

Integrative analysis of DNA methylation and transcriptomic data revealed genes that were both differentially methylated and expressed, highlighting metabolically regulated pathways under direct epigenetic control. In our dataset, approximately 70% of the annotated DMGs were located within gene body regions, whereas the remaining 30% were identified in promoter regions. This distribution suggests that, while promoter methylation typically serves a repressive function, gene methylation may have broader or alternative roles [60, 61]. Although promoter methylation is traditionally associated with transcriptional repression, gene body methylation has been linked to alternative splicing, transcriptional elongation, and even activation, depending on the genomic and chromatin contexts [62, 63]. Of note, many hypermethylated genes were also DEGs during the integrative analysis. This suggested that methylation and gene expression are not always inversely related, particularly in tissues with complex developmental programming, such as fat and muscle tissues. Previous studies in pigs and other livestock support this finding, indicating that gene body methylation can mark actively transcribed regions and correlate with tissue-specific gene functions [9, 64–66].

To clarify the functional impact of DNA methylation on gene regulation, we categorized the shared DMGs/DEGs based on tissue specificity (backfat, liver, and loin) and correlation patterns (positive, negative, or whole).




Fig. 5 Gene functional enrichment network between associated WGCNA modules (expression–methylation). **A** Yellow–green. **B** Blue–green. **C** Black–green. **D** Blue–grey. **E** Green–green. **F** Pink–grey. **G** Red–green. The node shapes indicate either genes (circles) or terms (diamonds). The node colors correspond to different gene types: DMG (red), DEG (yellow), and DMG/DEG (blue). Nodes with thick borders represent regulatory elements. Edge colors are based on associated modules

Functional enrichment analysis revealed distinct biological themes within each tissue, emphasizing the specificity of epigenetic regulation. In backfat tissue, genes such as *TNXB* (ENSSSCG00000023611), *ITGA8* (ENSSSCG00000011046), and *ITGA5* (ENSSSCG00000000293) are commonly associated with the negative regulation of ECM organization, cell–matrix adhesion, and cytoskeletal remodeling. *TNXB* encodes tenascin-X, a glycoprotein involved in matrix stabilization that has been shown to influence ECM remodeling during adipose tissue expansion [67]. Similarly, ITGA, an integrin subunit, mediates fibroblast-ECM interactions and has been implicated in tissue stiffness, thereby affecting preadipocyte differentiation [68]. In contrast, positively regulated genes in backfat, such as *PLEC* (ENSSSCG00000039094) and *FLNC* (ENSSSCG00000016578), promote sarcomeric cytoskeleton integrity and mechanical sensing in differentiating adipocytes, possibly reflecting functional convergence between muscle-like cytoskeletal pathways and adipocyte morphology [69, 70]. In loin muscle, negatively regulated genes, such as *MSMO1* (ENSSSCG00000008857) and *LMNB2* (ENSSSCG00000026784) are frequently involved in cholesterol biosynthesis, nuclear architecture, and collagen biosynthesis, respectively. MSMO1 plays a crucial role in sterol biosynthesis and is closely linked to lipid accumulation and cellular membrane integrity in intramuscular adipocytes [71]. In contrast, positively regulated genes in the loin were enriched for muscle fiber structure (*LMNB2*—ENSSSCG00000026784) and collagen biosynthesis (*COL1A1*—ENSSSCG00000036135), facilitating active muscle development and integrity during later growth stages. COL1A1, a type I collagen complex, is essential for ECM stiffness and structural support during muscle development, and is often co-expressed with COL1A2 in skeletal-muscle-related traits [72–74]. These patterns highlight that methylation-expression correlations differ by function and tissue type. In both backfat and loin, several genes exhibited changes in expression despite hypermethylation, emphasizing the complexity of regulation, particularly in the context of gene body methylation. Importantly, key pathways such as the PI3K-Akt signaling and cholesterol biosynthesis pathways emerged in both positive and negative contexts, reflecting stage-dependent epigenetic reprogramming rather than static repression.

Integration of epigenetic regulation network and tissue specificity through module correlation

The module-based integration of DNA methylation and gene expression provides deeper insights into how large-scale gene networks are regulated in a tissue- and growth-specific manner during pig growth. Unlike individual gene-level comparisons, the WGCNA-based approach allowed us to uncover how co-methylated and co-expressed gene clusters act as coordinated functional units, which may better reflect the underlying biological architecture of complex traits, such as fat deposition and muscle development [56]. The green methylation module, which showed a strong correlation with the liver (positive), backfat (negative), and loin (negative) phenotypes, emerged as a key epigenetic driver module. Its significant association with expression modules such as the black, red, and green modules, which were enriched for cholesterol biosynthesis, cell proliferation, and oxidative metabolism, suggests that DNA methylation may act as an upstream regulatory mechanism orchestrating liver-specific metabolic functions [25, 32, 62]. Of note, the directionality of the correlation between methylation and expression was not uniform across tissues, and both negative and positive relationships were observed. This variability supports previous findings that DNA methylation, especially in gene body regions, does not always suppress gene expression, and may instead reflect transcriptional activity or chromatin accessibility [60, 64].

Furthermore, the identification of tissue-specific module correlations, such as the blue and yellow expression modules correlating with backfat traits or the green-green module showing opposing regulation between the liver and loin, raises important questions about the mechanisms by which epigenetic states are established and maintained in a context-dependent manner. From a functional perspective, the identification of collagen-related genes (*COL1A1*, *COL26A1*, and *TNXB*) and cholesterol metabolism regulators (*MSMO1* and *HMGCR*) within highly correlated modules underscores the dual role of epigenetic regulation: one mediates structural remodeling in fat and muscle and the other coordinates metabolic homeostasis. Our findings suggest that their activity may be shaped by the methylation-driven reprogramming of module-wide transcription, which may serve as a fine-tuning mechanism for aligning tissue-specific functions with systemic metabolic demands. This perspective aligns with recent models that propose that epigenetic states modulate transcriptional response ranges rather than absolute on/off switches. Several of these genes also appeared in the DEG-DMGs that overlapped with the first integration, reinforcing their central role. The convergence of these findings from distinct analytical strategies lends robustness to their biological relevance and indicates their potential as biomarkers or targets in breeding programs focused on meat quality traits. Moreover, the tissue-specific nature of correlated modules suggests that epigenetic mechanisms are not uniformly deployed, but are tailored to support each tissue's unique functional requirements. For example, the contrasting patterns observed in the liver and backfat modules imply a context-dependent role for DNA

methylation in modulating metabolic gene networks. These findings echo prior observations in both livestock and human studies, where environmental cues, developmental timing, and tissue physiology jointly shaped epigenetic landscapes. Finally, it is worth noting that although WGCNA provides a powerful framework for identifying correlated gene clusters, it cannot confirm causal relationships. Although biologically suggestive, the observed correlations require further validation through experimental perturbation studies, such as CRISPR-based epigenetic editing or methylation inhibition. However, due to the limited amount of biological material and the comprehensive multi-tissue, multi-omics nature of this study, additional functional assays (e.g., protein-level or gene editing experiments) could not be performed. We acknowledge this as a limitation of the present work. Nonetheless, our findings represent a critical first step in establishing modular epigenetic networks as key regulators of tissue development and metabolic specialization in pigs. These results provide a valuable resource and experimental foundation for future targeted functional validation. Furthermore, epigenetic regulation operates through a multilayered machinery that involves not only DNA methylation but also histone modifications and chromatin architecture, and this strategy appears to be conserved across diverse biological systems. For example, in bacteria, DNA methylation and nucleoid-associated proteins cooperatively regulate gene expression and virulence, while in other livestock species such as cattle and sheep, the interplay between histone marks and chromatin accessibility has been shown to fine-tune developmental gene programs [75, 76]. Together, these examples, ranging from prokaryotes to mammals, highlight that DNA methylation is functionally intertwined with higher-order chromatin states. Therefore, future studies in pigs should integrate histone profiling and chromatin conformation analysis (e.g., Hi-C) with DNA methylation and transcriptomic data to capture a more comprehensive landscape of epigenetic regulation governing fat metabolism.

Potential environmental influences and comparative perspectives

Although all pigs were managed under uniform housing, feeding, and husbandry conditions, residual environmental factors (e.g., microenvironment, health status, early life influences) may still have contributed to the observed epigenetic and transcriptomic variation. Previous studies in livestock have shown that nutrition, thermal stress, pathogens, and other management-related factors can induce stable changes in DNA methylation and gene expression, thereby modulating growth, metabolism, and product quality [77–80]. These potential environmental influences should be considered when extrapolating our findings and motivate future experiments designed to explicitly manipulate and measure such factors.

Tissue and development stage specific epigenetic patterns similar to those observed here in porcine backfat, loin, and liver have been reported in other livestock species. In cattle and sheep, DNA methylation and transcriptome studies in subcutaneous and intramuscular adipose depots have linked depot-specific methylation signatures in lipid metabolic genes to differences in fat deposition, fatty acid composition, and carcass quality, paralleling the enrichment of lipid metabolism pathways detected in pig backfat [26, 29]. Likewise, comparative analyses of skeletal muscle methylomes in cattle and poultry have shown that methylation changes in myogenic, cytoskeletal, and energy metabolism genes accompany growth, fibre-type composition, and age-related transitions, which is consistent with the muscle-regulatory and energy homeostasis pathways enriched in the porcine loin across developmental stages [81–83]. The studies in livestock have shown that epigenetic regulation of hepatic genes is associated with systemic energy balance, feed efficiency, and responses to nutritional or environmental challenges. For example, DNA methylation changes in bovine have been linked to divergent residual feed intake and to differences in nutrient metabolism and metabolic stress across diets or thermal environments [84, 85]. Similar work in humans further indicates that epigenetic modifications in the liver contribute to the regulation of hepatic lipid metabolism and broader metabolic homeostasis [86]. Moreover, recent cross-species comparative analyses of DNA methylation in brain, liver, and skeletal muscle across multiple mammals, including cattle, sheep, goats, pigs, and mice, support the idea that modular epigenetic regulation of key metabolic tissues is a conserved feature underlying complex traits such as growth, feed efficiency, and adaptation in domesticated animals [87].

Conclusion

This study provides a comprehensive multi-omics overview of tissue-specific epigenetic regulation of fat metabolism in pigs. Through the integration of DNA methylation and gene expression data from the liver, backfat, and loin, we revealed that methylation exerts a fine-tuning role rather than a dominant regulatory effect. While the liver exhibited fewer methylation changes but extensive transcriptional variation, backfat and loin showed greater epigenetic plasticity linked to adipogenesis and muscle structure. Integrative network analysis identified tissue-dependent co-regulated modules, emphasizing extracellular matrix remodeling, cholesterol metabolism, and muscle fiber development

as key pathways influenced by methylation–expression interactions. While these findings advance our mechanistic understanding, several limitations temper immediate translational applications: (i) causality requires experimental validation; (ii) protein-level confirmation is needed; and (iii) generalizability across breeds remains to be established. The identified epigenetic markers provide a foundation for future hypothesis-driven studies and may eventually inform breeding strategies once validated in larger populations under diverse conditions.

Abbreviations

COL1A1	Collagen type I alpha 1 chain
COL26A1	Collagen type XXVI alpha 1 chain
DAVID	Database annotation visualization integrated discovery
DEG	Differentially expressed gene
DMG	Differentially methylated gene
DMR	Differentially methylated regions
ECM	Extracellular matrix
FC	Fold change
FLNC	Filamin C
GO	Gene Ontology
HMGCR	3-Hydroxy-3-methylglutaryl-CoA reductase
IMF	Intramuscular fat
ITGA5	Integrin subunit alpha 5
ITGA8	Integrin subunit alpha 8
KEGG	Kyoto Encyclopedia of Genes and Genomes
KNP	Korean native pig
LMNB2	Lamin B2
MAPK	Mitogen-activated protein kinase
MBD-seq	Methylated DNA-binding domain sequencing
MSMO1	Methylsterol monooxygenase 1
PLEC	Plectin
TMM	Trimmed mean of M values
TNXB	Tenascin XB
WGCNA	Weighted gene co-expression network analysis

Supplementary Information

The online version contains supplementary material available at https://doi.org/10.1186/s40104-025-01347-x.

Additional file 1: Table S1. Summary of RNA-seq and MBD-seq Quality Control and Mapping Statistics. Table S2. List of differentially methylated genes identified in the study. Table S3. Functional analysis results of differentially methylated genes based on enriched GO terms and KEGG pathways. Table S4. List of differentially expressed genes identified in the study. Table S5. Functional analysis results of differentially expressed genes based on enriched GO terms and KEGG pathways. Table S6. Functional enrichment of shared genes between differentially methylated and expressed genes based on gene element annotations. Table S7. Functional analysis of gene sets in the module integrating expression and DNA methylation data.

Additional file 2: Fig. S1. Multidimensional scaling plots of DNA methylation and gene expression data across tissues and developmental stages. (A) DNA methylation data. (B) Gene expression data. Fig. S2. Volcano plots of transcriptomic and epigenetic changes across porcine tissues. (A) DNA methylation data. (B) Gene expression. Yellow, brown, and purple dots represent genes from backfat, liver, and loin tissues, respectively. Each plot displays the $\log_2$ fold change on the *x*-axis and the $-\log_{10}$ *P*-value on the *y*-axis. Fig. S3. KEGG pathway enrichment analysis for differentially methylated genes. Fig. S4. KEGG pathway enrichment analysis for differentially expressed genes. Fig. S5. Integrative analysis of shared genes by genomic features between differential methylation and differential expression. (A, D) Back fat. (B, E) Loin. (C, F) Liver.

Acknowledgements
Not applicable.

Authors' contributions
DYK: Writing – original draft, Writing – review & editing, Visualisation, Software, Methodology, Formal analysis, Data curation. BHL: Writing – review & editing, Methodology. RKP: Writing – review & editing. WCP: Writing – review & editing, Data curation. JEP: Writing – review & editing, Data curation, Conceptualisation, Funding acquisition. JMK: Writing – review & editing, Conceptualisation, Project administration, Supervision, Funding acquisition.

Funding
This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education (Grant No. RS-2023-00245099), the Jeju RISE Center, funded by the Ministry of Education and the Jeju Special Self-Governing Province in 2025, as part of the "Regional Innovation System & Education (RISE): Glocal University 30" initiative, and the Cooperative Research Program for Agriculture Science & Technology Development, Rural Development Administration, Republic of Korea (Grant No. RS-2021-RD010153).

Data availability
The raw read data used in this study have been deposited in the NCBI Sequence Read Archive (SRA) under accession No. PRJNA768488 and PRJNA1251969.

Declarations

Ethics approval and consent to participate
All experimental protocols were approved by the Institutional Animal Care and Use Committee of the National Institute of Animal Science, South Korea (approval No. NIAS2016-848).

Consent for publication
Not applicable.

Competing interests
The authors declare no competing interests.

Author details
[1]Department of Animal Science and Technology, Chung-Ang University, Anseong, Gyeonggi-Do 17546, Republic of Korea. [2]Department of Animal Science, Iowa State University, Ames, IA 50011, USA. [3]Pere Virgili Institute for Health Research, Tarragona 43005, Spain. [4]Animal Genetics and Breeding Division, National Institute of Animal Science, RDA, 114, Sinbang 1-Gil, Seonghwan-Eup, Seobuk-Gu, Cheonan-Si, Chungcheongnam-Do 31000, Republic of Korea. [5]Department of Animal Biotechnology, College of Applied Life Science, Jeju National University, Jeju-Si 63243, Republic of Korea.

Received: 14 October 2025 Accepted: 25 December 2025
Published online: 11 March 2026

References

1. Lebret B, Lhuisset S, Labussiere E, Louveau I. Combining pig genetic and feeding strategies improves the sensory, nutritional and technological quality of pork in the context of relocation of feed resources. Meat Sci. 2023;197:109074. https://doi.org/10.1016/j.meatsci.2022.109074.
2. Khanal P, Maltecca C, Schwab C, Gray K, Tiezzi F. Genetic parameters of meat quality, carcass composition, and growth traits in commercial swine. J Anim Sci. 2019;97(9):3669–83. https://doi.org/10.1093/jas/skz247.
3. Dekkers JC. Marker-assisted selection for commercial crossbred performance. J Anim Sci. 2007;85(9):2104–14. https://doi.org/10.2527/jas.2006-683.
4. Malgwi IH, Halas V, Grunvald P, Schiavon S, Jocsak I. Genes related to fat metabolism in pigs and intramuscular fat content of pork: A focus on nutrigenetics and nutrigenomics. Animals (Basel). 2022;12(2):150. https://doi.org/10.3390/ani12020150.

5. Pandey S, Kim S, Kim ES, Keum GB, Doo H, Kwak J, et al. Exploring the multifaceted factors affecting pork meat quality. J Anim Sci Technol. 2024;66(5):863–75. https://doi.org/10.5187/jast.2024.e56.
6. Wood JD, Enser M, Fisher AV, Nute GR, Sheard PR, Richardson RI, et al. Fat deposition, fatty acid composition and meat quality: a review. Meat Sci. 2008;78(4):343–58. https://doi.org/10.1016/j.meatsci.2007.07.019.
7. Hausman GJ, Dodson MV, Ajuwon K, Azain M, Barnes KM, Guan LL, et al. Board-invited review: The biology and regulation of preadipocytes and adipocytes in meat animals. J Anim Sci. 2009;87(4):1218–46. https://doi.org/10.2527/jas.2008-1427.
8. Poklukar K, Čandek-Potokar M, Batorek Lukač N, Tomažin U, Škrlep M. Lipid deposition and metabolism in local and modern pig breeds: A review. Animals (Basel). 2020. https://doi.org/10.3390/ani10030424.
9. Liu J, Sebastià C, Jové-Juncà T, Quintanilla R, González-Rodríguez O, Passols M, et al. Identification of genomic regions associated with fatty acid metabolism across blood, liver, backfat and muscle in pigs. Genet Sel Evol. 2024;56:66. https://doi.org/10.1186/s12711-024-00933-3.
10. Xing K, Zhu F, Zhai L, Chen S, Tan Z, Sun Y, et al. Identification of genes for controlling swine adipose deposition by integrating transcriptome, whole-genome resequencing, and quantitative trait loci data. Sci Rep. 2016;6:23219. https://doi.org/10.1038/srep23219.
11. Zhao S, Wang J, Song X, Zhang X, Ge C, Gao S. Impact of dietary protein on lipid metabolism-related gene expression in porcine adipose tissue. Nutr Metab (Lond). 2010;7:6. https://doi.org/10.1186/1743-7075-7-6.
12. Chakravarthy MV, Pan Z, Zhu Y, Tordjman K, Schneider JG, Coleman T, et al. "New" hepatic fat activates pparalpha to maintain glucose, lipid, and cholesterol homeostasis. Cell Metab. 2005;1(5):309–22. https://doi.org/10.1016/j.cmet.2005.04.002.
13. Kristensen NB, Wu G. Metabolic functions of the porcine liver. In: Bach Knudsen IKE, Kjeldsen NJ, Poulsen HD, Jensen BB, editors. Nutritional physiology of pigs-online publication. Videncenter for Svineproduktion; 2012.
14. Frayn K, Arner P, Yki-Järvinen H. Fatty acid metabolism in adipose tissue, muscle and liver in health and disease. Essays Biochem. 2006;42(1):89–103. https://doi.org/10.1042/bse0420089.
15. Han Q, Huang X, He J, Zeng Y, Yin J, Yin Y. Intramuscular fat deposition in pig: a key target for improving pork quality. J Integr Agric. 2025;24(12):4461–83. https://doi.org/10.1016/j.jia.2024.03.005.
16. Yan E, Guo J, Yin J. Nutritional regulation of skeletal muscle energy metabolism, lipid accumulation and meat quality in pigs. Anim Nutr. 2023;14:185–92. https://doi.org/10.1016/j.aninu.2023.04.009.
17. Alfonso L, Mourot J, Insausti K, Mendizabal JA, Arana A. Comparative description of growth, fat deposition, carcass and meat quality characteristics of Basque and Large White pigs. Anim Res. 2005;54(1):33–42. https://doi.org/10.1051/animres:2005001.
18. Li M, Zhu L, Li X, Shuai S, Teng X, Xiao H, et al. Expression profiling analysis for genes related to meat quality and carcass traits during postnatal development of backfat in two pig breeds. Sci China C Life Sci. 2008;51(8):718–33. https://doi.org/10.1007/s11427-008-0090-0.
19. Duran-Montgé P, Theil PK, Lauridsen C, Esteve-Garcia E. Fat metabolism is regulated by altered gene expression of lipogenic enzymes and regulatory factors in liver and adipose tissue but not in semimembranosus muscle of pigs during the fattening period. Animal. 2009;3(11):1580–90. https://doi.org/10.1017/S1751731109990450.
20. Gong X, Zheng M, Zhang J, Ye Y, Duan M, Chamba Y, et al. Transcriptomics-based study of differentially expressed genes related to fat deposition in Tibetan and Yorkshire pigs. Front Vet Sci. 2022;9:919904. https://doi.org/10.3389/fvets.2022.919904.
21. Jones PA, Takai D. The role of DNA methylation in mammalian epigenetics. Science. 2001;293(5532):1068–70. https://doi.org/10.1126/science.1063852.
22. Reik W, Dean W, Walter J. Epigenetic reprogramming in mammalian development. Science. 2001;293(5532):1089–93. https://doi.org/10.1126/science.1063443.
23. Triantaphyllopoulos KA, Ikonomopoulos I, Bannister AJ. Epigenetics and inheritance of phenotype variation in livestock. Epigenetics Chromatin. 2016;9:1–18. https://doi.org/10.1186/s13072-016-0081-5.
24. Wang M, Feng S, Ma G, Miao Y, Zuo B, Ruan J, et al. Whole-genome methylation analysis reveals epigenetic variation in cloned and donor pigs. Front Genet. 2020;11:23. https://doi.org/10.3389/fgene.2020.00023.
25. Wang S, Zha L, Cui X, Yeh YT, Liu R, Jing J, et al. Epigenetic regulation of hepatic lipid metabolism by DNA methylation. Adv Sci (Weinh). 2023;10(20):e2206068. https://doi.org/10.1002/advs.202206068.
26. Li M, Wu H, Luo Z, Xia Y, Guan J, Wang T, et al. An atlas of DNA methylomes in porcine adipose and muscle tissues. Nat Commun. 2012;3:850. https://doi.org/10.1038/ncomms1854.
27. Gomez-Alonso MDC, Kretschmer A, Wilson R, Pfeiffer L, Karhunen V, Seppälä I, et al. DNA methylation and lipid metabolism: an ewas of 226 metabolic measures. Clin Epigenetics. 2021;13:7. https://doi.org/10.1186/s13148-020-00957-8.
28. Arora D, Park J-E, Lim D, Choi B-H, Cho I-C, Srikanth K, et al. Comparative methylation and rna-seq expression analysis in cpg context to identify genes involved in backfat vs. liver diversification in nanchukmacdon pig. BMC Genomics. 2021;22:801. https://doi.org/10.1186/s12864-021-08123-x.
29. Zhang S, Shen L, Xia Y, Yang Q, Li X, Tang G, et al. DNA methylation landscape of fat deposits and fatty acid composition in obese and lean pigs. Sci Rep. 2016;6:35063. https://doi.org/10.1038/srep35063.
30. Pan Z, Yao Y, Yin H, Cai Z, Wang Y, Bai L, et al. Pig genome functional annotation enhances the biological interpretation of complex traits and human disease. Nat Commun. 2021;12:5848. https://doi.org/10.1038/s41467-021-26153-7.
31. Foissac S, Djebali S, Munyard K, Vialaneix N, Rau A, Muret K, et al. Multi-species annotation of transcriptome and chromatin structure in domesticated animals. BMC Biol. 2019;17:108. https://doi.org/10.1186/s12915-019-0726-5.
32. Jacobsen MJ, Havgaard JH, Anthon C, Mentzel CMJ, Cirera S, Krogh PM, et al. Epigenetic and transcriptomic characterization of pure adipocyte fractions from obese pigs identifies candidate pathways controlling metabolism. Front Genet. 2019;10:1268. https://doi.org/10.3389/fgene.2019.01268.
33. Ci L, Sun H, Huang Y, Guo J, Albrecht E, Zhao R, et al. Maternal dietary fat affects the lt muscle fatty acid composition of progeny at weaning and finishing stages in pigs. Meat Sci. 2014;96(3):1141–6. https://doi.org/10.1016/j.meatsci.2013.10.033.
34. Collins CL, Pluske JR, Morrison RS, McDonald TN, Smits RJ, Henman DJ, et al. Post-weaning and whole-of-life performance of pigs is determined by live weight at weaning and the complexity of the diet fed after weaning. Anim Nutr. 2017;3(4):372–9. https://doi.org/10.1016/j.aninu.2017.01.001.
35. Hoile SP, Lillycrop KA, Grenfell LR, Hanson MA, Burdge GC. Increasing the folic acid content of maternal or post-weaning diets induces differential changes in phosphoenolpyruvate carboxykinase mrna expression and promoter methylation in rats. Br J Nutr. 2012;108(5):852–7. https://doi.org/10.1017/S0007114511006155.
36. Raj S, Skiba G, Weremko D, Fandrejewski H, Migdal W, Borowiec F, et al. The relationship between the chemical composition of the carcass and the fatty acid composition of intramuscular fat and backfat of several pig breeds slaughtered at different weights. Meat Sci. 2010;86(2):324–30. https://doi.org/10.1016/j.meatsci.2010.04.037.
37. Wu F, Vierck K, DeRouchey J, O'Quinn T, Tokach M, Goodband R, et al. A review of heavy weight market pigs: Status of knowledge and future needs assessment. Transl Anim Sci. 2017;1(1):1. https://doi.org/10.2527/tas2016.0004.
38. Ayuso D, González A, Peña F, Hernández-García FI, Izquierdo M. Effect of fattening period length on intramuscular and subcutaneous fatty acid profiles in Iberian pigs finished in the montanera sustainable system. Sustainability. 2020;12(19):7937. https://doi.org/10.3390/su12197937.
39. Kim YS, Kim SW, Weaver MA, Lee CY. Increasing the pig market weight: world trends, expected consequences and practical considerations. Asian-Australas J Anim Sci. 2005;18:590–600. https://doi.org/10.5713/ajas.2005.590.
40. Su-Young S, Kyoung-Mi P, Hoa V-B, Hyun-Woo S, Soo-Hyun C, Beom-Young P, et al. Pork quality characteristics of different parts from crossbred (Jeju Native Pig × Landrace) pigs. J Agric Life Sci. 2015;49(5):211–23. https://doi.org/10.14397/jals.2015.49.5.211.
41. Kumar H, Park W, Srikanth K, Choi BH, Cho ES, Lee KT, et al. Comparison of bacterial populations in the ceca of swine at two different stages and their functional annotations. Genes (Basel). 2019. https://doi.org/10.3390/genes10050382.

42. Kumar H, Srikanth K, Park W, Lee K-T, Choi B-H, Kim J-M, et al. Stage specific transcriptome analysis of liver tissue from a crossbred Korean native pig (KNP × Yorkshire). J Biomed Transl Res. 2018;19(4):116–24. https://doi.org/10.12729/jbtr.2018.19.4.116.
43. Seo YJ, Lim B, Kim DY, Lim KS, Kim JM. Regulation of swine growth by backfat tissue during growing and finishing stages. Animals (Basel). 2021. https://doi.org/10.3390/ani11123511.
44. Andrews S. Fastqc: A quality control tool for high throughput sequence data. Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc. Babraham Bioinformatics. 2010.
45. Krueger F, James F, Ewels P, Afyounian E, Weinstein M, Schuster-Boeckler B, et al. Trim galore. Babraham Bioinformatics. 2023. https://doi.org/10.5281/zenodo.7598955.
46. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for illumina sequence data. Bioinformatics. 2014;30(15):2114–20. https://doi.org/10.1093/bioinformatics/btu170.
47. Langmead B, Salzberg SL. Fast gapped-read alignment with bowtie 2. Nat Methods. 2012;9(4):357–9. https://doi.org/10.1038/nmeth.1923.
48. Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. Nat Biotechnol. 2019;37(8):907–15. https://doi.org/10.1038/s41587-019-0201-4.
49. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The sequence alignment/map format and samtools. Bioinformatics. 2009;25(16):2078–9. https://doi.org/10.1093/bioinformatics/btp352.
50. Liao Y, Smyth GK, Shi W. Featurecounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics. 2013;30(7):923–30. https://doi.org/10.1093/bioinformatics/btt656.
51. Lienhard M, Grimm C, Morkel M, Herwig R, Chavez L. Medips: genome-wide differential coverage analysis of sequencing data derived from DNA enrichment experiments. Bioinformatics. 2014;30(2):284–6. https://doi.org/10.1093/bioinformatics/btt650.
52. Robinson MD, McCarthy DJ, Smyth GK. Edger: a bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. 2010;26(1):139–40. https://doi.org/10.1093/bioinformatics/btp616.
53. Sherman BT, Hao M, Qiu J, Jiao X, Baseler MW, Lane HC, et al. David: A web server for functional enrichment analysis and functional annotation of gene lists (2021 update). Nucleic Acids Res. 2022;50(W1):W216–21. https://doi.org/10.1093/nar/gkac194.
54. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology: tool for the unification of biology. Nat Genet. 2000;25(1):25–9. https://doi.org/10.1038/75556.
55. Kanehisa M, Goto S. KEGG: Kyoto encyclopedia of genes and genomes. Nucleic Acids Res. 2000;28(1):27–30. https://doi.org/10.1093/nar/28.1.27.
56. Langfelder P, Horvath S. WGCNA: An R package for weighted correlation network analysis. BMC Bioinformatics. 2008;9:559. https://doi.org/10.1186/1471-2105-9-559.
57. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: a software environment for integrated models of biomolecular interaction networks. Genome Res. 2003;13(11):2498–504. https://doi.org/10.1101/gr.1239303.
58. Suzuki K, Ishida M, Kadowaki H, Shibata T, Uchida H, Nishida A. Genetic correlations among fatty acid compositions in different sites of fat tissues, meat production, and meat quality traits in duroc pigs. J Anim Sci. 2006;84(8):2026–34. https://doi.org/10.2527/jas.2005-660.
59. Zhang J, Meng S, Wang H, Zhang C, Sun Z, Huang L, et al. Comparison of growth performance, carcass properties, fatty acid profile, and genes involved in fat metabolism in nanyang and landrace pigs. Genes (Basel). 2024. https://doi.org/10.3390/genes15020186.
60. Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, et al. Conserved role of intragenic DNA methylation in regulating alternative promoters. Nature. 2010;466(7303):253–7. https://doi.org/10.1038/nature09165.
61. Flintoft L. DNA methylation: Looking beyond promoters. Nat Rev Genet. 2010;11(9):596. https://doi.org/10.1038/nrg2851.
62. Zhang Z, Chang L, Wang B, Wei Y, Li X, Li X, et al. Differential chromatin accessibility and gene expression associated with backfat deposition in pigs. BMC Genomics. 2024;25:902. https://doi.org/10.1186/s12864-024-10805-1.
63. Wang Y, Jin H, Tong X, Yu H, Li X, Zeng B. DNA methylation of postnatal liver development in pigs. Genes (Basel). 2024. https://doi.org/10.3390/genes15081067.
64. Kim DY, Lim B, Lim D, Park W, Lee KT, Cho ES, et al. Integrative methylome and transcriptome analysis of porcine abdominal fat indicates changes in fat metabolism and immune responses during different development. J Anim Sci. 2022. https://doi.org/10.1093/jas/skac302.
65. Zhou Y, Connor EE, Bickhart DM, Li C, Baldwin RL, Schroeder SG, et al. Comparative whole genome DNA methylation profiling of cattle sperm and somatic tissues reveals striking hypomethylated patterns in sperm. Gigascience. 2018. https://doi.org/10.1093/gigascience/giy039.
66. Luo R, Dai X, Zhang L, Li G, Zheng Z. Genome-wide DNA methylation patterns of muscle and tail-fat in dairymeade sheep and mongolian sheep. Animals (Basel). 2022. https://doi.org/10.3390/ani12111399.
67. Matsumoto K, Sato T, Oka S, Inokuchi J, Ariga H. Comparison of the compositions of phospholipid-associated fatty acids in wild-type and extracellular matrix tenascin-x-deficient mice. Biol Pharm Bull. 2004;27(9):1447–50. https://doi.org/10.1248/bpb.27.1447.
68. Roth T, Podesta F, Stepp MA, Boeri D, Lorenzi M. Integrin overexpression induced by high glucose and by human diabetes: potential pathway to cell dysfunction in diabetic microangiopathy. Proc Natl Acad Sci USA. 1993;90(20):9640–4. https://doi.org/10.1073/pnas.90.20.9640.
69. Castanon MJ, Wiche G. Identifying plectin isoform functions through animal models. Cells. 2021. https://doi.org/10.3390/cells10092453.
70. Deng Y, Yan J. Force-dependent structural changes of filamin c rod domains regulated by filamin c dimer. J Am Chem Soc. 2023;145(27):14670–8. https://doi.org/10.1021/jacs.3c02303.
71. Xin Y, Li C, Guo Y, Xiao R, Zhang H, Zhou G. Rna-seq analysis reveals a negative role of msmo1 with a synergized nsdhl expression during adipogenesis of 3t3-l1. Biosci Biotechnol Biochem. 2019;83(4):641–52. https://doi.org/10.1080/09168451.2018.1559719.
72. McKee TJ, Perlman G, Morris M, Komarova SV. Extracellular matrix composition of connective tissues: a systematic review and meta-analysis. Sci Rep. 2019;9:10542. https://doi.org/10.1038/s41598-019-46896-0.
73. Miyauchi Y, Kiyama M, Soga S, Nanri H, Ogiwara T, Yonamine S, et al. Downregulation of genes for skeletal muscle extracellular matrix components by cisplatin. Biol Pharm Bull. 2024;47(11):1846–50. https://doi.org/10.1248/bpb.b24-00428.
74. Wu Y, Xu Y. Integrated bioinformatics analysis of expression and gene regulation network of col12a1 in colorectal cancer. Cancer Med. 2020;9(13):4743–55. https://doi.org/10.1002/cam4.2899.
75. Kim KD, Tanizawa H, De Leo A, Vladimirova O, Kossenkov A, Lu F, et al. Epigenetic specifications of host chromosome docking sites for latent Epstein-Barr virus. Nat Commun. 2020;11:877. https://doi.org/10.1038/s41467-019-14152-8.
76. Ponsuksili S, Li S, Siengdee P, Hadlich F, Trakooljul N, Oster M, et al. DNA methylation in adipocyte differentiation of porcine mesenchymal stem cells and the impact of the donor metabolic type. Genomics. 2025;117(3):111050. https://doi.org/10.1016/j.ygeno.2025.111050.
77. Aravindh S, Silpa MV, Voggu SP, Rebez EB, Kalaignazhal G, Srinivas MV, et al. Epigenetic mechanisms associated with livestock adaptation to heat stress. Biology (Basel). 2025. https://doi.org/10.3390/biology14091154.
78. Dunislawska A, Pietrzak E, Wishna Kadawarage R, Beldowska A, Siwek M. Pre-hatching and post-hatching environmental factors related to epigenetic mechanisms in poultry. J Anim Sci. 2022. https://doi.org/10.1093/jas/skab370.
79. Ibeagha-Awemu EM, Yu Y. Consequence of epigenetic processes on animal health and productivity: is additional level of regulation of relevance? Anim Front. 2021;11(6):7–18. https://doi.org/10.1093/af/vfab057.
80. Ibeagha-Awemu EM, Zhao X. Epigenetic marks: regulators of livestock phenotypes and conceivable sources of missing variation in livestock improvement programs. Front Genet. 2015;6:302. https://doi.org/10.3389/fgene.2015.00302.
81. Chen S, Liu S, Shi S, Jiang Y, Cao M, Tang Y, et al. Comparative epigenomics reveals the impact of ruminant-specific regulatory elements on complex traits. BMC Biol. 2022;20:273. https://doi.org/10.1186/s12915-022-01459-0.
82. Kuraz Abebe B, Wang J, Guo J, Wang H, Li A, Zan L. A review of the role of epigenetic studies for intramuscular fat deposition in beef cattle. Gene. 2024;908:148295. https://doi.org/10.1016/j.gene.2024.148295.

83. Ran J, Li J, Yin L, Zhang D, Yu C, Du H, et al. Comparative analysis of skeletal muscle DNA methylation and transcriptome of the chicken embryo at different developmental stages. Front Physiol. 2021;12:697121. https://doi.org/10.3389/fphys.2021.697121.
84. Devos J, Behrouzi A, Paradis F, Straathof C, Li C, Colazo M, et al. Genetic potential for residual feed intake and diet fed during early- to mid-gestation influences post-natal DNA methylation of imprinted genes in muscle and liver tissues in beef cattle. J Anim Sci. 2021. https://doi.org/10.1093/jas/skab140.
85. Laporta J, Dado-Senn B, Guadagnin AR, Liu L, Penagaricano F. Preweaning heat stress alters liver transcriptome and DNA methylation in dairy calves. J Dairy Sci. 2025;108(4):4390–402. https://doi.org/10.3168/jds.2024-25975.
86. Axsom JE, Schmidt HD, Matura LA, Libonati JR. The influence of epigenetic modifications on metabolic changes in white adipose tissue and liver and their potential impact in exercise. Front Physiol. 2021;12:686270. https://doi.org/10.3389/fphys.2021.686270.
87. Chen S, Liu S, Shi S, Yin H, Tang Y, Zhang J, et al. Cross-species comparative DNA methylation reveals novel insights into complex trait genetics among cattle, sheep, and goats. Mol Biol Evol. 2024. https://doi.org/10.1093/molbev/msae003.