

A protein–foldamer supramolecular synthon for self-assembled hybrid architectures

Received: 14 November 2025

Accepted: 8 July 2026

Published online: 17 August 2026



Johannes Sigl¹, Vasily Morozov¹, Lingfei Wang¹, Janine Sachs², Eric Merlet², Eric Largy³, Niklas Geue^{4,5}, Nicklas Österlund^{4,5}, Sunbum Kwon^{1,2,7}, Florian Sanchez¹, Léo Candela⁶, Simon Huet⁶, Kevin Pagel^{4,5}, Yann Ferrand², Céline Douat¹, Cameron D. Mackereth³ & Ivan Huc¹✉

Constructing artificial assemblies that combine proteins and synthetic ligands has been hampered by the lack of protein–ligand interfaces that are sufficiently large and organized to enable precise structural control. Here ribosome display selection is used to identify a protein that binds a helical aromatic foldamer both tightly and selectively through a sizeable surface area. We used this complex as a supramolecular synthon to create well-defined hybrid foldamer–protein architectures. Examples include foldamers that bind two proteins and hold them at a precise distance, proteins that bind two foldamers and crystals in which proteins and foldamers are connected in cyclic or infinite arrays. The modularity of aromatic foldamers brings a further dimension to protein-based assemblies.

Artificial protein assemblies may be constructed by programming protein–protein interaction interfaces using either natural or de novo-designed proteins^{1–6}. Rationally organizing the spatial assembly of proteins using synthetic molecules⁶, instead of other proteins, is also desirable as illustrated by current efforts in the development of so-called molecular glues^{7,8}, proteolysis targeting chimeras (PROTACs)^{9,10} and hybrid protein frameworks^{11–13}. However, the co-assembly of proteins and synthetic molecules remains challenging to achieve with high structural precision. It has been limited by the lack of supramolecular synthons with a sufficiently large and organized protein–ligand interface for architectural control. Thus, synthetic macrocycles and molecular tweezers that bind to protein surfaces have been used to organize the packing of proteins in the solid state but their binding modes are variable^{11,13}. Conversely, small molecule ligands that selectively bind protein pockets may not prevent hinge motions, hampering the precise positioning of a large appendage such as another protein. Instead, flexible linkers are generally used, which reduces structural control⁸. In this Article, we introduce a complex between a synthetic helical aromatic foldamer and a 7-kDa artificial protein identified by ribosome display selection. The complex can serve as a supramolecular synthon for discrete aggregate and lattice construction. As solid-state and solution

structures reveal, the binding interface is sizeable ($>700 \text{ \AA}^2$), stable yet dynamic, and well-defined, allowing for the further elaboration of both the foldamer and protein components and for the construction of hybrid assemblies in which proteins are held at precise positions in space by synthetic foldamers.

Developing a supramolecular synthon that combines a protein and a synthetic molecule entailed a judicious choice of both components. Synthetic oligomers that fold—foldamers—provide a variety of medium-sized scaffolds many of which have shown potential to interact with proteins^{14,15}. In particular, aromatic oligoamide foldamers (AOFs) such as the oligoamides of 8-amino-2-quinoline carboxylic acid Q^{Xxx} (Fig. 1a) adopt particularly stable and predictable conformations and can be tailored to reach sizes that match those of small proteins¹⁶. For example, a simple octamer of Q^{Leu} remains helically folded at 120 °C in dimethyl sulfoxide¹⁷. The surface of AOFs can be decorated with biogenic-like side chains¹⁸ and their main chain can be programmed to form helices of variable diameter as well as sheets^{19,20}. Automated solid-phase synthesis (SPS) enables direct access to sequences in the 5–10-kDa range²¹. In principle, these combined features make AOFs suitable candidates for recognizing large surface areas of proteins. Efforts have until now focused on AOF design and screening, that is, on

¹Department of Pharmacy, Ludwig-Maximilians-Universität München, Munich, Germany. ²Univ. Bordeaux, CNRS, Bordeaux INP, CBMN, UMR 5248, Pessac, France. ³Univ. Bordeaux, CNRS, INSERM, ARNA, UMR 5320, U1212, Bordeaux, France. ⁴Department of Biology, Chemistry, and Pharmacy, Freie Universität Berlin, Berlin, Germany. ⁵Department of Molecular Physics, Fritz Haber Institute of the Max Planck Society, Berlin, Germany. ⁶Affilogic SAS, Nantes, France. ⁷Present address: Department of Chemistry, Chung-Ang University, Seoul, Republic of Korea. ✉e-mail: ivan.huc@cup.lmu.de




Fig. 1 | Foldamer building blocks, sequences and display selection results.

a, Monomer building blocks for AOFs with their respective letter code used in this study. Asp, aspartate-like side chain; Sul, sulfonate side chain; Pho, phosphonate-containing side chain; Orn, ornithine-like side chain; Dap, diaminopropionic acid-like side chain; Leu, leucine-like side chain; Phe, phenylalanine-like side chain; Hyd, hydroxy side chain; Dol, diol-containing side chain; Deg, diethylene glycol-containing side chain; Bio, biotin-containing terminal unit; PEG, diethylene glycol-containing terminal unit. Colours indicate the chemical nature of the side chains: orange, acidic; blue, basic; black, hydrophobic; green, polar neutral. Bold bonds define the inner rim of helical oligomers. **b**, AOF sequences synthesized, written left to right in the N- to C-terminal direction. In **4**, the unit D is a diacid, resulting in a symmetrical sequence with two C termini. **c**, Crystal structure of the dodecaamide **1b**, shown from side and top view. The right-

handed conformation is shown. The crystal also contains the left-handed helix. The bond order and hydrogen atoms are omitted for clarity. **d**, Crystal structure of the selected Nanofitin C10 at 1.4 Å (Supplementary Fig. 2) shown at the same scale as the structure of **1b** in **c**. Residues highlighted in pink, including five aromatic residues, are at positions randomized in the selection process. The coiled-coil extension was omitted for clarity. **e**, BLI sensorgrams of *P-1a* (blue curves) and *M-1a* (red curves) used as ligands on SA biosensors, against Nanofitin C10 (analyte) in a serial dilution from 250 nM to 62.5 nM in phosphate-buffered saline with 0.05% Tween 20 (PBST) revealing a dissociation constant (K_D) of 92 nM for *P-1a*–C10. **f**, Circular dichroism (CD) spectra of C10 recorded in ten steps heating the sample from 5 °C to 95 °C (top). Molar ellipticity (θ) recorded at 222 nm heating from 5 °C to 95 °C, then cooling back to 5 °C, showing complete CD signal recovery (bottom).

the identification of foldamers that bind to a biological protein target by varying helical²² or rod-like^{23,24} foldamer main chain and side chain components. However, structurally characterized complexes between proteins and helical AOFs are poorly stable (dissociation constant K_D in the micromolar range)²⁵ or involve flexible linkers proscribed for the current purpose²⁶. They thus do not constitute reliable supramolecular synthons. By contrast, the approach followed here consisted in identifying a protein that binds a given foldamer from a large library

of protein sequence variants, leading to a protein–foldamer complex in which both components are artificial.

To identify a protein that would tightly bind a foldamer, we turned to display selection methods, which brought their own challenges. Antibodies and nanobodies are typical proteins amenable to display selection. They have flexible loops that can create cavities, and successful examples of display selection of nanobodies against small molecules have been reported^{7,27}. However, their complex structure, relatively

large size and moderate stability are not ideal to serve as supramolecular synthons. Smaller proteins based on non-immunoglobulin scaffolds would fare better in this respect. Many such proteins have been used in display selection to bind to other protein targets^{28,29}. Yet examples of display selection against small synthetic molecules are rare; they typically involve loop binding regions rather than flat interfaces, and the structures of the complexes remain unknown^{30,31}. We opted for the selection of Nanofitins^{32,33}, which are based on the scaffold of Sac7d, a 66-residue cysteine-free protein stable over a broad range of temperature and pH³⁴. Sac7d naturally binds to dsDNA, whose size is similar to that of (Q^{xxx})_n helices. Binding is mediated by the slightly concave surface of an incomplete five-stranded β -barrel within which up to 14 residues have been randomly mutated for display selection. At their C-terminus, an α -helix caps the β -barrel and points away from the binding interface, thus providing a handle for fusion with other proteins. Furthermore, Nanofitin–protein complexes are amenable to X-ray crystallographic analysis^{35–37}.

Selection of a foldamer-binding protein

Biotinylated AOF **1a** (Fig. 1b), a Q^{xxx} dodecaamide five-turn helix, was chosen as the selection target because its chemically diverse side chain decoration potentially offers a variety of protein-binding epitopes, as highlighted by the X-ray crystallographic structure of its analogue **1b** (Fig. 1c)³⁸. Furthermore, the protein-binding capabilities of **1a** have been demonstrated in yeast lysate pull-down assays³⁸. Nanofitin selection was performed in four rounds of ribosome display, as previously described³², using biotinylated **1a** as a bait. The enrichment of naive libraries towards binders was monitored by enzyme-linked immunosorbent assay (ELISA) using the DNA pools collected after each round (Supplementary Fig. 1). Subsequent cloning and sequencing led to the identification of clone C10. Foldamer **1a** exists as a mixture of right-handed (*P*) and left-handed (*M*) diastereomeric helical conformers that do not interconvert and in which the chiral biotin has no effect on the proportion of *P* and *M* helices, which are thus present in identical amounts. The diastereoisomers can be separated by reversed-phase high-performance liquid chromatography (RP-HPLC) on a chiral stationary phase³⁸. C10 was recombinantly expressed and its affinity for *P*-**1a** and *M*-**1a** was assessed in parallel biolayer interferometry (BLI) assays. C10 was found to bind to *P*-**1a** with a *K*_D of 91.7 nM while no binding to *M*-**1a** was detected, hinting at a structurally well-defined interaction (Fig. 1e). We thus synthesized analogue **1c** whose N-terminal chiral C* camphanyl group quantitatively biases helix handedness to the *P* conformation³⁹. Initial attempts to crystallize C10 alone or in complex with **1b** or **1c** were unsuccessful. Yet by C-terminal fusion of the coiled-coil talin dimerization domain⁴⁰ (PDB 2QDQ) to C10, a high-resolution 1.4-Å X-ray crystallographic structure was obtained showing that the C10 apo-protein adopts the same tertiary fold as wild-type parent Sac7d (Fig. 1d, Supplementary Fig. 2 and Supplementary Table 1). The structure also showed that aromatic residues selected at randomly mutated positions (H22, F24, H26, Y33, H42) promote C10 dimerization in the solid state. Weak dimerization was also suspected in solution at concentrations required for NMR characterization (see below) and observed during size exclusion chromatography (SEC). Variable-temperature circular dichroism (CD) measurements highlighted the high thermostability of C10, showing a decrease of CD band intensity but no melting transition between 5 °C and 95 °C as well as complete CD signal recovery upon cooling (Fig. 1f), which boded well for its use in a supramolecular synthon.

NMR complex structure elucidation

A partial assignment of the NMR spectra of [¹³C, ¹⁵N]-labelled C10 was performed (Supplementary Fig. 3). The backbone amide signals of some residues were broadened and a few were missing. These were primarily residues from the region randomized during display selection. The broadening of these signals suggests that some aggregation involving

this region of the protein surface takes place at micromolar concentrations (Supplementary Fig. 3). By contrast, the spectra of solutions of the **1c**•C10 complex were sharp and its structure could be determined, allowing for an assessment of the nature of the binding mode in solution. Compared with usual ligands in protein–ligand complexes, the large size of the foldamer ligand necessitated the use of a modified structure determination process (see Methods for details). To assist further in identifying protein, foldamer and intermolecular cross-peaks, we used [¹³C, ¹⁵N]-labelled C10 combined with unlabelled foldamer **1c** at natural isotope abundance. This isotope labelling scheme allowed for the collection of edited or filtered NMR spectra so that signals of the protein only, of the foldamer only or of both the foldamer and protein are observed (Supplementary Figs. 4–6 and Supplementary Table 2). The resulting ensemble of 20 structures was calculated using restrained molecular dynamics in AMBER with 114 dihedral angle restraints and 737 distance restraints distributed throughout the complex (Supplementary Fig. 7 and Supplementary Tables 3 and 4).

The NMR structure of the **1c**•C10 complex revealed the architecture of the 1:1 assembly (Fig. 2a). The C-terminal aromatic cross-section of **1c** covers a large area of the protein β -sheet region (~830 Å², Fig. 2a), where most of the residues that were randomized during the selection process are located, while the N-terminus of the helix protrudes away from the protein. The interaction between the extended aromatic C-terminal region of **1c** with the flat C10 surface is somewhat reminiscent of G-quadruplex DNA–protein structures⁴¹. The upright binding mode of the foldamer helix on C10 differs from side-on helix orientations previously observed in more labile aromatic helical foldamer–protein complexes²⁵. In atomic detail, the **1c**•C10 binding interface primarily consists of (1) face-to-face π -stacking between the Q^{Dol}11 quinoline ring and H42 and T-shaped π – π interaction between Q^{Asp}12 and H26; (2) hydrophobic contacts between the two solvent-exposed terminal quinoline rings and residues V31 and L44, between the side chain of Q^{Dol}11 and Y33 and between Q^{Orn}9 and F24; (3) hydrogen bonding of Y33 to the main chain Q^{Dol}11–Q^{Asp}12 amide carbonyl and between T40 and the side chain of Q^{Dol}11; and (4) charge-reinforced hydrogen bonding of both Y8 and H42 to the C-terminal carboxylate of the foldamer and of H26 to the side chain carboxylate of Q^{Asp}12 (Fig. 2b–d). Interactions thus mainly involve residues located at the positions randomized during selection with the notable exception of Y8 and K9. The ensemble of 20 NMR structures determined for the **1c**•C10 complex shows limited variation in the relative position of the foldamer and protein, with an overall all-heavy atom RMSD of 1.7 ± 0.2 Å (Fig. 2a and Supplementary Table 3). Furthermore, molecular dynamics simulations of the complex confirm that the foldamer position and orientation on the C10 surface are essentially stable (Supplementary Video 1). The simulations also highlight the structural integrity of the complex in terms of residue fluctuations and hydrogen bond frequencies (Supplementary Figs. 8 and 9). Taken together, the solution structure data of **1c**•C10 validate the potential of this complex as a supramolecular synthon. Furthermore, the structure confirmed the location of the α -helix of C10 away from the foldamer-binding region and thus its potential use as a handle to fuse C10 with other proteins without altering foldamer binding.

Foldamer helix truncation

Large parts of the foldamer are not in contact with C10 in the complex, suggesting that a truncated version of **1** might bind equally well. We thus prepared pentaamide **2a** which comprises the five C-terminal units of **1**. For such short sequences, the equilibrium between *P* and *M* helices is expected to take place over the course of hours⁴². Indeed, upon mixing **2a** and C10, CD bands above 300 nm where only the foldamer absorbs were found to slowly emerge (Fig. 2f), indicating foldamer helix handedness bias upon binding to the protein. The emergence of the CD band occurred faster at 60 °C owing to enhanced helix dynamics, confirming that binding still takes place at such high temperatures.




Fig. 2 | Nuclear magnetic resonance (NMR) structure elucidation reveals the protein–foldamer complex binding epitope. **a**, NMR structure of the **1c**•C10 complex showing the aligned ensemble of 20 models. Structure statistics can be found in Supplementary Table 3. **b**, Schematic diagram of intermolecular contacts between C10 and the last four quinoline monomers of **1c** in the binding epitope identified with protein–ligand interaction profiler⁵⁰. Hydrogen bonds are shown as pink dashed lines and hydrophobic contacts as black solid lines. **c–e**, Three close-up structure views from different directions of important foldamer–protein contacts with annotated foldamer units and protein side chains.

Hydrogen bonds are shown as pink dashed lines, and hydrophobic contacts as black dashed lines. **f**, Time course of CD spectrum of pentameric sequence **2a** after addition of C10 at 30 μM, tris-buffered saline (TBS) pH 7.4, 25 °C. Note that the biotin moiety of **2a** does not bias foldamer helix handedness as seen in the flat curve for **2a** alone. **g**, BLI sensorgrams (blue solid lines) of C10 in solution and **2a** loaded on SA-biosensors and global fitting using a 1:1 binding model (black dashed lines). Kinetic experiments were performed in PBST at pH 7.4 with a serial dilution of C10 from 500 nM to 15.6 nM.

The positive band at 380 nm confirmed that the *P* helix is the favoured binding conformation³⁹. BLI measurements with biotinylated **2a** confirmed that it binds as well as **P-1a**. Note that helix handedness bias of **2a** is expected to remain marginal during the few minutes required for BLI measurements. We also prepared shorter foldamers having the last four, three or only two C-terminal units of **1a**. BLI and CD assays showed that strong binding was retained with four residues but worsened below that number (Supplementary Fig. 10). Altogether, these results support that foldamer–C10 interactions are limited to those revealed by the NMR structure and that C10 and a pentaamide foldamer helix provide

a robust supramolecular synthon for the elaboration of more complex architectures with controlled spatial organization.

Binding multiple proteins on the same foldamer

To test the generality and robustness of the supramolecular synthon upon modifications of the foldamer remote from the binding site, variants **3** and **4** were designed and synthesized (Fig. 1b), each presenting two binding epitopes through different modes of symmetric multimimerization. Variant **4** consists of a long *C*₂-symmetrical 31-mer single helix with two identical halves linked at their N termini by a central diacid unit




Fig. 3 | Binding multiple proteins on the same foldamer. **a**, Energy-minimized model of the C_2 -symmetrical $4\bullet(C10)_2$ complex. **b**, Size-exclusion chromatograms showing the formation of foldamer protein complexes as indicated by the lower elution volume for $4\bullet(C10)_2$ and $(P-3\bullet C10)_2$ compared with the protein alone. The split peak for $(P-3\bullet C10)_2$ was assigned to the presence of some $(P-3)_2\bullet C10$. **c,d**, Native mass spectrometry (MS) spectra of 1:2 and 1:1 mixtures of $4/C10$ and $P-3/C10$, respectively, at 100 μ M in 200 mM ammonium acetate pH 6.8 (top), detecting the formation of $4\bullet(C10)_2$ and $(P-3\bullet C10)_2$. The complexes were subjected to MSMS and collisional activation (bottom). Dashed arrows in the MSMS spectra indicate the dissociation produced from the m/z -selected

precursor ions. **e**, Collision cross-section in the gas phase ($^{TW}CCS_{N_2+He}$) of the $C10^{+5}$, $4\bullet(C10)_2^{+9}$ and $(P-3\bullet C10)_2^{+9}$ ions measured by ion mobility mass spectrometry. **f**, Asymmetric unit of the crystal structure of $(P-3\bullet C10)_2$ at 1.84 Å resolution containing two protein chains bridged by the foldamer duplex $(P-3)_2$. A pseudo (non-crystallographic) C_2 axis is indicated. **g**, Two different views of the C_4 -symmetrical higher order assembly observed in the $(P-3\bullet C10)_2$ X-ray crystallographic structure. The cyclic 8+8 aggregate comprised of four $(P-3\bullet C10)_2$ complexes is stabilized by inter-duplex π - π contacts between Q_{OMe} units and side chain-mediated charge-reinforced hydrogen bonds.




Fig. 4 | Binding multiple foldamers by protein dimerization. **a**, Crystal structure of the coil-extended C10 (C10CE) in complex with pentaamide **2b**. The ASU is shown from two perspectives, illustrating the cyclic arrangement of two foldamers and four C10CE proteins in the solid state. **b**, Detailed view of the foldamer packed between two C10 molecules, with the cognate-binding epitope

highlighted in teal on one protein and non-cognate interactions in purple on the opposite chain. **c**, Highlights of Glu, Lys and Arg inter-coiled-coil interactions, connecting two protein dimers. Residues marked with a star were reconstituted from the crystal structure and energy minimized.

D. Each half of **4** thus possesses a C-terminal C10-binding pentaamide elongated at its N terminus by a B* chiral unit, a number of hydrophilic Q^{Orn} and Q^{Deg} residues and a P residue, all assembled during SPS (Supplementary Fig. 11). The B* chiral control unit quantitatively biases helix handedness to the *P* conformation⁴³, and the benzylic amine of the *P* unit facilitates the post-SPS assembly of **4** by fragment condensation on the diacid of unit D activated as a bis-*N*-hydroxysuccinimide ester²⁵. The complexation of **4** with C10 was analysed by SEC. It resulted in a decrease of the retention time, indicating the formation of a presumably dumbbell-shaped ternary complex (Fig. 3a,b). The complex was also analysed in the gas phase using native nano-electrospray ionization mass spectrometry (nESI-MS). The 4•(C10)₂ complex was found to be the dominant species, and its fragmentation into C10+4•C10 could be monitored by tandem mass spectrometry (MSMS, Fig. 3c). Analogues of **4** having longer or shorter helices are thus expected to rigidly hold in space two C10 molecules (and any protein fused to C10) at a distance and twist angle precisely defined by the number of AOF residues.

Variant **3** uses an N-terminal extension of the pentaamide binding epitope with four 7-amino-8-methoxy quinoline carboxylic acid Q_{OMe} residues that code for a wider aromatic helix diameter, which in turn promotes dimerization into stable antiparallel double helices in water⁴⁴. The resulting duplex of sequence **3** binds two copies of C10, and the 2+2 quaternary complex can also be traced by SEC, showing an elution volume smaller than that of the protein or foldamer alone, yet larger than that of 4•(C10)₂, reflecting the effect of foldamer length on elution time (Fig. 3b). In the gas phase, the (P-3•C10)₂ complex was also observed by native nESI-MS (Fig. 3d) to be the dominant species. Its main fragmentation in MSMS was through a loss of a C10 unit and the formation of (P-3)•C10, and not through the disruption of the (P-3)₂ double helix highlighting its robustness. Furthermore, ion mobility measurements gave access to the collision cross-sections of C10, (P-3•C10)₂ and 4•(C10)₂, showing a trend consistent with the SEC data (Fig. 3e). Note that helix handedness control was not implemented in **3** and that, given its length, its helical conformation is kinetically locked (no CD band emerges upon adding C10 to **3**). In solutions of racemic (**3**)₂ duplexes, only the *P* helices thus bind to C10. We successfully crystallized and solved the structure of the (P-3•C10)₂ complex, thus resolving the racemic (**3**)₂ duplexes by crystallization. The structure at 1.84 Å resolution confirmed the expected dumbbell geometry, with two C10 molecules spatially separated upon binding to the opposite sides of double helical (**3**)₂ (Fig. 3f, Supplementary Video 2 and Supplementary Table 1). The asymmetric unit (ASU) contains

two independent foldamer–C10 interfaces which show highly conserved interactions essentially identical to those of the NMR structure (Supplementary Figs. 12 and 13, Supplementary Table 5), highlighting again the robustness of the supramolecular synthon.

The X-ray crystallographic structure of C10 bound to the foldamer was also compared with that of the apo-C10, showing that the main chains are essentially superimposable, an illustration of the robustness of the Nanofitin scaffold, while surface side chains, including those involved in foldamer recognition, may have different orientations (Supplementary Fig. 12). Future developments may exploit the use of longer helical segments to adjust protein–protein distance–foldamer length spontaneously doubles upon duplex formation, and the fact that heteromeric foldamer double helices form selectively when complementary strands are used⁴⁴.

In the solid state, the (P-3•C10)₂ complexes form a higher-order tetrameric [(P-3•C10)₂]₄ assembly organized in a fourfold cyclic arrangement (Fig. 3g and Supplementary Video 3). The assembly is held together by face-to-face contacts between Q_{OMe} residues belonging to different duplexes and an extensive array of inter-duplex charge-reinforced hydrogen bonds (Supplementary Fig. 14). In solution, the 8+8 aggregate was not detected by electron microscopy, and SEC retention volume shifts seem to be too small for an ~80-kDa object. It was also not visible in native nESI-MS spectra measured from micromolar solutions (Fig. 3d). We concluded that the assembly forms only at the mM concentration of the crystallization drops and possibly represents a minor component in solution, even at these higher concentrations, that would only be stabilized in the solid state. It nevertheless represents a starting point for designing stable foldamer assemblies able to bind multiple—here eight—proteins.

Binding multiple foldamers on a protein dimer

We next investigated modifications on C10 to explore how to spatially organize the supramolecular synthon via multimerization of the protein. Coil coiling of α-helices is arguably the best understood aggregation pattern and the easiest to design^{1,2,45}. Implementation of a coiled coil on C10 is straightforward, thanks to its C-terminal α-helix that points away from the foldamer-binding region. We opted again for a simple fusion with the antiparallel coiled-coil domain of talin mentioned above, albeit with some design improvements. In the first design, two consecutive glycine residues (G66&67) present at the junction between C10 and the coiled coil were responsible for a local disruption of α-helix that kinked the structure (Supplementary Fig. 1). In this




Fig. 5 | Crystal structure of the coiled-coil-mediated C10CE dimer in complex with the C_2 symmetrical foldamer **4 ($4 \cdot (C10CE)_2$) at 2.53 Å. **a**, Continuous ribbon built of protein dimers connected to the C_2 -symmetrical foldamer **4**. The ASU (highlighted in gold) contains one protein monomer and one-half of **4**. The imposed local C_2 axes of the complex coincide with crystallographic**

symmetry axes, giving rise to a highly ordered assembly in the space group $I4_122$. **b–d**, Orthoscopic views of the crystal lattice along different directions: along (010) (**b**), along (110) (**c**) and along (001) (**d**). In all cases, the unit cell is shown in black. Proteins are shown as blue ribbons and foldamers as red sticks.

second design, these two glycine residues were replaced by alanine residues to better promote α -helicity. Crystallization efforts with this coil-extended C10 construct (C10CE) in complex with the initial target sequence, for example, **1b** or **1c**, repeatedly yielded crystals of the foldamer alone, presumably owing to its lower solubility. Therefore, pentamer **2b** (Fig. 1b) was synthesized, carrying a solubilizing PEG termination as well as two phosphonate-containing Q^{Pho} residues that were expected to improve solubility with respect to the original carboxylate Q^{Asp} units. Native nESI-MS showed the formation of the **2b**•C10CE complex along with the (**2b**•C10CE)₂ dimer. The proportions also seen in the spectrum of C10CE alone suggested some lability of the coiled-coil dimerization in the gas phase. The (**2b**•C10CE)₂ dimer was found to fragment via coiled-coil dissociation (Supplementary Figs. 15, 16).

Diffraction quality crystals of the C10CE in complex with **2b** were eventually obtained, and the X-ray crystallographic structure was elucidated (Fig. 4, Supplementary Fig. 17 and Supplementary Table 1). In the ASU, two coiled-coil protein dimers are in complex with two molecules of **2b** through their C-terminal cross-section, providing further evidence that this robust, cognate interaction is highly conserved (Fig. 4b, Supplementary Fig. 13 and Supplementary Table 5). In addition, the two C10 domains establish contacts with the N-terminal cross-section of the **2b** helices (Fig. 4b). This interaction only concerns the N-terminal quinoline ring and its short PEG chain, which involve hydrophobic contacts with Y33, H22 and T40 and a charge-reinforced hydrogen bond between H42 and a phosphonate side chain. We presume that they are weaker than the cognate foldamer–C10 recognition and only occur at the mM concentration of crystallization drops at substoichiometric amounts of foldamer. There is thus little risk that such non-cognate interactions

would compete with the 91.7-nM cognate C10–foldamer recognition. The 4+2 cyclic assembly—four proteins and two foldamers—mediated by the non-cognate interactions was not detected by native nESI-MS from micromolar solutions (Supplementary Fig. 15). It is possibly a crystallization artefact—it crystallizes although it represents a minor component in solution. It may also be the result of a substoichiometric amount of foldamer. Nevertheless, the outcome is an X-shaped, figure-of-8, discrete macrocyclic assembly between two foldamers and two C10CE coiled-coil dimers (Fig. 4a, Supplementary Video 4). At the centre of the cyclic structure, favourable contacts between the two coiled coils are mediated by hydrogen bonds between multiple Glu, Lys and Arg residues that may be amenable to optimization (Fig. 4c). Altogether, the structure validates, and goes beyond, the initial design principles, and it suggests directions for further developments.

Binding multiple proteins with multiple foldamers in one system

Based on the established principles of enforced multimerization around the foldamer–C10 supramolecular synthon, we lastly combined the rigid homochiral divalent single helix of **4** with the head-to-tail coiled-coil-mediated dimer of C10CE. Crystallization trials quickly yielded diffraction quality crystals and revealed a structure of high symmetry, with the ASU containing one protein monomer and half of the C_2 -symmetrical **4** (Fig. 5, Supplementary Table 1). This structure provided yet one more illustration of the conservation of C10–foldamer interactions (Supplementary Fig. 1–3, Supplementary Table 5). Through application of the symmetry operations, a 1D polymeric zig-zag-like network that extends in a sort of flat ribbon is built (Fig. 5a). The ribbon consists mainly

of cognate foldamer–C10-binding interfaces and the α -helical coiled coil included in the design, along with C10–C10 contacts associated with crystal packing. The coincidence of the symmetry elements of the building blocks with the crystallographic rotational axes can be interpreted as an enforcement of global symmetry by predisposed local symmetries⁴⁶. Here the C_2 symmetry axis of the foldamer and the C_2 symmetry axis of the C10 coiled-coil dimer are perpendicular, hence the formation of a flat assembly, but they do not intersect, giving rise to a polymeric structure rather than a macrocycle. This assembly is the direct outcome of the distance and angular twist between the protein-binding epitopes of **4** encoded in the number of Q^{xxx} residues and of the distance and angular twist between the foldamer-binding epitopes of the C10 coiled-coil dimer encoded by the length of the α -helices. In the gas phase, the 1D polymer was not observed by native nESI-MS (Supplementary Fig. 16).

While the crystal lattice appears to be tightly packed when viewed down its c -axis (Fig. 5d), it is in fact highly porous, with the foldamer spacing the proteins ~4.3 nm apart, while the coiled-coil domain keeps the foldamers ~8 nm apart (Fig. 5a–c). Computational analysis of the pores showed that the resulting channels in a and b directions reach a bottleneck radius of 15.5 Å and that the largest cavities can accommodate theoretical spheres with a diameter of ~5 nm (Supplementary Fig. 18)⁴⁷, a value comparable to that of other highly porous frameworks¹¹. A typical way to assess solvent content in protein crystal structures is through the Matthews coefficient, which describes the unit cell volume per protein mass (but does not tell anything about pore size). The mean Matthews coefficient of Protein Data Bank (PDB) protein structures is $2.67 \text{ Å}^3 \text{ Da}^{-1}$ (ref. 48) to be compared with $3.51 \text{ Å}^3 \text{ Da}^{-1}$ for the porous lattice mentioned above¹¹ and $5.13 \text{ Å}^3 \text{ Da}^{-1}$ for the structure of **4**•(C10CE)₂. This contrasts with the low porosity of the structure obtained with the pentaamide helix **2b**•(C10CE)₂ (Fig. 4), which has a calculated maximum bottleneck size of 8.6 Å, and reflects the benefit of the space between proteins imposed by **4**. Considering that the length of the foldamer is freely adjustable (within synthetic limits), spacing within such lattices is, in principle, tunable, highlighting the relevance of protein–synthetic foldamer assemblies in the context of porous solid design.

In summary, in contrast to smaller molecules or peptides, aromatic foldamers constitute large and rigid-enough synthetic molecules to allow for successful display selection of protein binders and the generation of a well-defined and robust binding interface. It appears that the foldamer aromatic cross-section acted as a hot spot—one such protein–foldamer contact had been observed before with carbonic anhydrase II (ref. 49), and this feature may be exploited in future studies to target proteins with flat surfaces. The identification of C10 also suggests that targets other than aromatic foldamers may be amenable to the selection of protein binders, to bridge further the synthetic and protein chemical spaces. In turn, both the foldamer and protein can be further designed without altering their binding interactions so as to produce discrete or extended aggregates and lattices combining both biogenic and synthetic components. Extensions of this work will take advantage of the inherent modularity and synthetic accessibility of foldamers, of the diversity of protein scaffolds amenable to display selection such as the Nanofitin described in our study²⁹ and of the advanced computational approaches to design protein structures and interaction^{1–3}. These extensions include the design, either structure-guided or using *ab initio* protein design tools, of foldamer–protein nanocages and porous 3D lattices in which the foldamers may bring functional groups not available to proteins. They also included the use of foldamers to bring together or, on the contrary, to keep at a well-defined distance, naturally occurring proteins that have been tagged with foldamer-binding domains to alter their biological functions.

Online content

Any methods, additional references, Nature Portfolio reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions

and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41557-026-02222-6>.

References

- Regan, L. & DeGrado, W. F. Characterization of a helical protein designed from first principles. *Science* **241**, 976–978 (1988).
- Thomson, A. R. et al. Computational design of water-soluble α -helical barrels. *Science* **346**, 485–488 (2014).
- Hsia, Y. et al. Design of a hyperstable 60-subunit protein icosahedron. *Nature* **535**, 136–139 (2016).
- Suzuki, Y. et al. Self-assembly of coherently dynamic, auxetic, two-dimensional protein crystals. *Nature* **533**, 369–373 (2016).
- Lai, Y. T., Cascio, D. & Yeates, T. O. Structure of a 16-nm cage designed by using protein oligomers. *Science* **336**, 1129–1129 (2012).
- Luo, Q., Hou, C., Bai, Y., Wang, R. & Liu, J. Protein assembly: versatile approaches to construct highly ordered nanostructures. *Chem. Rev.* **116**, 13571–13632 (2016).
- Shui, S., Buckley, S., Scheller, L. & Correia, B. E. Rational design of small-molecule responsive protein switches. *Protein Sci.* **32**, e4774 (2023).
- Soini, L., Leysen, S., Davis, J. & Ottmann, C. Molecular glues to stabilise protein–protein interactions. *Curr. Opin. Chem. Biol.* **69**, 102169 (2022).
- Liu, Z. et al. An overview of PROTACs: a promising drug discovery paradigm. *Mol. Biomed.* **3**, 46 (2022).
- Troup, R. I., Fallan, C. & Baud, M. G. J. Current strategies for the design of PROTAC linkers: a critical review. *Explor. Target. Anti-tumor Ther.* **1**, 273–312 (2020).
- Engilberge, S., Rennie, M. L., Dumont, E. & Crowley, P. B. Tuning protein frameworks via auxiliary supramolecular interactions. *ACS Nano* **13**, 10343–10350 (2019).
- van Dun, S., Ottmann, C., Milroy, L. G. & Brunsveld, L. Supramolecular chemistry targeting proteins. *J. Am. Chem. Soc.* **139**, 13960–13968 (2017).
- Mockler, N. M., Raston, C. L. & Crowley, P. B. Making and breaking supramolecular synthons for modular protein frameworks. *Chem. Eur. J.* **31**, e202500732 (2025).
- Gopalakrishnan, R., Frolov, A. I., Knerr, L., Drury, W. J. III & Valeur, E. Therapeutic potential of foldamers: from chemical biology tools to drug candidates? *J. Med. Chem.* **59**, 9599–9621 (2016).
- Pasco, M., Dolain, C. & Guichard, G. in *Comprehensive Supramolecular Chemistry II* 89–125 (Elsevier, 2017).
- Wang, S., Sigl, J., Allmendinger, L., Maurizot, V. & Huc, I. Design of an abiotic unimolecular three-helix bundle. *Chem. Sci.* **16**, 1136–1146 (2025).
- Jiang, H., Leger, J. M. & Huc, I. Aromatic delta-peptides. *J. Am. Chem. Soc.* **125**, 3448–3449 (2003).
- Zwillinger, M. et al. Development of aromatic foldamer building blocks bearing multiple biogenic side chains. *J. Org. Chem.* **90**, 3043–3052 (2025).
- Gong, B. et al. Creating nanocavities of tunable sizes: hollow helices. *Proc. Natl Acad. Sci. USA* **99**, 11583–11588 (2002).
- Atcher, J. et al. Large-amplitude conformational changes in self-assembled multi-stranded aromatic sheets. *Angew. Chem. Int. Ed.* **60**, 2574–2577 (2021).
- Corvaglia, V., Sanchez, F., Menke, F. S., Douat, C. & Huc, I. Optimization and automation of helical aromatic oligoamide foldamer solid-phase synthesis. *Chem. Eur. J.* **29**, e202300898 (2023).
- Wang, L. et al. Structure-based design of an aromatic helical foldamer–protein interface. *Chem. Sci.* **16**, 12385–12396 (2025).
- Azzarito, V. et al. Stereocontrolled protein surface recognition using chiral oligoamide proteomimetic foldamers. *Chem. Sci.* **6**, 2434–2443 (2015).

24. Stillman, N. H. et al. Protein mimetic 2D FAST rescues alpha synuclein aggregation mediated early and post disease Parkinson's phenotypes. *Nat. Commun.* **15**, 3658 (2024).
25. Deepak, D. et al. DNA mimic foldamer recognition of a chromosomal protein. *Angew. Chem. Int. Ed.* **64**, e202422958 (2025).
26. Reddy, P. S. et al. Structure elucidation of helical aromatic foldamer–protein complexes with large contact surface areas. *Chem. Eur. J.* **25**, 11042–11047 (2019).
27. Kang, S. et al. COMBINES-CID: an efficient method for de novo engineering of highly specific chemically induced protein dimerization systems. *J. Am. Chem. Soc.* **141**, 10948–10952 (2019).
28. Gebauer, M. & Skerra, A. Engineered protein scaffolds as next-generation antibody therapeutics. *Curr. Opin. Chem. Biol.* **13**, 245–255 (2009).
29. Helma, J., Cardoso, M. C., Muyldermans, S. & Leonhardt, H. Nanobodies and recombinant binders in cell biology. *J. Cell Biol.* **209**, 633–644 (2015).
30. Arrata, I., Barnard, A., Tomlinson, D. C. & Wilson, A. J. Interfacing native and non-native peptides: using Affimers to recognise α -helix mimicking foldamers. *Chem. Commun.* **53**, 2834–2837 (2017).
31. Wang, L. et al. Display selection of peptide ligands for helical aromatic foldamers. *Org. Biomol. Chem.* **23**, 4641–4647 (2025).
32. Huet, S., Gorre, H., Perrocheau, A., Picot, J. & Cinier, M. Use of the nanofitin alternative scaffold as a GFP-ready fusion tag. *PLoS ONE* **10**, e0142304 (2015).
33. Mouratou, B. et al. Remodeling a DNA-binding protein as a specific in vivo inhibitor of bacterial secretin PulD. *Proc. Natl Acad. Sci. USA* **104**, 17983–17988 (2007).
34. Robinson, H. et al. The hyperthermophile chromosomal protein Sac7d sharply kinks DNA. *Nature* **392**, 202–205 (1998).
35. Garlich, J. et al. Discovery of APL-1030, a novel, high-affinity nanofitin inhibitor of C3-mediated complement activation. *Biomolecules* **12**, 432 (2022).
36. Correa, A. et al. Potent and specific inhibition of glycosidases by small artificial binding proteins (affitins). *PLoS ONE* **9**, e97438 (2014).
37. Behar, G. et al. Tolerance of the archaeal Sac7d scaffold protein to alternative library designs: characterization of anti-immunoglobulin G Affitins. *Protein Eng. Des. Sel.* **26**, 267–275 (2013).
38. Kwon, S. et al. Interrogating the potential of helical aromatic foldamers for protein recognition. *Org. Biomol. Chem.* **22**, 9342–9347 (2024).
39. Kendhale, A. M. et al. Absolute control of helical handedness in quinoline oligoamides. *J. Org. Chem.* **76**, 195–200 (2011).
40. Gingras, A. R. et al. The structure of the C-terminal actin-binding domain of talin. *EMBO J.* **27**, 458–469 (2008).
41. Chen, M. C. et al. Structural basis of G-quadruplex unfolding by the DEAH/RHA helicase DHX36. *Nature* **558**, 465–469 (2018).
42. Vallade, M. et al. Assessing interactions between helical aromatic oligoamide foldamers and protein surfaces: a tethering approach. *Bioconjugate Chem.* **30**, 54–62 (2019).
43. Bindl, D., Heinemann, E., Mandal, P. K. & Huc, I. Quantitative helix handedness bias through a single H vs. CH3 stereochemical differentiation. *Chem. Commun.* **57**, 5662–5665 (2021).
44. Koehler, V. et al. High-affinity hybridization of complementary aromatic oligoamide strands in water. *Angew. Chem. Int. Ed.* **62**, e202311639 (2023).
45. Gradišar, H. et al. Design of a single-chain polypeptide tetrahedron assembled from coiled-coil segments. *Nat. Chem. Biol.* **9**, 362–366 (2013).
46. Laniado, J. & Yeates, T. O. A complete rule set for designing symmetry combination materials from protein molecules. *Proc. Natl Acad. Sci. USA* **117**, 31817–31823 (2020).
47. Pletzer-Zelgert, J. et al. LifeSoaks: a tool for analyzing solvent channels in protein crystals and obstacles for soaking experiments. *Acta Crystallogr. D.* **79**, 837–856 (2023).
48. Shao, C., Liu, Z., Yang, H., Wang, S. & Burley, S. K. Outlier analyses of the Protein Data Bank archive using a probability-density-ranking approach. *Sci. Data* **5**, 180293 (2018).
49. Buratto, J. et al. Structure of a complex formed by a protein and a helical aromatic oligoamide foldamer at 2.1 Å resolution. *Angew. Chem. Int. Ed. Engl.* **53**, 883–887 (2014).
50. Schake, P., Bolz, S. N., Linnemann, K. & Schroeder, M. PLIP 2025: introducing protein–protein interactions to the protein–ligand interaction profiler. *Nucleic Acids Res.* **53**, W463–W465 (2025).

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2026

Methods

Detailed methods are provided in the Supplementary Information.

Foldamer synthesis

All foldamers were obtained by manual or automated solid-phase supported synthesis from Fmoc-amino acid monomers—Fmoc-Q^{xxx}-OH, Fmoc-Q^{OMe}-OH, Fmoc-P-OH and Fmoc-B^{*}-OH—bearing suitable acid-labile side chain-protecting groups, for example, Boc for Q^{Orn}, Q^{Dap} and Q^{OMe} (Fig. 1a). The monomers used in this study are not commercially available and were synthesized following well-established protocols. In general, the carboxylic acid function of the monomers was activated as its acid chloride by applying in situ Appel's conditions in the presence of triphenylphosphine and trichloroacetonitrile. Coupling reactions to the free amine of an oligomer were performed in the presence of 2,4,6-collidine as a base. Subsequent deprotection of the main chain amine, that is, Fmoc removal, was carried out with 2% DBU (1,8-diazabicyclo[5.4.0]undec-7-ene) in *N*-methyl-2-pyrrolidone. Removal of the silyl ether protection of Q^{Dol} was carried out on-resin with 1 M tetrabutylammonium fluoride in THF. Cleavage from the solid support was performed using a mixture of TFA, triisopropylsilane and H₂O (95:2.5:2.5; v/v/v) for Wang resin, thereby leading to the deprotection of the other side chains, or with 1% TFA in dichloromethane for SASRIN resin, leaving the other side chain-protecting groups intact. Fragment condensation en route to the C₂-symmetrical helical foldamer **4** was performed in DMF with the di-*N*-hydroxysuccinimide active ester of pyridine-2,6-dicarboxylic acid **D** and diisopropylethylamine as a base. All foldamer sequences were purified by RP-HPLC.

Nanofitin selection

The combinatorial library of Nanofitins for ribosome display was assembled by two successive overlapping PCRs from degenerate oligonucleotides encoding NNS triplets (N=A, C, T or G and S=C or G). Finally, a PCR step added the 5'- and 3'-flanking regions necessary for ribosome display. The PCR-amplified library was transcribed, and the selection was performed in four rounds at 4 °C against 15 pmol of the immobilized biotinylated foldamer **1a**. Amplified DNA material from the last round was cloned between BamHI and HindIII restriction sites of a plasmid derived from pQE-30 (Qiagen), and the ligation mixture was transformed into *Escherichia coli* DH5α LacIq strains (Invitrogen). Expression of the Nanofitin clones was induced by the addition of isopropyl β-D-1-thiogalactopyranoside (IPTG). The selection target **1a** was immobilized on streptavidin (SA) MaxiSorp plates, and crude *E. coli* extracts were applied to wells with and without immobilized antigen for 1 h. Detection was carried out by the addition of an RGS-His antibody HRP conjugate (Qiagen), followed by the addition of *o*-phenylenediamine dihydrochloride substrate (Sigma-Aldrich) solution. Absorbance at 450 nm was measured using a Varioskan enzyme-linked immunosorbent assay plate reader (Thermo Scientific), revealing Nanofitin C10 as a binder with a strong signal-to-background ratio.

Nanofitin expression and purification

Nanofitin C10 and the coil-extended versions were expressed recombinantly in *E. coli* BL21(DE3) as N-terminally His-tagged or MBP-His-tagged fusion proteins. Plasmids (pET-28a(+), pMAL-cSE) were purchased ready to use as lyophilized solids (GenScript). For unlabelled proteins, cultures were grown in LB. For the expression of the [¹³C, ¹⁵N]-isotope-labelled C10, M9 minimal medium supplemented with D-glucose-¹³C₆, ¹⁵NH₄Cl, and trace element solutions was used. Overexpression was induced with IPTG. Purification was carried out by Ni-NTA IMAC. Tag removal was achieved by cleavage with HRV-3C protease overnight. Final purification was carried out by SEC.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

X-ray crystallographic structures have been deposited at the PDB under 9QDO, 9QT7, 9QOG and 9QNU. The ensemble of 20 structures calculated for the **1c**•C10 complex has been deposited at the PDB under accession 9T00. The chemical shift assignments for the unbound C10 and **1c**•C10 complex were deposited in the BMRB (<https://bmr.bio.org/>) under accession numbers 53300 and 53317, respectively. The raw data that served to create the graphs in Figs. 1, 2 and 3 as well as BLI data have been deposited at Open Data LMU under https://data.ub.uni-muenchen.de/701/1/C10_BLI-raw_Data.zip (<https://doi.org/10.5282/ubm/data.818>). All other data are available in the article or the Supplementary Information. Source data are provided with this paper.

Acknowledgements

The synchrotron data were collected at beamlines P13 and P14 operated by EMBL Hamburg at the PETRA III storage ring (DESY, Hamburg, Germany). We would like to thank D. von Stetten, G. Grebnev, I. Bento and S. Storm (all from European Molecular Biology Laboratory, Hamburg) for the assistance in using the beamlines. We thank M. Loos, I. Alonso and M. Vançon (all of the Department of Pharmacy, Ludwig-Maximilians-Universität München) for helping with the synthesis of short oligomers as well as the Max Planck Institute for Biochemistry for granting access to their crystal broad screening facility. We thank the Core Facility BioSupraMol, Freie Universität Berlin, for instrument time and support. This work benefited from the facilities and expertise of the Biophysical and Structural Chemistry platform (BPCS) at IECB, CNRS UMS3033, Inserm US001 and Bordeaux University. We thank Axelle Grélard and Estelle Morvan (both from BPCS) for access to the NMR spectrometers and technical assistance.

Author contributions

I.H. and J. Sigl conceived the project. V.M., C.D.M., C.D. and S.H. developed the methodology. J. Sigl, V.M., L.W., J. Sachs, E.M., E.L., N.G., N.Ö., S.K., F.S., L.C., S.H., C.D. and C.D.M. carried out the investigation. I.H., Y.F. and C.D.M. acquired funding for the study. The project was supervised by I.H., Y.F., C.D.M., S.H. and K.P. J. Sigl, C.D.M., C.D. and I.H. wrote the original draft of the paper. All authors reviewed and edited the paper.

Funding

This study was funded by the Free State of Bavaria under the Excellence Strategy of the Federal Government and the Länder through the ONE MUNICH Project Munich Multiscale Biofabrication (I.H.) and by CNRS-LMU IRP Foldsfun (Y.F. and I.H.). A Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Walter Benjamin grant (559720072) was awarded to N.G. A Wenner-Gren Foundation fellowship was awarded to N.Ö. Open access funding provided by Ludwig-Maximilians-Universität München.

Competing interests

L.C. and S.H. are employees of Affilogic. The other authors declare that they have no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41557-026-02222-6>.

Correspondence and requests for materials should be addressed to Ivan Huc.

Peer review information *Nature Chemistry* thanks the anonymous reviewers for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- ☒ ☐ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☒ ☐ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☒ ☐ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☒ ☐ A description of all covariates tested
- ☒ ☐ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☒ ☐ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☒ ☐ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection NMR data were collected using Topspin version 4.

Data analysis

Data Analysis/Collection, Processing & Modelling

- * NMRPipe: Processing of multidimensional NMR spectra.
- * Sparky (Version 3.190) / NMRFAM-SPARKY (Version 1.470): Analysis and assignment of NMR spectra.
- * OriginPro (Version V.2019b): Processing and plotting of chromatography, CD spectroscopy, and BLI data.
- * Octet Analysis Studio (Version V.13): Processing of BLI data and calculation of kinetic binding values.
- * CrysAlisPro: Processing of X-ray diffraction data.
- * XDS: Processing of X-ray diffraction data.
- * PHASER: Solution of crystal structures via molecular replacement.
- * PHENIX refine: Execution of iterative crystallographic structure refinement.
- * COOT: Facilitation of real-space model building and evaluation of electron density.
- * elBOW: Generation of ligand coordinate and geometry restraint files.
- * Grade2: Generation of ligand geometry restraints.
- * PRODRG: Creation of parameter and topology files for foldamers.
- * ARIA (Version 2.3): Iterative assignment of NOESY spectra and calibration of distance restraints in conjunction with CNS.
- * CNS (Version 1.2): Calculation of initial NMR structures from linear conformations.
- * TALOS-N: Prediction of protein backbone dihedral angles from NMR chemical shifts.
- * Antechamber: Generation of partial charges and AMBER-compatible atom types.
- * prepren: Generation of topology and parameter prepin files.
- * parmchk2: Retrieval of force field parameters for molecular dynamics.

- * Multiwfn (Version 3.8): Execution of RESP charge calculations.
- * Orca (Version 6.0): Execution of single-point energy calculations applying density functional theory (DFT).
- * Amber (Version 24): Execution of restrained molecular dynamics simulations.
- * Leap: Preparation of complexes for MD simulations within Amber.
- * pmemd: Execution of energy minimization steps on CPU.
- * pmemd.cuda: Execution of MD production runs on GPU.
- * cpptraj (Version 6.18.1): Analysis of molecular dynamics trajectories.
- * PLIP (Version 2.3.0): Profiling of protein-ligand interactions.
- * PyMOL (Version 3.1.6): Visualization of molecules, alignment of structures, and calculation of solvent-accessible surface areas.
- * ChemDraw (Version 25.0): Depiction of chemical structures.
- * R (Version 4.5.1): Analysis of trajectory and ensemble data.
- * Maestro MacroModel (Version 11.5): Execution of molecular modeling and energy minimizations.
- * LifeSoaks: Calculation of pore bottlenecks via the proteins.plus interface.

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

Data and materials availability: Crystallographic structures have been deposited at the Protein Data Bank under 9QDO, 9QT7, 9QOG & 9QNU. The ensemble of 20 structures calculated for the 1c-C10 complex have been deposited at the PDB under accession 9T00. The chemical shift assignments for the unbound C10 and 1c•C10 complex were deposited in the BMRB (<https://bmr.io/>) under accession numbers 53300 and 53317, respectively. BLI data have been deposited at Open Data LMU under https://data.ub.uni-muenchen.de/701/1/C10_BLI-raw_Data.zip. All other data are available in the main text or the supplementary materials.

Data and materials availability: Crystallographic structures have been deposited at the Protein Data Bank under 9QDO, 9QT7, 9QOG & 9QNU. The ensemble of 20 structures calculated for the 1c-C10 complex have been deposited at the PDB under accession 9T00. The chemical shift assignments for the unbound C10 and 1c•C10 complex were deposited in the BMRB (<https://bmr.io/>) under accession numbers 53300 and 53317, respectively. BLI data have been deposited at Open Data LMU under https://data.ub.uni-muenchen.de/701/1/C10_BLI-raw_Data.zip. All other data are available in the main text or the supplementary materials.

Research involving human participants, their data, or biological material

Policy information about studies with [human participants or human data](#). See also policy information about [sex, gender \(identity/presentation\), and sexual orientation](#) and [race, ethnicity and racism](#).

Reporting on sex and gender

Use the terms sex (biological attribute) and gender (shaped by social and cultural circumstances) carefully in order to avoid confusing both terms. Indicate if findings apply to only one sex or gender; describe whether sex and gender were considered in study design; whether sex and/or gender was determined based on self-reporting or assigned and methods used. Provide in the source data disaggregated sex and gender data, where this information has been collected, and if consent has been obtained for sharing of individual-level data; provide overall numbers in this Reporting Summary. Please state if this information has not been collected. Report sex- and gender-based analyses where performed, justify reasons for lack of sex- and gender-based analysis.

Reporting on race, ethnicity, or other socially relevant groupings

Please specify the socially constructed or socially relevant categorization variable(s) used in your manuscript and explain why they were used. Please note that such variables should not be used as proxies for other socially constructed/relevant variables (for example, race or ethnicity should not be used as a proxy for socioeconomic status). Provide clear definitions of the relevant terms used, how they were provided (by the participants/respondents, the researchers, or third parties), and the method(s) used to classify people into the different categories (e.g. self-report, census or administrative data, social media data, etc.) Please provide details about how you controlled for confounding variables in your analyses.

Population characteristics

Describe the covariate-relevant population characteristics of the human research participants (e.g. age, genotypic information, past and current diagnosis and treatment categories). If you filled out the behavioural & social sciences study design questions and have nothing to add here, write "See above."

Recruitment

Describe how participants were recruited. Outline any potential self-selection bias or other biases that may be present and how these are likely to impact results.

Ethics oversight

Identify the organization(s) that approved the study protocol.

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

- ☒ Life sciences ☐ Behavioural & social sciences ☐ Ecological, evolutionary & environmental sciences

Life sciences study design

All studies must disclose on these points even when the disclosure is negative.

Sample size	Describe how sample size was determined, detailing any statistical methods used to predetermine sample size OR if no sample-size calculation was performed, describe how sample sizes were chosen and provide a rationale for why these sample sizes are sufficient.
Data exclusions	Describe any data exclusions. If no data were excluded from the analyses, state so OR if data were excluded, describe the exclusions and the rationale behind them, indicating whether exclusion criteria were pre-established.
Replication	Describe the measures taken to verify the reproducibility of the experimental findings. If all attempts at replication were successful, confirm this OR if there are any findings that were not replicated or cannot be reproduced, note this and describe why.
Randomization	Describe how samples/organisms/participants were allocated into experimental groups. If allocation was not random, describe how covariates were controlled OR if this is not relevant to your study, explain why.
Blinding	Describe whether the investigators were blinded to group allocation during data collection and/or analysis. If blinding was not possible, describe why OR explain why blinding was not relevant to your study.

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems		Methods	
n/a	Involved in the study	n/a	Involved in the study
☒	☐ Antibodies	☒	☐ ChIP-seq
☒	☐ Eukaryotic cell lines	☒	☐ Flow cytometry
☒	☐ Palaeontology and archaeology	☒	☐ MRI-based neuroimaging
☒	☐ Animals and other organisms		
☒	☐ Clinical data		
☒	☐ Dual use research of concern		
☒	☐ Plants		

Plants

Seed stocks	Report on the source of all seed stocks or other plant material used. If applicable, state the seed stock centre and catalogue number. If plant specimens were collected from the field, describe the collection location, date and sampling procedures.
Novel plant genotypes	Describe the methods by which all novel plant genotypes were produced. This includes those generated by transgenic approaches, gene editing, chemical/radiation-based mutagenesis and hybridization. For transgenic lines, describe the transformation method, the number of independent lines analyzed and the generation upon which experiments were performed. For gene-edited lines, describe the editor used, the endogenous sequence targeted for editing, the targeting guide RNA sequence (if applicable) and how the editor was applied.
Authentication	Describe any authentication procedures for each seed stock used or novel genotype generated. Describe any experiments used to assess the effect of a mutation and, where applicable, how potential secondary effects (e.g. second site T-DNA insertions, mosaicism, off-target gene editing) were examined.