



Diagnosis and management of neurogenic orthostatic hypotension

Woohee Ju¹, Dong In Sinn²

¹Department of Neurology, Seoul National University Hospital, Seoul National University College of Medicine, Seoul, Korea

²Department of Neurology, School of Medicine, Stanford University, Palo Alto, CA, USA

Received: June 13, 2023

Revised: October 17, 2023

Accepted: October 18, 2023

Correspondence to

Dong In Sinn

Department of Neurology, School of Medicine, Stanford University, 291 Campus Drive, Stanford, Palo Alto 94305 CA, USA

Tel: +1-650-723-6469

Fax: +1-650-725-0390

E-mail: dsinn@stanford.edu

ORCID

Dong In Sinn

<https://orcid.org/0000-0002-9807-8273>

Woohee Ju

<https://orcid.org/0000-0003-0522-1956>

Orthostatic hypotension is a sustained and pathological drop in blood pressure upon standing. Orthostatic hypotension can be due to non-neurogenic conditions or autonomic disorders. Impaired baroreflex-mediated vasoconstriction and insufficient release of norepinephrine play key roles in the pathophysiology of neurogenic orthostatic hypotension. Its common symptoms mainly related to inadequate cerebral blood flow include dizziness, lightheadedness, and syncope. It is crucial to differentiate neurogenic orthostatic hypotension from non-neurogenic orthostatic hypotension. For the management of neurogenic orthostatic hypotension, physicians should implement non-pharmacological methods and, if possible, reverse combined non-neurological conditions. Depending on severity of symptoms, pharmacological intervention may be tried after or with non-pharmacological methods. Its management should be individualized based on intensity of symptoms, comorbid conditions, drug side effects, and etiology. In this review, we discuss the definition, pathophysiology, clinical approach, and management of neurogenic orthostatic hypotension.

Key words: Orthostatic hypotension; Autonomic disorders; Norepinephrine

INTRODUCTION

Immediately after standing up, gravity moves 450-500 mL of blood downward in the intravascular space. With continued standing, there is a gradual shift of fluid from the intravascular to the interstitial space. Therefore, orthostatic stress (standing) causes a large fluid shift of approximately 500-1,000 mL, which results in increased blood pooling in the dependent parts of the body below the heart. The main areas of blood pooling are the intestine and the large muscle groups of the legs.¹⁻⁴ To keep adequate blood supply to the brain, the autonomic nervous system plays a key role by activating various compensatory mechanisms, such as vasoconstriction of blood vessels in the non-essential areas.

The arterial baroreflex is the main mechanism for autonomic blood pressure (BP) control. Mechanoreceptors in the carotid artery monitor changes in the stretch of the arterial walls and send afferent signals to the brainstem. After processing the signals, the brainstem sends out efferent signals to the main target organs changing heart rate (HR) and blood vessel tone accordingly.^{5,6} Furthermore, prolonged upright posture stimulates the renin-angiotensin-aldosterone system and vasopressin secretion regulating blood volume on a long-term basis and ultimately maintaining cerebral blood flow.^{6,7} Failure of these compensatory mechanisms can lead to orthostatic hypotension (OH). Inadequate cerebral perfusion causes dizziness, lightheadedness, vision changes and syncope/pre-syncope. In this review article, we will discuss abnormal hemodynamic responses to standing up including OH, its pathophysiology, symptoms and management.

ABNORMAL HEMODYNAMIC CHANGES UPON STANDING UP

There exist four types of abnormal drops in BP after standing up: classic OH, initial OH, delayed OH, and delayed BP recovery.^{3,8} Exaggerated postural tachycardia or postural orthostatic tachycardia syndrome is not discussed here because there is no pathological drop in BP by its definition.⁸

Classic orthostatic hypotension

Classic OH is defined as a sustained reduction of systolic blood pressure (SBP) by at least 20 mmHg or of diastolic blood pressure (DBP) by at least 10 mmHg within 3 minutes of standing or tilt table testing with at least 60 degrees of head-up tilt (Fig. 1A, B).^{1-3,6-8} In hypertensive patients (a supine SBP ≥ 150 mmHg or DBP ≥ 90 mmHg), a cut-off value of 30/15 mmHg fall is more appropriate to define OH, since BP fall after standing up largely depends on baseline BP (Fig. 1C).^{8,9}

Delayed orthostatic hypotension

When patients report symptoms consistent with orthostatic intolerance, orthostatic vital signs with 3 minutes of standing at clinics may not reveal any abnormalities. In such cases, delayed OH can be considered especially when there is latency of symptoms after standing up. Delayed OH is defined as

a sustained drop in BP after standing for 3 minutes or long (Fig. 1C).^{3,8-10} It is considered a milder form of sympathetic adrenergic failure,^{8,10-12} and its progression to OH has been associated with α -synucleinopathies, diabetes mellitus, more severe abnormalities on autonomic reflex test, and higher mortality rates.¹³ However, the frequency, severity, and type of orthostatic intolerance do not differ significantly between classic and delayed OH.^{11,14}

Initial orthostatic hypotension

Initial orthostatic hypotension (I-OH) is a 'transient' reduction of SBP by at least 40 mmHg or of DBP by at least 20 mmHg. The transient drop occurs in 15 seconds of standing and BP recovers in 30 seconds (Fig. 1D). It occurs more frequently in adolescents and young adults.^{8,15} Patients may experience transient symptoms such as lightheadedness, nausea, and blurred vision right after standing up or when making a few steps after standing.¹⁶ A mismatch between cardiac output and peripheral vascular resistance is the most likely mechanism.⁸ I-OH is usually a benign phenomenon. Its symptoms can disappear quickly by sitting or squatting. The usual blood pressure cuff on the arm or the wrist may not detect it. An active standing test with a continuous beat-to-beat BP measurement is required. Also, an active standing test has high specificity but low sensitivity for I-OH.

Delayed blood pressure recovery

Delayed BP recovery is characterized by an immediate and transient drop in BP when standing, followed by a gradual recovery. The recovery in SBP to <20 mmHg from the supine baseline occurs 30 seconds or longer after standing and within 3 minutes (Fig. 1E).^{3,10} In a large cohort study, patients on anti-hypertensive medications or with diabetic autonomic neuropathy often present transient postural symptoms of delayed BP recovery. It is associated with an increased risk of unexplained falls and fractures.¹⁷⁻²⁰

Clinicians should be aware that classic OH, delayed OH, and delayed BP recovery are not mutually exclusive conditions.²¹ Various factors such as hydration, food-intake, ambient temperature, medications, and time spent in sitting or flat positions before standing up can affect postural hemodynamic changes.




Fig. 1. Different changes in blood pressure (BP) and heart rate (HR) during tilt-table tests (A-C) or during active standing tests (D, E). Green lines indicate HR. Upper red lines indicate systolic BP (SBP) and lower lines reflect diastolic BP (DBP). In (D) and (E), only SBP is shown. Orthostatic hypotension (OH) is defined as a sustained drop in SBP by at least 20 mmHg or in DBP by at least 10 mmHg (A, B). An absolute HR increase of less than 15 beats per minute or a ratio of HR to SBP ($\Delta\text{OH}/\Delta\text{SBP}$) less than 0.5 can differentiate neurogenic OH (A) from non-neurogenic OH (B). Delayed OH is a sustained decline in BP occurring after 3 minutes of standing. When supine SBP is greater than 150 mmHg, 30/15 criteria may be more appropriate for OH or delayed OH instead of 20/10 (C). Initial OH is a 'transient' drop in SBP by at least 40 mmHg or in DBP by at least 20 mmHg during active standing (D). Delayed BP recovery is a transient decline in BP by 20/10 during active standing, but BP recovers 30 seconds or longer after standing up and within 3 minutes (E).

www.kci.go.kr

CLINICAL PRESENTATIONS

Symptoms of orthostatic hypotension

OH can be either symptomatic or asymptomatic. Its symptoms result from hypoperfusion to the brain and other organs above the heart. Common symptoms include light-headedness, dizziness, and falls with or without loss of consciousness.^{1,2,6-10} Visual symptoms can occur due to impaired perfusion to the eyes and/or the occipital lobes. Auditory symptoms such as hearing changes and tinnitus may occur because the cochleae are end-artery organs vulnerable to ischemia. Patients might report vague symptoms of fatigue, cognitive slowing, and generalized weakness that improve upon lying down.^{9,10,22} Pain due to ischemia in the trapezius and the adjacent muscles can occur over the suboccipital and paracervical areas making its distribution look like a coat-hanger (coat-hanger pain).^{9,10,23,24} Moreover, a study has also found that a 30.0% proportion of patients with OH report shortness of breath (platypnea). Its main mechanism is a ventilation-perfusion mismatch with hypoperfusion of the lung apices.^{9,25} In general, the most important factor that can determine if certain symptoms are due to OH is that they occur in upright positions, and improve quickly after sitting or lying down.

Blood pressure threshold for symptoms

In a study with Parkinson's disease (PD) with OH, OH symptoms were associated with standing mean arterial pressure less than 75 mmHg, regardless of degrees of BP drop.²⁶ The authors suggested pharmacological intervention for OH should be initiated for patients with standing mean arterial pressure below 75 mmHg. In studies with OH from various etiologies, there were no hemodynamic parameters that could predict symptoms consistently suggesting a weak correlation between symptoms and hemodynamic measurements.^{27,28}

CLINICAL APPROACH

In-clinic monitoring

At clinics, BP and HR are measured after at least 3-5 minutes of resting in a supine position, ideally followed by measurements at 1 and 3 minutes of standing.^{2,6} When this method

is not feasible at a clinic, criteria of 15/7 mmHg instead of 20/10 mmHg can be used for sit-to-stand measurements to diagnose OH with acceptable sensitivity and specificity.²⁹ However, it's unclear whether this criterion can be applied to patients with hypertension in a seated position.

At-home monitoring

Studies suggest that home BP monitoring by patients is a highly feasible and efficient screening method for OH and that it is comparable to measurements supervised by healthcare providers.^{30,31} To minimize the effects of drugs on BP, early morning before taking any medications is an ideal time. Patients can measure BP/HR after lying down and resting. After that, they can obtain standing BP/HR after 3 minutes of standing.⁹ To detect OH, an orthostatic BP/HR log for 2 weeks is encouraged. Patients may need to measure BP when they experience symptoms (if feasible) or when they change medications for OH.

Neurogenic vs. non-neurogenic orthostatic hypotension

Non-neurological diseases such as acute anemia, volume depletion, adrenal insufficiency, heart failure, thyroid disorders, end-stage renal disease with hemodialysis, medications, and severe gastrointestinal diseases can cause OH. It is crucial to differentiate neurogenic OH (N-OH) from non-neurogenic OH to find its cause and manage related symptoms. HR changes may be helpful for the differentiation. Absent or minimal responses of HR to OH usually indicate N-OH. An absolute increase of HR by less than 15 beats per minute implies a higher probability of N-OH. A relative HR change to SBP drop ($\Delta\text{HR}/\Delta\text{SBP}$ ratio) less than 0.5 has very high sensitivity and specificity for diagnosing N-OH, especially in autonomic neurodegenerative diseases (Fig. 1A, B).^{32,33} However, HR can be affected by cardiac medications, pacemakers, cardiac ablation, and aging. Autonomic neuropathies sparing cardiac sympathetic innervation may show robust HR response to OH. The presence of N-OH does not exclude non-neurogenic components of OH. Combined non-neurogenic components, particularly medications, can aggravate OH in patients with N-OH.

Etiologies of neurogenic orthostatic hypotension

Causes of N-OH can be divided into central and peripheral diseases. Among peripheral causes, diabetic mellitus is

the most common condition. OH can be easily observed in patients with diabetes.³⁴ N-OH can be seen in acute immune-mediated diseases such as autoimmune autonomic ganglionopathy, seronegative autoimmune autonomic neuropathy, and Guillain-Barré syndrome.³⁵ Hereditary neuropathies such as familial amyloidosis polyneuropathy³⁶ as well as acquired neuropathies due to B12 deficiency, Sjogren disease, systemic lupus erythematosus, thyroid disease, and human immunodeficiency virus may result in N-OH. As central causes, there are spinal cord diseases (e.g., spinal cord injury, syringomyelia, transverse myelitis)^{6,37} and neurodegenerative diseases. Autonomic neurodegenerative diseases, collectively called α -synucleinopathies, include PD, multiple system atrophy (MSA), dementia with Lewy bodies, and pure autonomic failure (PAF) sharing pathological findings of abnormal α -synuclein accumulation.^{6,10,38}

PATHOPHYSIOLOGICAL CONCEPTS

Arterial baroreflex and inadequate release in norepinephrine

The major pathophysiological mechanism of N-OH is an inadequate release of norepinephrine after standing up. Fluid shift and blood pooling in the dependent areas after standing up produce a cascade of hemodynamic changes, including decreased venous return, a reduction in stroke volume by up to 40.0%,³⁹ an overall decline in cardiac output, and eventually lower BP.^{4,6,10,40} The mechanoreceptors of the arterial baroreflex in the carotid sinus and the aortic arch sense changes in tension of the arterial walls secondary to changes in BP. They send afferent signals to the brainstem via the vagus and glossopharyngeal nerves.^{4,6,40,41} The signals are processed in the brainstem in coordination with the higher-order central autonomic network.^{40,42} The main effector organs of the baroreflex are the blood vessels and the heart. The rostral ventrolateral medulla neurons send efferent signals to the intermediolateral cell column in the spinal cord (T1-L2) where sympathetic preganglionic nerves exit through the ventral root and form synapses with postganglionic sympathetic adrenergic fibers.¹⁰ There is also a separate efferent pathway adjusting cardiovagal tone to the heart. The activated adrenergic fibers increase release of norepinephrine (NE), a key neurotransmitter for sympathetic

vasoconstriction, at the neurovascular junction. In healthy people, plasma NE levels are doubled after standing from supine levels. However, NE increase is blunted in patients with N-OH (Fig. 2).⁴³

Supine hypertension

Approximately half of the patients with N-OH experience supine hypertension.^{10,44} Also, 30.0% of patients with PD and MSA have mild to severe supine hypertension, regardless of their age, disease duration, or stage.⁴⁵ Supine hypertension is largely asymptomatic especially during daytime. But it can lead to pressure-induced natriuresis at night with subsequent nocturia and poor sleep quality. Due to the forced natriuresis, OH can get worse the following morning with more severe symptoms.^{44,46} Supine hypertension increases the risks of cardiovascular diseases and renal failure as long-term complications. Therefore, patients with N-OH should be regularly screened for supine hypertension, especially when starting a long-acting medication for N-OH or when reporting possibly related symptoms such as frequent awakening and nocturia.^{44,47} The long-term benefits of the management of supine hypertension are still under a matter of investigation. Its managements are discussed in detail on the scientific statement of the professional societies.⁴⁸

Denervation hypersensitivity

Chronic reduction of NE release may cause denervation hypersensitivity in the peripheral α 1-adrenergic receptors. Denervation hypersensitivity exaggerates responses of the receptors to its agonists.^{7,49} Denervation hypersensitivity is believed to contribute to supine hypertension and unpredictable drug responses in patients with N-OH.

MANAGEMENTS

The consensus statements for N-OH management recommend a stepwise and individualized approach. Experts suggest a 4-step treatment algorithm: a review of the patient's medications, non-pharmacologic measures, initiation of a single pharmacologic agent, and combination pharmacologic agents.⁹ When feasible, factors that can aggravate OH should be corrected. Medications such as diuretics, tricyclics, peripheral α 1-blockers, beta-blockers, or calcium




Fig. 2. Norepinephrine (NE) changes after standing up. In a supine/resting state, there is a baseline release of NE from the post-ganglionic adrenergic fibers. After standing up, signal traffic in the autonomic ganglia increases due to the enhanced release of acetylcholine from the pre-ganglionic sympathetic fibers. It activates the post-ganglionic fibers resulting in a surge in plasma NE level and vasoconstriction (A). In post-ganglionic degeneration in pure autonomic failure, the baseline NE level is decreased. Standing up does not bring up the NE level significantly (B). In pre-ganglionic degeneration in multiple system atrophy, the baseline level is normal or slightly lower than normal. But standing up does not increase the level like normal subjects (C).

channel blockers with vasodilating effects, and nitrates should be stopped or minimized.^{6,8-10} Anemia may worsen OH as well.^{9,50} Anemia reduces systemic vascular resistance through the induction of nitric oxide synthesis and low blood viscosity.⁵¹ If OH symptoms persist and affect daily activities after correction of the precipitating factors, non-pharmacological measures need to be considered. In milder cases, pharmacological agents can be used when non-pharmacological treatments are insufficient. But, in severe cases, both non-pharmacological and pharmacological measures may need to be started spontaneously.

Non-pharmacological managements

1. Physical counter-maneuver

Physical counter-maneuvers such as squatting, leg-crossing, thigh contraction, and bending at the waist can reduce venous pooling and increase total peripheral resistance quickly through mechanical compression of the vein.^{3,4,52-54} Cardiac output and BP typically increase within 3-5 seconds,ulti-

mately maintaining cerebral perfusion pressure and extending the duration of the upright position.⁵⁵

2. Salt supplementation

An increase in water and salt intake may expand the intravascular volume and reduce the severity of OH. With systemic autonomic failure, there is continued urinary sodium loss even with a low-salt diet⁵⁶ indicating impaired renal control of sodium balance. Excessive urinary sodium loss can result in urinary loss of water (natriuresis) contributing to a substantial decrease in intravascular volume in patients with autonomic failure.^{57,58} To increase fluid volume and improve orthostatic tolerance, it is recommended to have a high salt intake (6-10 grams per day) along with increased water consumption.^{9,49,59} However, the recommended amount should be individualized particularly in patients with combined heart failure, liver disease, severe peripheral edema, or chronic kidney disease. Long-term effects of a high-salt diet in patients with autonomic failure are not well-studied yet.⁵⁸

3. Water

To expand circulating fluid, water intake needs to increase up to 2-2.5 L of water per day.^{2,12,60} Gulping a big glass of water (about 500 mL) over less than 10 minutes raises plasma NE level and increases BP quickly with peak effects in 20-30 minutes.⁶¹ Patients may use this method as a rescue when they experience worsening symptoms or in the morning before their morning OH medications take effects. This pressor response lasts for 1 to 1.5 hours. It is triggered by low osmolality of water. The transient receptor potential vanilloid 4 channel receptors in the portal system sense low osmolality and trigger the osmo-pressor response. Therefore, the temperature of water seems irrelevant,^{62,63} although some patients occasionally report better responses to cold water. Young healthy people do not have such responses. Older people without autonomic failure may have similar responses to a lesser degree. Only patients with autonomic failure have a pronounced BP response to rapid water drinking.^{61,64}

4. Compressive garments

Compressive garments for OH include custom-fitted elastic stockings (e.g., waist-high, thigh-high) and abdominal binder.^{9,10,65} Considering significant blood pooling in the splanchnic and muscular circulation, waist-high compression garments are the most effective, followed by thigh-high compression stockings.⁶⁶ However, their compliance is poor, and they can be cumbersome to put on, for individuals with impaired motor coordination.⁹ In such cases, an abdominal binder is a good alternative.⁶⁷ Patients should be instructed to apply or readjust the abdominal binder in a supine position.

5. Lifestyle modification

Activities such as prolonged standing, alcohol consumption, hot showers, and saunas should be avoided as they augment vasodilation and venous pooling.^{9,10,68} Additionally, high-carbohydrate foods can lead to postprandial hypotension due to shift of blood into the splanchnic circulation.⁶⁹ Therefore, patients who experience worsening OH symptoms after a meal should try frequent and smaller meals. In severe cases, medications such as caffeine, acarbose, and octreotide can be considered. For example, acarbose, an anti-diabetic drug, inhibits α -glucosidase and delays the release of glucose from larger carbohydrates.⁴⁹

Pharmacological managements

If non-pharmacological treatments are not sufficient, pharmacological treatments can be used in combination.¹² There are two categories of medications: NE enhancers and replacers. In theory, the selection of OH drugs depends on the phenotypes of N-OH: preganglionic and postganglionic types. In post-ganglionic degeneration with limited sympathetic reserve as seen in PAF and PD (Fig. 2B), NE replacers such as midodrine and droxidopa are preferred. In pre-ganglionic degeneration with relatively intact peripheral sympathetic reserve as seen in MSA (Fig. 2C), NE enhancers such as pyridostigmine or NE reuptake inhibitors to potentiate endogenous NE release may be a better choice.⁴⁹ Residual sympathetic reserve can be assessed based on resting serum NE levels, degrees of HR increase to OH (cardiac sympathetic reserve), and sympathetic adrenergic response during autonomic function tests.^{6,49,70} Other factors including comorbidities, degrees of supine hypertension, drug side effects, and drug prices may affect choices of drugs.

1. Midodrine

Midodrine is a short-acting $\alpha 1$ adrenoreceptor agonist. Desglymidodrine, its active metabolite, constricts arterial and venous vessels, thus increasing BP in a dose-dependent pattern. The half-life of desglymidodrine is approximately 4 hours, and the peak plasma level is reached in 1 hour after oral administration.⁷¹ In a randomized controlled trial of 97 patients with OH, 10 mg of midodrine increased standing SBP by 22 mmHg compared with placebo.⁷² Other randomized trial (10 mg dose three times per day)⁷³ and double-blind, dose-response trial (2.5, 10, and 20 mg)⁷¹ showed benefits of midodrine in OH symptoms. The U.S. Food and Drug Administration granted accelerated approval in 1996 but requested a post-market trial.¹⁰ In a subsequent phase 4 cross-over randomization study with patients who had been on stable doses of midodrine for at least 3 months, standing time on a tilt-table test was significantly shorter with placebo, which demonstrated sustained effects of midodrine.⁷⁴ Due to concern about iatrogenic supine hypertension, its last dose of the day should be away from bedtime by 4 hours.⁵³ Its common side effects include scalp itching and piloerection.⁴⁹ Other potential side effects include hypertension, tingling sensation, and urinary retention. However, midodrine has minimal central side effects because it does

not cross the blood-brain barrier. Midodrine is the most used medication for OH due to its availability, minimal drug interaction, and quick onset with predictable effects. It can be used as needed for planned activities requiring prolonged standing.

2. Fludrocortisone

Fludrocortisone, which is a synthetic mineralocorticoid analog, potentiates $\alpha 1$ receptor activity and expands plasma volume through the urinary reabsorption of water and salt.¹⁰ The effects of volume expansion last for only a few weeks.^{3,9,10} It is believed that fludrocortisone makes $\alpha 1$ adrenergic receptors more sensitive to endogenous NE through changes in DNA transcription and protein translation. Its side effects include hypokalemia, edema, heart failure, supine hypertension, and headaches.^{2,75} In a cohort study of 129 patients with MSA, increased mortality was observed in those treated with midodrine, fludrocortisone, or a combination of both, regardless of disease severity.⁷⁶ Additionally, fludrocortisone users had higher rates of all-cause hospitalizations, particularly among congestive heart failure patients, compared to midodrine users.⁷⁷

3. Droxidopa

Droxidopa is an oral synthetic amino acid that provides exogenous NE, as it is converted to NE through decarboxylation. Since droxidopa is a NE replacer, it is more effective in patients with reduced peripheral sympathetic reserve. Low resting plasma NE levels (<219.5 pg/mL) reflecting post-ganglionic degeneration were associated with a greater pressor effect of droxidopa.⁷⁸ Its phase 3 studies showed significant improvement in OH-related symptoms in the first week. However, the positive effects did not last longer.⁷⁹⁻⁸¹ Droxidopa was granted accelerated approval in 2014, which requires post-market clinical trials to demonstrate its sustainability. Therefore, the rationale and design of the neurogenic orthostatic hypotension to assess sustained effects of droxidopa therapy study have been completed recently. Its results from the first phase of an open-label trial for 12 weeks have been published.⁸² Results of the double-blinded trial will follow.⁸³ The most common adverse effects of droxidopa were headache and dizziness.⁸¹

4. NE reuptake inhibitors: atomoxetine and ampreloxadine

When norepinephrine is released, approximately 90.0% of NE is taken back to the presynaptic nerves by NE reuptake transporters (NET) quickly. Atomoxetine and ampreloxadine inhibit the NET and enhance the actions of NE on the peripheral adrenergic receptors (NE enhancer). Thus, NET inhibitors are more effective in pre-ganglionic degeneration where peripheral sympathetic reserve and baseline release of NE are still relatively intact.¹⁰ Higher resting plasma NE levels are moderately correlated with the pressor response to atomoxetine.⁸⁴ A randomized controlled trial suggests that atomoxetine may serve as an alternative therapeutic option to midodrine in the treatment of orthostatic hypotension in autonomic failure.⁸⁵ Although ampreloxadine showed promising results in a phase 2 study,⁸⁶ a phase 3 trial in patients with α -synucleinopathies was terminated prematurely due to the lack of positive outcomes. A different phase 3 study of ampreloxadine only with MSA (NCT03750552) has been recently launched.

5. Pyridostigmine

Pyridostigmine is a reversible cholinesterase inhibitor that potentiates cholinergic neurotransmission in the autonomic ganglia, both sympathetic and parasympathetic.^{87,88} In a double-blind, randomized study of 58 patients with neurogenic OH, pyridostigmine alone or in combination with midodrine (5 mg) improved standing BP, particularly by increasing diastolic BP, without exacerbating supine hypertension.⁸⁹ However, another double-blind crossover study found that pyridostigmine was less effective than fludrocortisone in treating OH in Parkinson's disease.⁹⁰ For individuals with severe autonomic failure, although neither pyridostigmine nor atomoxetine improved BP compared with placebo, the combination of pyridostigmine (60 mg) and atomoxetine (18 mg) increased seated BP in a synergistic manner.⁹¹ Because of its modest efficacy and small clinical sample size, pyridostigmine is currently used as a second-line or adjunctive treatment option. Additionally, side effects such as nausea, diarrhea, diaphoresis, and hypersalivation can occur because of its nonselective effects on both nicotinic and muscarinic acetylcholine receptors.^{10,88} Further more comprehensive research involving older individuals is required to confirm its effectiveness and tolerability.

CONCLUSION

Patients with autonomic disorders may have N-OH. To find its specific etiology and determine appropriate management, it is important to identify symptoms of OH and differentiate N-OH from non-neurogenic OH. Symptoms of N-OH result from insufficient blood supply to organs, mainly the brain, secondary to low-standing BP. There should be relief after sitting or lying down. Heart rate changes during OH may help distinguish between N-OH and non-neurogenic OH. The main pathophysiological mechanism of N-OH is insufficient increase in NE release from the sympathetic adrenergic system to the blood vessels. The patterns of plasma NE levels differ depending on phenotypes of N-OH (pre-ganglionic vs. post-ganglionic degeneration). A stepwise approach is recommended to manage N-OH symptoms. It is important to consider combined medical conditions and correct potential contributors to OH. Before initiating pharmacological treatment, non-pharmacological approaches should be explored and tailored to the patient's situation.

Conflicts of Interest

None.

REFERENCES

- Robertson D. The pathophysiology and diagnosis of orthostatic hypotension. *Clin Auton Res* 2008;18:2-7.
- Shibao C, Lipsitz LA, Biaggioni I. Evaluation and treatment of orthostatic hypotension. *J Am Soc Hypertens* 2013;7:317-324.
- Wieling W, Kaufmann H, Claydon VE, van Wijnen VK, Harms MPM, Juraschek SP, et al. Diagnosis and treatment of orthostatic hypotension. *Lancet Neurol* 2022;21:735-746.
- Low PA, Singer W. Management of neurogenic orthostatic hypotension: an update. *Lancet Neurol* 2008;7:451-458.
- Lanfranchi PA, Somers VK. Arterial baroreflex function and cardiovascular variability: interactions and implications. *Am J Physiol Regul Integr Comp Physiol* 2002;283:R815-R826.
- Magkas N, Tsioufis C, Thomopoulos C, Dilaveris P, Georgiopoulos G, Sanidas E, et al. Orthostatic hypotension: from pathophysiology to clinical applications and therapeutic considerations. *J Clin Hypertens (Greenwich)* 2019;21:546-554.
- Kanjwal K, George A, Figueredo VM, Grubb BP. Orthostatic hypotension: definition, diagnosis and management. *J Cardiovasc Med (Hagerstown)* 2015;16:75-81.
- Freeman R, Wieling W, Axelrod FB, Benditt DG, Benarroch E, Biaggioni I, et al. Consensus statement on the definition of orthostatic hypotension, neurally mediated syncope and the postural tachycardia syndrome. *Clin Auton Res* 2011;21:69-72.
- Gibbons CH, Schmidt P, Biaggioni I, Frazier-Mills C, Freeman R, Isaacson S, et al. The recommendations of a consensus panel for the screening, diagnosis, and treatment of neurogenic orthostatic hypotension and associated supine hypertension. *J Neurol* 2017;264:1567-1582.
- Freeman R, Abuzinadah AR, Gibbons C, Jones P, Miglis MG, Sinn DI. Orthostatic hypotension: JACC state-of-the-art review. *J Am Coll Cardiol* 2018;72:1294-1309.
- Tzur I, Barchel D, Khateb Z, Swarka M, Izhakian S, Gorelik O. Delayed versus classic orthostatic hypotension: clinical and prognostic implications. *Blood Press* 2020;29:209-219.
- Brignole M, Moya A, de Lange FJ, Deharo JC, Elliott PM, Fanciulli A, et al. Practical instructions for the 2018 ESC guidelines for the diagnosis and management of syncope. *Eur Heart J* 2018;39:e43-e80.
- Gibbons CH, Freeman R. Clinical implications of delayed orthostatic hypotension: a 10-year follow-up study. *Neurology* 2015;85:1362-1367.
- Byun JI, Moon J, Kim DY, Shin H, Sunwoo JS, Lim JA, et al. Delayed orthostatic hypotension: severity of clinical symptoms and response to medical treatment. *Auton Neurosci* 2018;213:81-85.
- van Wijnen VK, Harms MP, Go-Schön IK, Westerhof BE, Krediet CT, Stewart J, et al. Initial orthostatic hypotension in teenagers and young adults. *Clin Auton Res* 2016;26:441-449.
- Wieling W, Krediet CT, van Dijk N, Linzer M, Tschakovsky ME. Initial orthostatic hypotension: review of a forgotten condition. *Clin Sci (Lond)* 2007;112:157-165.
- Vinik AI, Ziegler D. Diabetic cardiovascular autonomic neuropathy. *Circulation* 2007;115:387-397.
- Finucane C, O'Connell MD, Donoghue O, Richardson K, Savva GM, Kenny RA. Impaired orthostatic blood pressure recovery is associated with unexplained and injurious falls. *J Am Geriatr Soc* 2017;65:474-482.
- Doyle K, Lavan A, Kenny RA, Briggs R. Delayed blood pressure recovery after standing independently predicts fracture in community-dwelling older people. *J Am Med Dir Assoc* 2021;22:1235-1241.e1.
- Mol A, Slangen LRN, van Wezel RJA, Maier AB, Meskers CGM.

- Orthostatic blood pressure recovery associates with physical performance, frailty and number of falls in geriatric outpatients. *J Hypertens* 2021;39:101-106.
21. van Twist DJL, Mostard GJM, Sipers WMWH. Delayed recovery from initial orthostatic hypotension: an expression of frailty in the elderly. *Clin Auton Res* 2020;30:105-106.
 22. Centi J, Freeman R, Gibbons CH, Nearingard S, Canova AO, Cronin-Golomb A. Effects of orthostatic hypotension on cognition in Parkinson disease. *Neurology* 2017;88:17-24.
 23. Robertson D, Kincaid DW, Haile V, Robertson RM. The head and neck discomfort of autonomic failure: an unrecognized aetiology of headache. *Clin Auton Res* 1994;4:99-103.
 24. Bleasdale-Barr KM, Mathias CJ. Neck and other muscle pains in autonomic failure: their association with orthostatic hypotension. *J R Soc Med* 1998;91:355-359.
 25. Gibbons CH, Freeman R. Orthostatic dyspnea: a neglected symptom of orthostatic hypotension. *Clin Auton Res* 2005;15:40-44.
 26. Palma JA, Gomez-Esteban JC, Norcliffe-Kaufmann L, Martinez J, Tijero B, Berganzo K, et al. Orthostatic hypotension in Parkinson disease: how much you fall or how low you go? *Mov Disord* 2015;30:639-645.
 27. Freeman R, Illigens BMW, Lapusca R, Campagnolo M, Abuzinadah AR, Bonyhay I, et al. Symptom recognition is impaired in patients with orthostatic hypotension. *Hypertension* 2020;75:1325-1332.
 28. Arbogast SD, Alsheklee A, Hussain Z, McNeeley K, Chelmsky TC. Hypotension unawareness in profound orthostatic hypotension. *Am J Med* 2009;122:574-580.
 29. Shaw BH, Garland EM, Black BK, Paranjape SY, Shibao CA, Okamoto LE, et al. Optimal diagnostic thresholds for diagnosis of orthostatic hypotension with a 'sit-to-stand test'. *J Hypertens* 2017;35:1019-1025.
 30. Cremer A, Rousseau AI, Boulesteau R, Kuntz S, Tzourio C, Gosse P. Screening for orthostatic hypotension using home blood pressure measurements. *J Hypertens* 2019;37:923-927.
 31. Gibbon JR, Parry SW, Witham MD, Yarnall A, Frith J. Feasibility, reliability and safety of self-assessed orthostatic blood pressure at home. *Age Ageing* 2022;51:afac153.
 32. Norcliffe-Kaufmann L, Kaufmann H, Palma JA, Shibao CA, Biaggioni I, Peltier AC, et al. Orthostatic heart rate changes in patients with autonomic failure caused by neurodegenerative synucleinopathies. *Ann Neurol* 2018;83:522-531.
 33. Lamotte G, Sandroni P. Updates on the diagnosis and treatment of peripheral autonomic neuropathies. *Curr Neurol Neurosci Rep* 2022;22:823-837.
 34. Freeman R. Autonomic peripheral neuropathy. *Neurol Clin* 2007;25:277-301.
 35. Zaeem Z, Siddiqi ZA, Zochodne DW. Autonomic involvement in Guillain-Barré syndrome: an update. *Clin Auton Res* 2019;29:289-299.
 36. Palma JA, Gonzalez-Duarte A, Kaufmann H. Orthostatic hypotension in hereditary transthyretin amyloidosis: epidemiology, diagnosis and management. *Clin Auton Res* 2019;29:33-44.
 37. Goldstein DS, Sharabi Y. Neurogenic orthostatic hypotension: a pathophysiological approach. *Circulation* 2009;119:139-146.
 38. Mendoza-Velásquez JJ, Flores-Vázquez JF, Barrón-Velázquez E, Sosa-Ortiz AL, Illigens BW, Siepmann T. Autonomic dysfunction in α -synucleinopathies. *Front Neurol* 2019;10:363.
 39. Harms MP, Wesseling KH, Pott F, Jenstrup M, Van Goudoever J, Secher NH, et al. Continuous stroke volume monitoring by modelling flow from non-invasive measurement of arterial pressure in humans under orthostatic stress. *Clin Sci (Lond)* 1999;97:291-301.
 40. Idiaquez JF, Idiaquez J, Casar JC, Biaggioni I. Neurogenic orthostatic hypotension. Lessons from synucleinopathies. *Am J Hypertens* 2021;34:125-133.
 41. Smit AA, Halliwill JR, Low PA, Wieling W. Pathophysiological basis of orthostatic hypotension in autonomic failure. *J Physiol* 1999;519:1-10.
 42. Benarroch EE. The central autonomic network: functional organization, dysfunction, and perspective. *Mayo Clin Proc* 1993;68:988-1001.
 43. Cryer PE, Weiss S. Reduced plasma norepinephrine response to standing in autonomic dysfunction. *Arch Neurol* 1976;33:275-277.
 44. Mantovani G, Marozzi I, Rafanelli M, Rivasi G, Volpato S, Ungar A. Supine hypertension: a state of the art. *Auton Neurosci* 2022;241:102988.
 45. Fanciulli A, Göbel G, Ndayisaba JP, Granata R, Duerr S, Strano S, et al. Supine hypertension in Parkinson's disease and multiple system atrophy. *Clin Auton Res* 2016;26:97-105.
 46. Jordan J, Tank J, Heusser K, Reuter H, Biaggioni I. What do we really know about supine hypertension in patients with orthostatic hypotension. *Curr Opin Cardiol* 2019;34:384-389.
 47. Fanciulli A, Jordan J, Biaggioni I, Calandra-Buonaura G, Cheshire WP, Cortelli P, et al. Consensus statement on the definition of neurogenic supine hypertension in cardiovascular autonomic failure by the American Autonomic Society (AAS) and the European Federation of Autonomic Societies (EFAS) : endorsed by

- the European Academy of Neurology (EAN) and the European Society of Hypertension (ESH). *Clin Auton Res* 2018;28:355-362.
48. Jordan J, Fanciulli A, Tank J, Calandra-Buonaura G, Cheshire WP, Cortelli P, et al. Management of supine hypertension in patients with neurogenic orthostatic hypotension: scientific statement of the American Autonomic Society, European Federation of Autonomic Societies, and the European Society of Hypertension. *J Hypertens* 2019;37:1541-1546.
 49. Biaggioni I. Management of orthostatic hypotension. Introduction. In: Biaggioni I, Browning K, Fink G, Jordan J, Low PA, Paton JFR, eds. *Primer on the autonomic nervous system*. 4th ed. Vol. 1. Oxford (UK): Elsevier; 2023;699-704.
 50. Ando Y, Asahara K, Obayashi K, Suhr O, Yonemitsu M, Yamashita T, et al. Autonomic dysfunction and anemia in neurologic disorders. *J Auton Nerv Syst* 1996;61:145-148.
 51. Bolek H, Bolek EC. Orthostatic hypotension among elderly patients in Italian internal medicine wards: an observational study-comment. *Intern Emerg Med* 2020;15:1353-1354.
 52. Wieling W, van Dijk N, Thijs RD, de Lange FJ, Krediet CT, Halliwill JR. Physical countermeasures to increase orthostatic tolerance. *J Intern Med* 2015;277:69-82.
 53. Palma JA, Kaufmann H. Management of orthostatic hypotension. *Continuum (Minneapolis)* 2020;26:154-177.
 54. van Dijk N, de Bruin IG, Gisolf J, de Bruin-Bon HA, Linzer M, van Lieshout JJ, et al. Hemodynamic effects of leg crossing and skeletal muscle tensing during free standing in patients with vasovagal syncope. *J Appl Physiol* (1985) 2005;98:584-590.
 55. Krediet CT, de Bruin IG, Ganzeboom KS, Linzer M, van Lieshout JJ, Wieling W. Leg crossing, muscle tensing, squatting, and the crash position are effective against vasovagal reactions solely through increases in cardiac output. *J Appl Physiol* (1985) 2005;99:1697-1703.
 56. Wilcox CS, Aminoff MJ, Slater JD. Sodium homeostasis in patients with autonomic failure. *Clin Sci Mol Med* 1977;53:321-328.
 57. Williams EL, Raj SR, Schondorf R, Shen WK, Wieling W, Claydon VE. Salt supplementation in the management of orthostatic intolerance: vasovagal syncope and postural orthostatic tachycardia syndrome. *Auton Neurosci* 2022;237:102906.
 58. Stock JM, Chelimsky G, Edwards DG, Farquhar WB. Dietary sodium and health: how much is too much for those with orthostatic disorders? *Auton Neurosci* 2022;238:102947.
 59. Mtinangi BL, Hainsworth R. Early effects of oral salt on plasma volume, orthostatic tolerance, and baroreceptor sensitivity in patients with syncope. *Clin Auton Res* 1998;8:231-235.
 60. Snapper H, Cheshire WP. Oral and intravenous hydration in the treatment of orthostatic hypotension and postural tachycardia syndrome. *Auton Neurosci* 2022;238:102951.
 61. Jordan J, Shannon JR, Black BK, Ali Y, Farley M, Costa F, et al. The pressor response to water drinking in humans: a sympathetic reflex? *Circulation* 2000;101:504-509.
 62. Lipp A, Tank J, Franke G, Arnold G, Luft FC, Jordan J. Osmosensitive mechanisms contribute to the water drinking-induced pressor response in humans. *Neurology* 2005;65:905-907.
 63. McHugh J, Keller NR, Appalsamy M, Thomas SA, Raj SR, Diedrich A, et al. Portal osmopressor mechanism linked to transient receptor potential vanilloid 4 and blood pressure control. *Hypertension* 2010;55:1438-1443.
 64. May M, Jordan J. The osmopressor response to water drinking. *Am J Physiol Regul Integr Comp Physiol* 2011;300:R40-R46.
 65. Podoleanu C, Maggi R, Brignole M, Croci F, Incze A, Solano A, et al. Lower limb and abdominal compression bandages prevent progressive orthostatic hypotension in elderly persons: a randomized single-blind controlled study. *J Am Coll Cardiol* 2006;48:1425-1432.
 66. Ricci F, De Caterina R, Fedorowski A. Orthostatic hypotension: epidemiology, prognosis, and treatment. *J Am Coll Cardiol* 2015;66:848-860.
 67. Bourne KM, Sheldon RS, Hall J, Lloyd M, Kogut K, Sheikh N, et al. Compression garment reduces orthostatic tachycardia and symptoms in patients with postural orthostatic tachycardia syndrome. *J Am Coll Cardiol* 2021;77:285-296.
 68. Lahrmann H, Cortelli P, Hiltz M, Mathias CJ, Struhal W, Tassinari M. EFNS guidelines on the diagnosis and management of orthostatic hypotension. *Eur J Neurol* 2006;13:930-936.
 69. Figueroa JJ, Basford JR, Low PA. Preventing and treating orthostatic hypotension: as easy as A, B, C. *Cleve Clin J Med* 2010;77:298-306.
 70. Shibao C, Lipsitz LA, Biaggioni I. ASH position paper: evaluation and treatment of orthostatic hypotension. *J Clin Hypertens (Greenwich)* 2013;15:147-153.
 71. Wright RA, Kaufmann HC, Perera R, Opfer-Gehrking TL, McElligott MA, Sheng KN, et al. A double-blind, dose-response study of midodrine in neurogenic orthostatic hypotension. *Neurology* 1998;51:120-124.
 72. Jankovic J, Gilden JL, Hiner BC, Kaufmann H, Brown DC, Coghlan CH, et al. Neurogenic orthostatic hypotension: a double-blind, placebo-controlled study with midodrine. *Am J Med* 1993;95:38-48.

73. Low PA, Gilden JL, Freeman R, Sheng KN, McElligott MA. Efficacy of midodrine vs placebo in neurogenic orthostatic hypotension. A randomized, double-blind multicenter study. Midodrine Study Group. *JAMA* 1997;277:1046-1051.
74. Smith W, Wan H, Much D, Robinson AG, Martin P. Clinical benefit of midodrine hydrochloride in symptomatic orthostatic hypotension: a phase 4, double-blind, placebo-controlled, randomized, tilt-table study. *Clin Auton Res* 2016;26:269-277.
75. Ten Harkel AD, Van Lieshout JJ, Wieling W. Treatment of orthostatic hypotension with sleeping in the head-up tilt position, alone and in combination with fludrocortisone. *J Intern Med* 1992;232:139-145.
76. Pavy-Le Traon A, Foubert-Samier A, Ory-Magne F, Fabbri M, Senard JM, Meissner WG, et al. Ambulatory blood pressure and drug treatment for orthostatic hypotension as predictors of mortality in patients with multiple system atrophy. *Eur J Neurol* 2022;29:1025-1034.
77. Grijalva CG, Biaggioni I, Griffin MR, Shibao CA. Fludrocortisone is associated with a higher risk of all-cause hospitalizations compared with midodrine in patients with orthostatic hypotension. *J Am Heart Assoc* 2017;6:e006848.
78. Palma JA, Norcliffe-Kaufmann L, Martinez J, Kaufmann H. Supine plasma NE predicts the pressor response to droxidopa in neurogenic orthostatic hypotension. *Neurology* 2018;91:e1539-e1544.
79. Biaggioni I, Freeman R, Mathias CJ, Low P, Hewitt LA, Kaufmann H. Randomized withdrawal study of patients with symptomatic neurogenic orthostatic hypotension responsive to droxidopa. *Hypertension* 2015;65:101-107.
80. Kaufmann H, Freeman R, Biaggioni I, Low P, Pedder S, Hewitt LA, et al. Droxidopa for neurogenic orthostatic hypotension: a randomized, placebo-controlled, phase 3 trial. *Neurology* 2014;83:328-335.
81. Hauser RA, Isaacson S, Lisk JP, Hewitt LA, Rowse G. Droxidopa for the short-term treatment of symptomatic neurogenic orthostatic hypotension in Parkinson's disease (nOH306B). *Mov Disord* 2015;30:646-654.
82. Hauser RA, Favit A, Hewitt LA, Lindsten A, Gorny S, Kymes S, et al. Durability of the clinical benefit of droxidopa for neurogenic orthostatic hypotension during 12 weeks of open-label treatment. *Neurol Ther* 2022;11:459-469.
83. Owen R, Peng G, Gorny S, Cram C, Rowse G. Rationale and design of the neurogenic orthostatic hypotension to assess sustained effects of droxidopa therapy (RESTORE) study (P5. 131). *Neurology* 2016;86:131.
84. Shibao CA, Palma JA, Celedonio JE, Martinez J, Kaufmann H, Biaggioni I. Predictors of the pressor response to the norepinephrine transporter inhibitor, atomoxetine, in neurogenic orthostatic hypotension. *Hypertension* 2021;78:525-531.
85. Ramirez CE, Okamoto LE, Arnold AC, Gamboa A, Diedrich A, Choi L, et al. Efficacy of atomoxetine versus midodrine for the treatment of orthostatic hypotension in autonomic failure. *Hypertension* 2014;64:1235-1240.
86. Kaufmann H, Vickery R, Wang W, Kanodia J, Shibao CA, Norcliffe-Kaufmann L, et al. Safety and efficacy of amprelosetine in symptomatic neurogenic orthostatic hypotension: a phase 2 trial. *Clin Auton Res* 2021;31:699-711.
87. Palma JA, Kaufmann H. Epidemiology, diagnosis, and management of neurogenic orthostatic hypotension. *Mov Disord Clin Pract* 2017;4:298-308.
88. Lamotte G, Lenka A. Orthostatic hypotension in parkinson disease: what is new? *Neurol Clin Pract* 2022;12:e112-e115.
89. Singer W, Sandroni P, Opfer-Gehrking TL, Suarez GA, Klein CM, Hines S, et al. Pyridostigmine treatment trial in neurogenic orthostatic hypotension. *Arch Neurol* 2006;63:513-518.
90. Schreglmann SR, Büchele F, Sommerauer M, Epprecht L, Kägi G, Hägele-Link S, et al. Pyridostigmine bromide versus fludrocortisone in the treatment of orthostatic hypotension in Parkinson's disease - a randomized controlled trial. *Eur J Neurol* 2017;24:545-551.
91. Okamoto LE, Shibao CA, Gamboa A, Diedrich A, Raj SR, Black BK, et al. Synergistic pressor effect of atomoxetine and pyridostigmine in patients with neurogenic orthostatic hypotension. *Hypertension* 2019;73:235-241.